CORRECTION Open Access



Correction to: LINC00460/DHX9/IGF2BP2 complex promotes colorectal cancer proliferation and metastasis by mediating HMGA1 mRNA stability depending on m6A modification

Pingfu Hou^{1,2†}, Sen Meng^{1†}, Minle Li^{1†}, Tian Lin¹, Sufang Chu¹, Zhongwei Li¹, Junnian Zheng^{1,2*}, Yumina Gu^{3*} and Jin Bai^{1,2*}

Correction to: J Exp Clin Cancer Res 40, 52 (2021) https://doi.org/10.1186/s13046-021-01857-2

Following publication of the original article [1], the authors identified minor errors in Fig. 3, specifically:

• Fig. 3d: incorrect image was used for migration cells image of shLINC00460#2 (1st row, 3rd column)

The authors provided the Journal with the original data files. The corrected figure is provided here. The correction does not have any effect on the results or conclusions of the paper. The original article has been corrected.

Author details

¹Cancer Institute, Xuzhou Medical University, 84 West Huaihai Road, Xuzhou 221002, Jiangsu Province, China. ²Center of Clinical Oncology,

The original article can be found online at https://doi.org/10.1186/s13046-021-01857-2

Full list of author information is available at the end of the article

Affiliated Hospital of Xuzhou Medical University, Xuzhou, China. ³Department of Interventional Radiography, Affiliated Hospital of Xuzhou Medical University, 99 West Huaihai Road, Xuzhou 221002, Jiangsu Province, China.

Published online: 16 November 2021

Reference

 Hou P, Meng S, Li M, et al. LINC00460/DHX9/IGF2BP2 complex promotes colorectal cancer proliferation and metastasis by mediating HMGA1 mRNA stability depending on m6A modification. J Exp Clin Cancer Res. 2021;40:52. https://doi.org/10.1186/s13046-021-01857-2.



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativeccommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*}Correspondence: jnzheng@xzhmu.edu.cn; guyumingxyfy@foxmail.com; bj@xzhmu.edu.cn

[†]Pingfu Hou, Sen Meng and Minle Li contributed equally to this work.

² Center of Clinical Oncology, Affiliated Hospital of Xuzhou Medical University, Xuzhou, China

³ Department of Interventional Radiography, Affiliated Hospital of Xuzhou Medical University, 99 West Huaihai Road, Xuzhou 221002, Jiangsu Province, China




Fig. 3 LINC00460 promotes CRC cell migration, invasion and supports the proper function of the EMT transcription program and the location of LINC00460 in CRC cells. **a, b, c, d** The ability of migration and invasion in HCT116 and SW480 cells \pm LINC00460 KD/OE (***p < 0.001). **e, f, g, h** Relative mRNA expression levels of LINC00460 and EMT markers in HCT116 and SW480 cells \pm LINC00460 KD/OE. The relative mRNA expression levels were normalized to GAPDH (*p < 0.05, **p < 0.01, ***p < 0.001). **i, j** Western blots of EMT markers in HCT116 and SW620 cells \pm LINC00460 KD/OE. GAPDH was used as a loading control