


REVIEW

Open Access



Developing a diagnostic framework for patients presenting with Exercise Induced Leg Pain (EILP): a scoping review

Fatma Bosnina¹, Nat Padhiar^{1,2*} , Stuart Miller¹, Krishna Girotra¹, Chrysovalanto Massoura¹ and Dylan Morrissey^{1,3}

Abstract

Background Numerous conditions are grouped under the generic term exercise-induced leg pain (EILP), yet clear diagnostic guidelines are lacking. This scoping review was conducted to clarify the definition and diagnostic criteria of nine commonly occurring EILP conditions.

Methods Three online databases were searched from inception to April 2022 for any English language original manuscripts identifying, describing, or assessing the clinical presentation and diagnostic criteria of the nine most common conditions that cause EILP. We included manuscripts considering all adults with any reported diagnostic criteria for EILP in any setting. Methodological quality was assessed using the Mixed Method Appraisal tool. Condition definitions were identified and categorised during data charting. Twenty-five potential elements of the history, 24 symptoms, 41 physical signs, 21 investigative tools, and 26 overarching diagnostic criteria, were identified and coded as counts of recommendation per condition, alongside qualitative analysis of the clinical reasoning. Condition definitions were constructed with 11 standardised elements based on recent consensus exercises for other conditions.

Results One hundred nineteen retained manuscripts, of which 18 studied multiple conditions, had a median quality of 2/5. A combination of the history, pain location, symptoms, physical findings, and investigative modalities were fundamental to identify each sub-diagnosis alongside excluding differentials. The details differed markedly for each sub-diagnosis. Fifty-nine manuscripts included data on chronic exertional compartment syndrome (CECS) revealing exertional pain (83% history), dull aching pain (76% symptoms), absence of physical signs (78% physical findings) and elevated intercompartment pressure (93% investigative modality). Twenty-one manuscripts included data on medial tibial stress syndrome (MTSS), revealing persistent pain upon discontinuation of activity (81% history), diffuse medial tibial pain (100% pain location), dull ache (86% symptoms), diffuse tenderness (95% physical findings) and MRI for exclusion of differentials (62% investigative modality). Similar analyses were performed for stress fractures (SF, $n=31$), popliteal artery entrapment syndrome (PAES, $n=22$), superficial peroneal nerve entrapment syndrome (SPNES, $n=15$), lumbar radiculopathy ($n=7$), accessory/low-lying soleus muscle syndrome (ALLSMS, $n=5$), myofascial tears ($n=3$), and McArdle's syndrome ($n=2$).

Conclusion Initial diagnostic frameworks and definitions have been developed for each condition of the nine most common conditions that cause EILP, suitable for clinical consideration and consensus confirmation.

*Correspondence:

Nat Padhiar

nat.padhiar@london-sportswise.co.uk

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords Exercise induced leg pain, Chronic exertional compartment syndrome, Medial tibial stress syndrome, Tibial stress fractures, Popliteal artery entrapment syndrome, Superficial peroneal nerve entrapment syndrome, Lumbar-sacral radiculopathy, McArdle's syndrome, Myofascial tears and accessory/low-lying soleus muscle syndrome

Background

Exercise-induced leg pain (EILP) is an umbrella term to describe pain in the leg induced by physical activities. It is commonly seen in young active individuals involved in competitive or endurance sports (including running, football, rugby, and dancing). The incidence of EILP among athletes has been reported to be between 12.8% and 82.4% [1–4]; however, it has not been reported for the general population [1–3]. The common element of EILP is a history of activity-induced leg pain, which commonly increases until activity cessation [5, 6]. In some cases, pain is also experienced at rest and manageable during activity. Apart from pain, other symptoms can include burning, cramping, swelling, muscle weakness, fatigue, malaise, paraesthesia, numbness, poikilothermia, and tightness [2, 7, 8].

There are more than 40 conditions (Additional file 1) that fall into the generic category of EILP, which requires clinicians to be vigilant when diagnosing patients presenting with exercise-related leg pain [2, 4–6, 9, 10]. The most common EILP entities fall into one of the following subdivisions [5]: (a) muscular origin pain, such as chronic exertional compartment syndrome (CECS) and myofascial tears; (b) pain of bony origin, such as medial tibial stress syndrome (MTSS) and tibial stress fracture (TSF); (c) pain due to nerve entrapment or compression, such as superficial peroneal nerve entrapment syndrome (SPNES); (d) radicular pain or radiculopathy; (e) pain of vascular origin due to temporary compression of either an artery or a vein, such as popliteal artery entrapment syndrome (PAES); (f) muscle disorders, such as McArdle's syndrome; and (g) accessory or low-lying soleus muscle syndrome (ALLSMS) (also known as ASM) [2, 4–6, 9, 10]. These nine specific EILP conditions (CECS, MTSS, tibial SF, SPNES, myofascial tears, radiculopathy, PAES, McArdle syndrome, & ALLSMS) were the focus of this scoping review.

A detailed and methodical clinical history, knowledge of the conditions, appropriate examination and anatomical knowledge are regarded as key to reaching an appropriate diagnosis, while remaining aware that some of the conditions have overlapping signs and symptoms.

Because of coexisting conditions, a lack of consensus on the diagnostic guidelines and limited diagnostic test accuracy, EILP subdiagnoses cannot be clearly established in many cases [5, 10]. Clinical examination or physical findings are variably helpful, with the exception

of nerve- or bone-related pathology. Investigations or further investigative modalities are often essential to confirm the diagnosis [10]. It is essential to be able to differentiate between the causes of EILP among patients to reach the most appropriate diagnosis to construct an appropriate treatment and management plan [5, 10].

There is a copious and varied body of literature describing EILP-related conditions and their clinical presentation; however, there is limited guidance facilitating differentiation between even the more common conditions. This variation contributes to diagnostic confusion, variable care, and suboptimal outcomes. Therefore, the aim of this scoping review was to collate and evaluate the published diagnostic criteria of the nine most commonly occurring EILP conditions to provide clear guidance about differential diagnosis.

Following an introductory literature search [11] and discussions between the authors, the research question—‘What are the diagnostic criteria for patients with EILP?’—was formulated to reflect the scoping review population, concept, and context.

Methods

This scoping review was conducted and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews (PRISMA-ScR) [12]. The methodological framework of this review is based on Arksey and O'Malley's approach with modifications recommended by Levac et al., as well as the work of the Joanna Briggs Institute and previously published relevant scoping reviews [13–17]. The completed PRISMA-ScR Checklist is available (See Additional file 2) [12].

A scoping review was the most appropriate methodological approach to achieve this aim due to the iterative, exploratory methodology the topic needed [18–20]. A scoping review was also viewed as ideal to clarify key concepts, characterise findings from the diverse literature and outline the range of available evidence [11, 14, 15].

Protocol and registration

The protocol was registered as an open-ended registration with Open Science Framework (OSF) (<https://doi.org/10.17605/OSF.IO/4K27B>) [21]. The plan was modified after searching three databases (PubMed, Embase, Scopus), and the team decided not to search five databases, as we found three conclusive databases. Therefore,

the protocol was modified on August 2022 (<https://doi.org/10.17605/OSF.IO/DBPJE>) [22].

Eligibility criteria

We included manuscripts considering adult people of any gender that contained a description of EILP diagnostic criteria in any format at any consultation in all health care settings. However, the diagnostic criteria did not need to be the focus of the manuscript to be included, i.e., if the recruitment involved detail of how a condition was screened for, or if in the introduction of a systematic review it was described, then that paper was included. There was no limitation on manuscript type. See Table 1 for the full inclusion and exclusion criteria.

Information sources and search

Three online databases were searched from inception to April 2022 (i.e., PubMed, Embase, Scopus) for relevant manuscripts. The keywords and constructs used to execute each search systematically were determined based on a preliminary search [11, 23] of the existing relevant reviews [4, 5, 10, 24, 25] and in consultation with

the team members. A list of search terms (conditions, exercise, anatomical location, outcomes, and exclusions) is provided in Table 2, and the full search strategy for each database is available (see Additional file 3). A decision was made to limit searches to the English language, guided and managed by the first author, and organised using the reference management software EndNote 20.

Selection of sources of evidence

After deduplication using EndNote 20, two levels of screening were conducted. The titles and corresponding abstracts of all returned manuscripts were evaluated against the eligibility criteria using online Rayyan [26]. Second, full-text copies of all manuscripts included at the title and abstract screening stage were retrieved and screened to determine the initial study selection. Both stages were performed independently by three of the authors (FB, KG, CM), with each paper being screened by two authors. Any disagreement over the eligibility of a particular manuscript was resolved through discussion among the team to determine final manuscript selection meeting the eligibility criteria. If an agreement was not reached, a fourth author was consulted. Further searching of reference lists of the included full-text manuscripts and citations of these were explored to identify additional relevant manuscripts not returned in the primary search and to ensure that the search strategy was capturing relevant manuscripts [12–14].

Data charting

A customised coded Excel sheet was designed to collate the charted data in a consistent format (Additional file 4). Following an initial discussion and brief exploration of the returned manuscripts, the main headers

Table 1 Eligibility criteria

Inclusion criteria	Exclusion criteria
Human participants	Animal models
English language	Not English language
Adult participants (18–60)	Children or elderly participants
Description of clinical diagnosis of EILP	Systemic diseases causing leg pain
Manuscripts looking at diagnosis, assessment, and clinical presentation	Manuscripts containing only a description of therapeutic approaches

Table 2 Search terms

Search terms	
Construct	Keywords
Conditions	AND "shin splints" OR EILP OR "exercise induced leg pain" OR "compartment syndrome" OR "chronic exertional compartment syndrome" OR "exertional compartment syndrome" OR CECS OR "anterior compartment syndrome" OR "posterior compartment syndrome" OR "lateral compartment syndrome" OR "stress fracture" OR "tibial stress fracture" OR "fibular stress fracture" OR "stress syndrome" OR "Medial tibial stress syndrome" OR "tibial stress syndrome" OR "tibial stress injury" OR MTSS OR "stress injury" OR "radicular leg" OR "leg radiculopathy" OR "lumbar radiculopathy" OR "lumbar radicular" OR "nerve entrapment syndrome" OR "neuropathy" OR "peroneal neuropathy" OR "superficial peroneal neuropathy" OR "arterial entrapment syndrome" OR "popliteal artery entrapment syndrome" OR (exercise pain) OR "Myopathy" OR "Glycogen Storage Disease Type V" OR "McArdle disease" OR "McArdle syndrome"
Exercise	AND Exercise* OR active* OR sport* OR athlete* OR train* OR exertion OR practice* OR physical
Anatomical location	AND Leg OR compartment* OR calf OR shin* OR "lower limb" OR tibia* OR fibula* OR fascia
Outcomes	AND Diagnoses* OR "differential diagnosis" OR presentation OR "Clinical presentation" OR detection OR "clinical reasoning" OR findings
Exclusions	NOT In title: acute OR disease OR diabetes* OR foot OR feet OR thigh OR shoulder OR arm OR hand OR wrist OR back OR Raynaud's OR cervical radiculopathy

were chosen: i) definition of the condition, ii) anatomical location, iii) prevalence and incidence, iv) history, v) symptoms, vi) physical findings, vii) aggravating activity, viii) diagnostic criteria used within the literature, ix) investigative tools used, and x) therapeutic approaches undertaken and their efficacy. Within each of these main headings, subheadings were iteratively developed to classify the approaches described in each manuscript. Details of the population demographics, such as age and sex, were included wherever reported, alongside year of publication, author name, and study design [5, 6]. Three authors charted data from the first 10 manuscripts independently to determine whether their approach to the charting process was consistent [13, 27]. The full data charting process was conducted by one author (FB) for consistency, while two other authors (KG and CM) undertook approximately 50% each, so that data from each paper were checked by at least two reviewers.

Quality assessment

The methodological quality of retained manuscripts was evaluated by three authors (FB, KG and CM) using the Mixed Methods Appraisal Tool (MMAT) 2018 Version [28]. The MMAT measures five various methodological classifications, enabling use across different study designs, and is valid, reliable, and efficient [28, 29]. Inconsistencies in the MMAT scoring between the authors were resolved by agreement between them and accessing another author as needed.

Results synthesis

Condition definitions

In addition to the synthesis of reported diagnostic approaches, definitions for the nine conditions were recorded from each included manuscript, and a single definition for each was formulated by first identifying the most common definitions and their sources. The literature was searched to construct operational definitions to provide in the review. The content was then cross-referenced to a standardised structure derived from analysis of all definitions and consideration of previous reports of consensus definitions for chronic conditions [30]. The domains included [condition name], causing [selection of symptoms/impairments including frequency and severity], for at least [minimum duration], at [location], resulting in [disability], in [population], for [x duration], when [exercise relationship], because [proposed mechanism], confirmed by [y and z test(s)], and [other conditions] particularly need to be excluded.

Data coding and synthesis of main domains

Based on discussions among the study team and reviewing relevant conceptual and methodological articles [15–17, 31, 32], data were coded.

The five main domains identified initially within the data charting process were history, symptoms, physical findings, investigations and overarching diagnostic criteria. The number of manuscripts that reported the use of an approach within the diagnosis was coded and counted under each subdomain element. Synthesis of reporting across manuscripts was undertaken for each of the nine EILP-related conditions and is presented as the percentage of manuscripts reporting a diagnosis approach within the number of manuscripts reporting a diagnosis for the specific EILP diagnosis. As such, the total number of manuscripts differed across injuries, and a single manuscript might include data on multiple conditions.

The synthesised data were presented in condition-specific graphs, which show the proportion of manuscripts reporting a given diagnostic criterion. Data were combined for the nine conditions into a separate figure for each of the five main domains charted (all details are available in Additional file 4).

Results

Selection of sources of evidence

The search yielded 5022 manuscripts after deduplication. After the title and abstract screening stage, 303 manuscripts were reviewed in full. Of these 303 manuscripts, 119 were retained, and 184 were excluded because they did not meet the eligibility criteria as outlined in Fig. 1. Eleven of the full text articles were not available through online referral sources, and the authors did not respond to requests to share the full text.

Characteristics of sources of evidence

A summary of the types of manuscripts included in this review is shown in Table 3. Noncomparative studies including opinion pieces/narrative reviews accounted for 45.4% of the included manuscripts [4, 6, 33–84], case reports (22%) [7, 85–109], and case series studies (17%) [110–129]. Five percent of the included manuscripts were systematic reviews [5, 130–134]. Case control studies accounted for 2.5% [135–137], as well as retrospective case series studies (2.5%) [24, 138, 139]. Cross-sectional studies accounted for 2.5% [140–142]. Experimental quantitative trials accounted for 2.5% (randomised and nonrandomised controlled trials) [143–145]. In all, 0.8% of the manuscripts were observational cohort studies [146].




Fig. 1 Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) flow chart

Table 3 Type of manuscripts included and evidence level of different study types

Type of manuscripts	Number	Evidence level
Opinion piece/narrative review	54 (45.4%)	5
Case reports	26 (22%)	5
Case series	20 (17%)	4
Systematic reviews	6 (5%)	2
Case-control studies	3 (2.5%)	4
Retrospective case series	3 (2.5%)	4
Cross-sectional	3 (2.5%)	4
Randomised controlled trials (RCT)	2 (1.7%)	2
Nonrandomised controlled trials (non-RCT)	1 (0.8%)	3
Cohort studies	1 (0.8%)	2
	119 total	Median 4

Quality assessment

The median quality of the included manuscripts was 0.4, with a mean of 2.01 (IQR=3/0-3) using the MMAT (for details, see Additional file 5). Most of the nonsystematic reviews and opinion pieces/narrative reviews were rated as poor to moderate quality (0-1 scores), while case-series studies, case control studies, and experimental quantitative trials had higher MMAT scores (3-4 scores).

Data extraction

Additional file 4 presents the completed data charting table and the individual charted data from each resource of evidence.

Synthesis of results

Condition definitions

Synthesised definitions of the nine most common conditions causing EILP are provided in Table 4. Not all included manuscripts provide a specific definition of the involved condition, with 101 manuscripts describing only one condition, while 18 described more than one condition. Additionally, none of the included manuscripts defined myofascial tears, and the definition was synthesised from other literature resources. The purpose of this work was to produce definitions based on the current literature to represent current thinking but in a format suitable to inform later consensus work on the definitions.

Synthesis of main domains and diagnostic criteria

The results synthesis displays the frequency of each diagnostic element within the main domains for all conditions individually. The elements within the domains with the highest frequency for a certain condition are the ones reported the most within the included manuscripts; lower frequencies indicate that the element is less frequently reported within the manuscripts for diagnosing that condition.

Twenty-five potential subdomains/elements for history, 24 for symptoms, 41 for physical signs, and 21 for investigative tools were identified within the retained manuscripts. The fifth main domain identified was the overarching diagnostic criteria, some of which reproduced data in the other four main domains but had been highlighted by manuscript authors.

History domain The most reported elements of the medical history indicating *CECS* were exertional pain, gradual onset during the exercise, and the relief of pain within a few minutes after discontinuing the activity (frequency = 83%, 80%, and 76% of papers, respectively), as shown in Fig. 2. Pain relief soon after activity cessation appeared to be specific to *CECS* cases, with it only minimally being reported as a diagnostic tool for *PAES*, *SPNES*, *tibial SF* and *MTSS* (23%, 13%, 3%, and 5%, respectively). The most reported history elements for *MTSS* were diffuse pain spread within the distal two-thirds of the tibia, pain persistence from a few hours to days, and gradual symptom onset during activity (100%, 81%, 76%, respectively). The first two elements were specific to the *MTSS* presentation pattern. However, for *tibial stress fracture*, local pain spread (<5 cm), gradual onset of pain during exercise, and exertional pain were reported to be the history pattern in 74%, 55%, and 48% of patients, respectively. Localised pain was specific to *tibial stress fractures* (74%).

There were no history elements unique for *SPNES* and *radiculopathy*, as the main reported elements for *SPNES* were exertional pain, no medical history related, and gradual pain onset (60%, 47%, 40%, respectively), while the most common elements for *radiculopathy* were no prior medical history, nonexertional pain, and pain during rest (57%, 28.6%, 28.6%, respectively). For *PAES*, the typical elements were exertional pain type, absence of related medical history, and gradual onset during activity (77%, 50%, and 46%, respectively). The most frequent history elements for *myofascial tears* were direct/indirect trauma, nonexertional pain, and pain at rest (100%, 67%, 33%, respectively), with the history of trauma being particular to this condition. Cases of *McArdle's syndrome* were found to have a well-defined pattern; however, the elements were not particular to this syndrome, including exercise intolerance, exertional pain and pain relieved by rest (100%, 100%, and 50%, respectively). *ALLSMS* history was unclear, with the reported elements of swelling after exercising, pain being relieved by rest and exertional pain ranging from 20%-60%.

Symptoms domain The most reported symptoms are shown in Fig. 3, with a dull ache, tightness, and paraesthesia being the main symptoms for *CECS* (76%, 54%, and 44% respectively). A dull ache was also the main symptom reported for *MTSS* and *tibial stress fracture* (86% and 77%, respectively). However, *tibial stress fracture* also presented with sharp pain in some (23%) patients. Neurological symptoms, including numbness, tingling, and weakness (67%, 40%, 40% respectively), were the most common indications for *SPNES*. *Myofascial tears* can present with different types of pain (dull aching 67%, radiating 67%, and sharp 33%), while other commonly reported symptoms include altered sensation and tenderness after exercising (both 33%). The presence of low back pain (LBP), sharp pain, and radiating pain could be an indicator of *radiculopathy*, all of which returned 100%. The highly reported symptoms for *PAES* were dull ache, intermittent claudication, and paraesthesia (59%, 50%, 41%, respectively). LBP and intermittent claudication (100% and 50%, respectively) were specific symptoms for *radiculopathy* and *PAES*, respectively. The most specific reported symptoms for *McArdle's syndrome* were early fatigue, myalgia, and weakness (100%, 50%, and 50%, respectively). *ALLSMS* reports included dull aching, burning, and tenderness (40%, 40%, 20%, respectively); however, it could also present with no symptoms (20%).

Findings of the physical examination procedures domain Physical examination for patients with EILP was typically categorised as either observation, palpation, range of motion (ROM), neurological testing, or special

Table 4 Definitions of the nine most common EILP conditions

Condition	Synthesised definition and key early sources
Chronic exertional compartment syndrome	<p>... is defined as increased pressure in the lower leg compartment causing severe pain, tightness, and paraesthesia which in severe cases that would occur every time sufferers exercise for at least 6–12 months. The anterior compartment is most affected and symptoms usually bilateral, affecting the ability to do competitive sports. It is more common in young males. Symptoms may persist for months to years and usually build during activity until they must stop, pain then settle with rest, recurring on return to activity. The mechanism is tightness in the affected compartment due to an increase in intramuscular pressure, resulting in a lack of perfusion and altered physiology. CECS diagnosis is confirmed by measurement of dynamic intracompartment pressure. SPNES and MTSS need to be excluded.</p> <p>22 definitions reported in 59 (37%) relevant publications 2 common partial definitions, first reported in [147, 148]</p>
Medial tibial stress syndrome	<p>... is defined as periostitis causing diffuse pain for months to years at the distal posteromedial border of the tibia, which can be unilateral or bilateral, and lead to compromised ability to exercise. It is more common in young active individuals involved in endurance activity (runners, dancers, and military recruits), and can last for a few hours to days per episode. The mechanism is subcutaneous periostitis associated with altered biomechanics of the lower limb, training errors and increased training intensity. MTSS diagnosis is confirmed by clinical history, site of pain, palpation, MRI scan and bone scan. Stress fracture and CECS of the deep posterior compartment need to be excluded.</p> <p>11 definitions reported in 21 (52%) relevant publications 2 common definitions, first reported in [132, 146]</p>
Tibial stress fracture	<p>... is defined as cortical or full bone fracture causing localised excruciating pain both at rest and with weight-bearing activity with nocturnal pain for days to few weeks. This occurs on a daily basis until it heals, typically at the middle to lower one-third of the tibia most commonly but can occur anywhere. The acute pain can cause gross disability and usually affect young age group 10–30 years of age and it heals in a minimum of 6–8 weeks with immobilisation. The mechanism is an imbalance between osteoblast and osteoclast activity leading to bone breakage due to overtraining, overuse, and repeated overloading. Tibial stress fracture diagnosis is confirmed by localised tenderness, plus positive hop and/or fulcrum test. Bone tumours and frank fracture need to be excluded.</p> <p>10 definitions reported in 31 (32%) relevant publications 2 common definitions, first reported in [48, 149]</p>
Superficial peroneal nerve entrapment syndrome	<p>... is defined as mechanical compression of the superficial peroneal nerve causing moderate to severe pain, paraesthesia, numbness, and the feeling of a 'restless leg' for few months to years at unilateral anterior compartment of the leg. It is more common in young active adults and each episode can last for minutes to an hour. The symptoms occur with activity and are relieved with a variable period of rest. The mechanism is scarring/entrapment around the opening of the fascia where the nerve becomes a cutaneous sensory nerve supplying the dorsum of the foot. SPNES diagnosis is confirmed by diagnostic local anaesthetic test. CECS needs to be excluded.</p> <p>3 definitions reported in 15 (20%) relevant publications 2 common definitions, first reported in [150, 151]</p>
Myofascial tear	<p>... is defined as a single event causing a tear within the myofascial causing pain with activity as the main symptom and can be severely disabling. Located at the interface between the aponeurosis and the fibre or muscle fasciculus (and its corresponding perimysium).</p> <p>It is more common in young active adults and can be acute or chronic. In chronic cases, pain occurs with activity and is relieved with rest. The mechanism is the tendon or aponeurosis affected either focally by small muscle fibre tears or by major tears that produce a muscle gap but not a tendinous gap. Myofascial tears diagnosis is confirmed by dynamic ultrasound or, more reliably for deep tears, MRI. Muscle tears, haematoma, radiculopathy, and sural nerve entrapment syndrome need to be excluded.</p> <p>0 definition reported in 3 (0%) relevant publications 1 common definition, first reported in [152, 153]</p>

Table 4 (continued)

Condition	Synthesised definition and key early sources
Lumbar radiculopathy	<p>... is defined as mechanical compression of a nerve root at the level of the spinal cord as it exits the foramen or lateral recess causing sharp pain ... radiates down the legs, paraesthesia, numbness, spontaneous cramp, lack of power, fatigue, and tiredness in one or both legs during activity and at rest for months to years. It usually affects the posterior dermatomes and myotomes of the legs but can occur in other areas leading to restricted movement, disturbed sleep, and altered ability to exercise. It is more common in males 30–50 years old. The mechanism is mechanical compression of a nerve root at the level of the spinal cord as it exits the foramen or lateral recess. Lumbar radiculopathy diagnosis is confirmed by lumbar-sacral MRI scan and sometimes EMG/nerve conduction study. Gluteal and piriformis syndromes, myopathy, and in some cases unusual and uncommon CECS and PAES need to be excluded.</p> <p>3 definitions reported in 7 (42%) relevant publications 1 common definition, first reported in [154]</p>
Popliteal artery entrapment syndrome	<p>... is defined as arterial insufficiency in the affected limb which arises with entrapment of the artery, commonly giving leg symptoms with exertion causes pain, poikilothermia along with tightness, paraesthesia, and numbness can also occur for weeks to three months at the superficial posterior compartment, and it is usually unilateral, leading to intermittent claudication and temperature changes. It is more common in young active runners and each episode can last for few minutes. The mechanism is an abnormal relationship between the popliteal artery and the surrounding myofascial structures in the popliteal fossa. PAES diagnosis is confirmed by MRI, MRA, CT, Angiography, Duplex ultrasound scan. Radiculopathy, soleus sling syndrome, and CECS affecting the superficial posterior compartment need to be excluded.</p> <p>11 definitions reported in 22 (50%) relevant publications 1 common definitions, first reported in [80]</p>
McArdle's syndrome	<p>... is defined as autosomal recessive metabolic myopathy causes pain, tightness, swelling, malaise, and lethargy for years at multiple muscle compartments of the upper and lower limbs, leading to fixed weakness, malaise, fatigue, and tiredness. It is more common in young active population; it is a long-term condition that occurs during exercise. The mechanism is an autosomal recessive metabolic myopathy causing exercise-induced rhabdomyolysis due to a deficiency of muscle phosphorylase. McArdle's syndrome diagnosis is confirmed by resting creatine kinase (CK) followed by 3 consecutive postexercise CK and muscle biopsy. Other medical myopathies and CECS need to be excluded.</p> <p>1 definition reported in 2 (50%) relevant publications 1 common definition, first reported in [57]</p>
Accessory/low-lying soleus muscle syndrome	<p>... is defined as a space occupying mass which can cause nerve compression which mimics tarsal tunnel syndrome and can also cause increase in the intercompartmental pressure mimicking CECS causes soft tissue swelling that may be painful during physical activity for few months to years. Other symptoms that may be attributed to impingement on neurovascular structures and paraesthesia and numbness affecting the plantar aspect of the foot. It occurs at the superficial posterior compartment and can be unilateral or bilateral and more common in young active adults.</p> <p>The mechanism is a rare supernumerary anatomical variant. ALLSMS diagnosis is confirmed by MRI and ultrasound scans. CECS affecting the superficial posterior compartment, PAES, and radiculopathy need to be excluded.</p> <p>1 definition reported in 5 (20%) relevant publications 2 common definitions, first reported in [155, 156]</p>

tests. Therefore, Fig. 4 is presented to illustrate the most common physical findings used to establish a diagnosis. Interestingly, the absence of physical findings at rest was most reported for CECS (78%), followed by altered sensation and tenderness after exercising (46% and 41%, respectively). Diffuse tenderness on palpation presented in more than 95% of MTSS literature but absent in all others was specific for MTSS. Other findings, such as weakness and swelling, were reported with a lower frequency for MTSS (both 14.3%). The presence of local tenderness, a positive tuning fork and a hopping test were reported as being indicative of a *tibial stress fracture* (77%, 65%,

and 48%, respectively). The main reported physical findings for SPNES were weakness, positive Tinel's test, and altered sensation (67%, 53%, and 47%, respectively). The most reported findings for *radiculopathy* were weakness, positive slump test, femoral stretch, and straight leg raise, all of which returned 100%. Signs of swelling, tenderness after exercise and tense muscle compartment (67%) were the most common findings of *myofascial tears*. The most reported findings for PAES were reduced pulses, positional loss of pulses using arterial Doppler ultrasonography, and the absence of signs while resting (68%, 59%, 50%, respectively). Weakness was the only physical




Fig. 2 This bar chart details the frequency with which each of the 25 most common elements of the patient's history were reported as being relevant to each condition. Column headers indicate the condition and the number of relevant manuscripts in brackets. CECS=chronic exertional compartment syndrome, MTSS=medial tibial stress syndrome, tibial SF= tibial stress fracture, SPNES=superficial peroneal nerve entrapment syndrome, PAES=popliteal artery entrapment syndrome, ALLSMS=accessory/low-lying soleus muscle syndrome. Px=Pain. Hx=History




Fig. 3 This bar chart details the frequency with which each of the 24 most common elements of the patient's symptoms were reported as being relevant to each condition. Column headers indicate the condition and the number of relevant manuscripts in brackets. CECS=chronic exertional compartment syndrome, MTSS=medial tibial stress syndrome, tibial SF= tibial stress fracture, SPNES=superficial peroneal nerve entrapment syndrome, PAES=popliteal artery entrapment syndrome, ALLSMS=accessory/low-lying soleus muscle syndrome. Px=Pain. LBP=low back pain




Fig. 4 This bar chart details the frequency with which each of the 41 most common elements of the patient's physical findings were reported as being relevant to each condition. Column headers indicate the condition and the number of relevant manuscripts in brackets. CECS = chronic exertional compartment syndrome, MTSS = medial tibial stress syndrome, tibial SF = tibial stress fracture, SPNES = superficial peroneal nerve entrapment syndrome, PAES = popliteal artery entrapment syndrome, ALLSMS = accessory/low-lying soleus muscle syndrome. Px = Pain. ROM = Range of motion

finding reported for *McArdle's syndrome*, as well as the absence of signs during rest (100% and 50%, respectively). The appearance of a soft tissue mass in the ankle region, alongside tenderness after exercising, and pain during ankle plantar flexion (60%, 60%, 20%, respectively) were the main findings indicating *ALLSMS*.

Further investigations domain Further investigative modalities, as displayed in Fig. 5, were used chiefly as the concluding step in the diagnostic process as positive or negative confirmation of the clinical diagnosis. MRI was frequently used for all conditions to confirm a particular diagnosis (*ALLSMS* 80%, *tibial SF* 78%, *radiculopathy* 71%, *myofascial tears* 66%, *PAES* 50%, *SPNES* 20%), or to rule out other pathologies (*MTSS* 62%, *CECS* 31%), except for the absence of its use as a diagnostic tool for *McArdle's*. Conversely, *McArdle's* diagnosis was reported to be confirmed by more invasive tastings, i.e., genetic screening, blood test, and biopsy (100%, 50%, and 50%, respectively). The use of dynamic intracompartment pressure (ICP) tests returned a 66% frequency for *CECS*, with low to no returns for the others. Further measurement of the ankle brachial pressure index (ABPI) was only used to confirm cases of *PAES* (31%) alongside the

use of some imaging modalities (ultrasonography 68%, MRI 50%, CT angiogram 50%) after provoking symptoms with exercises and manoeuvres as determined by the history. The diagnostic process of *stress fractures* has been reported to be based on the use of imaging modalities, such as X-ray, MRI, and bone scan (87%, 87%, 54%, respectively). In addition, X-ray can be a useful indication of the presence of *ALLSMS* (20%), and ultrasound scans can reveal *myofascial tears* (66%). Moreover, nerve conduction studies could assist in the diagnosis of *radiculopathy* and *SPNES* (71% and 53%, respectively).

Diagnostic criteria

The overall diagnostic criteria for these nine conditions are presented in Fig. 6, which represents the frequency of each of the 26 most common diagnostic criteria reported for each of the conditions, including those reported in the previous four domains. In other words, the resultant overarching diagnostic criteria for these nine conditions consist of 24 criteria reported in the included manuscripts as individual criteria plus the other four domain elements (history, symptoms, physical findings, further investigations).




Fig. 5 This bar chart details the frequency with which each of the 21 most common elements of the further investigation's modalities were reported as being relevant to each condition. Column headers indicate the condition and the number of relevant manuscripts in brackets. CECS = chronic exertional compartment syndrome, MTSS = medial tibial stress syndrome, tibial SF = tibial stress fracture, SPNES = superficial peroneal nerve entrapment syndrome, PAES = popliteal artery entrapment syndrome, ALLSMS = accessory/low-lying soleus muscle syndrome. CT = computed tomography scan. USS = Ultrasound scan. MRI = Magnetic resonance imaging. ABPI = Ankle brachial pressure index. ICP = Dynamic intracompartment pressure

Discussion

This comprehensive scoping review details the definitions and key diagnostic criteria for the nine most common conditions that cause EILP based on the literature. The findings are collated and evaluated by condition.

Single definitions for each of the nine conditions were synthesised by identifying the most common definitions and their sources within the literature and then checked according to a standardised domain. These are qualitative outputs of the data synthesis, using criteria from the clinical case definition of post-COVID-19 syndrome in a previous Delphi consensus [30]. A Delphi study of health-care professionals and patients was conducted in a three-stage process to define core health-related domains for tendinopathy, and they found that nine core domains for tendon research should guide the reporting of outcomes in clinical trials [157, 158]. This imposes the importance of having clear definitions domains for such clinical conditions. The diversity in definition between manuscripts was notable, with some manuscripts not providing any definition and most lacking important details (including frequency, least, duration, mechanism) that would enable comparison between studies and evidence synthesis. By combining all of these in a standardised structure, we aim to provide a clear foundation for promoting discussions

in the future and use of these synthesised definitions for further consensus processes.

The diagnosis of CECS revolved primarily around the pain pattern of the relief within a few minutes after discontinuing the activity, the absence of physical findings when not exercising and invasive ICP measuring procedure. Elevated ICP is proposed to be a diagnostic tool for CECS within the results. This was accepted by Pedowitz et al., who demonstrated elevated values in the affected compartments and a resting pressure greater than 15 mmHg and 5 min postexercise pressures greater than 20 mmHg [139]. However, current studies outlining these criteria have no high-level evidence. Therefore, current ICP pressure criteria for CECS diagnosis are consequently unreliable, and emphasis should remain on a clear history [81, 159], which was also presented within this review [81, 159].

The diagnosis of MTSS was based mainly on a history of persistent, cumulative pain that does not immediately cease after stopping activity and may persist between a few hours and a few days, as reported in 81% of the papers. This is accompanied by diffuse tenderness of the tibial periosteum during physical examination (95% reporting frequency). Imaging, mainly MRI, was also used, albeit recommended in only 62% of manuscripts to




Fig. 6 This bar chart details the frequency with which each of the 26 most common diagnostic criteria were reported as being relevant to each condition. Column headers indicate the condition and the number of relevant manuscripts in brackets. CECS = chronic exertional compartment syndrome, MTSS = medial tibial stress syndrome, tibial SF = tibial stress fracture, SPNES = superficial peroneal nerve entrapment syndrome, PAES = popliteal artery entrapment syndrome, ALLSMS = accessory/low-lying soleus muscle syndrome. CK = Creatine kinase. So2 = Oxygen saturation. NIR'S = Near-infrared spectrometry. ABPI = Ankle brachial pressure index. ICP = Dynamic intracompartment pressure. DTR = Deep tendon reflex. Hx = History

confirm MTSS in severe cases and to exclude other conditions [5, 140, 160–162].

For tibial stress fracture, diagnosis is first based on a history of gradual onset of focal pain during exercise, which matches the presence of localised pain on palpation and positive imaging findings featured in all manuscripts. Other specific tests have been used and reported to be useful in this review, for example, the hop test or fulcrum test. However, the hop test was found to be 100% sensitive and 45% specific for diagnosing medial tibial stress fracture, whereas the fulcrum test was not proposed to be statistically specific and sensitive for the diagnosis of medial tibial stress fracture [38, 110]. Therefore, the history, palpation and imaging modalities are consistent, and core elements are needed to detect stress fracture [38, 110, 163, 164].

The key distinctive element for the diagnosis of PAES was suspicion from the history of intermittent claudication type of pain, positional loss of pulse in the tibialis posterior artery during physical examination, and the absence of pain while resting. Sinha et al. identified that provocation Doppler ultrasonography, ankle brachial pressure index measurement, and MRI & MR angiography have 94% and 90% sensitivity for diagnosing PAES [165]. However, these were only stipulated in ~ 50% of the

relevant manuscripts, suggesting that there is inconsistency of diagnostic approaches that merits further study.

The diagnosis of SPNES was based mainly on a history of nonspecific exertional neurological type of pain over the anterior compartment, accompanied by weakness, positive Tinel's test, and altered sensation. Nerve conduction studies and electromyography were not found to be very useful unless pre- and postexercise symptom provocation tests were employed. MRI could be useful to demonstrate anatomical relations causing nerve compression [10, 166].

Lumbar radiculopathy diagnosis revolves primarily around the history of radiating pain, worsening of pain by coughing, bending, or sitting, neurological symptoms, and altered neurological findings during special testing, supporting the literature [154, 167]. However, this review found that nonexertional pain was relatively common in radiculopathy. This anomaly shows that the onset of pain may be important for radiculopathy diagnosis and that radicular pain can occur without exercise, unlike the other conditions, which are primarily exertional in nature, although pain may linger after exertion for a while in conditions such as MTSS. Radiculopathy diagnosis is mainly confirmed by imaging modalities, particularly MRI, as reported in ~ 70% of the relevant manuscripts.

For *myofascial tears*, the diagnostic criteria reported within this review are quite reflective of the literature (history of traumatic onset and sharp pain) [168, 169]. However, there was conspicuously little literature for this condition, despite it being a more frequent cause of EILP than other muscle/myofascial pathologies, which led to inconsistencies in these findings. This clearly indicates a lack of a diagnostic guide for the condition. Further research is required to establish a more reliable diagnostic approach for myofascial tears. This gap within the literature may clarify the unexpected dominant diagnostic tool for the condition according to the scoping review: imaging modalities. However, these factors play a role in the diagnosis of all conditions; therefore, they are not primary for the diagnostic procedure.

The key distinctive element for the diagnosis of *McArdle's syndrome* is genetic PYGM testing and blood screening (including serum creatine kinase and genetic testing), giving it a clearly different diagnostic approach to the rest of the EILP conditions and is well represented in this review as expected. It was also characterised by early fatigue and myalgia due to genetic abnormalities. Lucia et al. reported that the diagnosis is usually confirmed by muscle biopsy, which shows a negative histochemical reaction for myophosphorylase and no myophosphorylase activity [57]. McArdle's syndrome placed a large importance on further investigations for pathogenic mutations and the presence of certain symptoms and signs, including exercise intolerance and fixed weakness (all of which were returned within 100% of the literature).

For *ALLSMS*, diagnosis relied heavily on the absence of traumatic history, the lack of major presenting symptoms and the presence of anatomical variation on examination with (20%) return. Imaging modalities were recommended to confirm the diagnosis. This was a similar diagnostic approach to the literature, with physical findings revealing an extra muscle or low-lying muscle that can be clearly identified by imaging [102]. Accessory and low-lying soleus muscles are present as normal anatomical variations in many asymptomatic subjects. A significant increase in intensity and exercise workload, in some cases, may lead to pain and disability. These findings are also in agreement with the literature [170].

For each condition, a combination of key aspects of the history, physical findings, and further investigative modalities were fundamental to reach a diagnostic conclusion and formulate a diagnosis. This diagnostic framework will be useful for clinicians dealing with these conditions to improve differential diagnosis and reduce diagnostic variability. These findings must, however, be treated with caution, as study quality is low. Therefore, the diagnostic procedure will be easier when there are established diagnostic guidelines.

Limitations of the literature

There was a paucity of literature for some of the included conditions (myofascial tears, McArdle's syndrome, ALLSMS), as indicated by the low numbers of returns for these conditions. There was a lack of detailed diagnostic studies reporting sensitivity and specificity for any condition and a lack of longitudinal studies and trials, meaning that the strength of our conclusions is limited. Considering the level of evidence for the included manuscripts, the median is four, which indicates that most of the included manuscripts were of low quality. Additionally, there was a lack of consensus on definitions, which we hope our review can facilitate. Moreover, the prevalence of these conditions is not clear within the literature; therefore, there is a need for better epidemiological studies. Finally, the literature does not fully reflect clinical practice, as many patients present with an unclear picture; therefore, a consideration of how to deal with atypical presentations is an important consideration for future research that is not currently represented in the literature.

Limitations of the review

There was only a partial critical appraisal (quality assessment, not risk of bias) of sources conducted due to the type of review. However, the mixed methods appraisal tool (MMAT) was used to assess the quality of the included manuscripts. Moreover, the authorship of the included manuscripts was not considered within this review, and this could be quite influential for the number of results due to the weight of each manuscript, especially where the same author has written multiple papers using the same diagnostic criteria. In addition, the domains we used for defining the conditions do not include the patient's description and patient voice, and this may be worth considering in future work [157, 158]. Moreover, the data included in this review did not compare the evidence of different diagnostic modalities to assist in the diagnosis.

Conclusion

In this scoping review, we developed definitions with a standardised structure and propose an initial diagnostic framework for nine common EILP sub-conditions. Patient history, anatomical symptom location, physical findings and investigative modalities represent crucial diagnostic elements. Many of the findings are consistent across the literature, but others are less so—highlighting both the inconsistencies between different clinical groups' diagnostic approach to EILP and the need for efforts to build consensus. Notwithstanding the overlapping symptoms, primary patterns of presentation vary greatly between the conditions, and clinicians should be able to differentiate using the findings in this

scoping review. All the conditions reported the necessity to exclude other differentials, reinforcing the view that EILP diagnosis is often one of exclusion. Adoption of this diagnostic framework and application of the associated definitions should help reduce care variability, improve EILP diagnosis and therefore improve management and outcome.

Future research recommendations

1. Consensus studies on the resulting definitions and diagnostic criteria could be a potential for future research. We are hopeful that at the end of the current 4-phase study (SR is phase 1), we will have guidelines on diagnostic criteria for EILP.
2. Prevalence of the various conditions needs to be addressed.
3. Intra-compartment pressure measurement in supporting the diagnosis of CECS requires a particular study.
4. Use of dynamic MRI & MRA in the assessment of PAES needs evaluation.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13047-023-00680-6>.

Additional file 1. Collection of EILP diagnosis since 1986.

Additional file 2. Preferred reporting items for systematic reviews and meta-analyses extension for scoping reviews (PRISMA-ScR) checklist. The completed PRISMA-ScR checklist.

Additional file 3. Full search strategy. The full search strategy used within each database.

Additional file 4. Developing a framework for the diagnostic criteria of patients with EILP scoping review data charting table. The data charting table for this review.

Additional file 5. Developing a framework for the diagnostic criteria of patients with EILP scoping review mixed method appraisal tool. The mixed method appraisal tool (MMAT) scoring for this review.

Acknowledgements

Not applicable.

Authors' contributions

Main Authors: Fatma Bosnina, Nat Padhiar, Dylan Morrissey
Statistical analysis: Fatma Bosnina, Stuart Miller
Data collection: Fatma Bosnina, Chrysovalanto Massoura, Krishna Girotra

Funding

The first author is sponsored by the Government of National Unity Libyan Embassy – London Academic Attaché for this research.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

Not applicable.

Author details

¹Sports & Exercise Medicine, William Harvey Research Institute, Barts & The London School of Medicine & Dentistry, Queen Mary University of London, London, UK. ²London Sportswise, London, UK. ³Barts Health NHS Trust Physiotherapy Department, London, UK.

Received: 21 June 2023 Accepted: 31 October 2023

Published online: 21 November 2023

References

1. Orava S, Puranen J. Athletes' leg pains. *Br J Sports Med*. 1979;13(3):92–7.
2. Rajasekaran S, Kvinlaug K, Finnoff JT. Exertional leg pain in the athlete. *PM R*. 2012;4(12):985–1000. <https://doi.org/10.1016/j.pmrj.2012.10.002>.
3. Taunton JE, Ryan MB, Clement DB, McKenzie DC, Lloyd-Smith DR, Zumbo BD. A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med*. 2002;36(2):95–101. Available from: <http://bjsm.bmj.com/>. [Cited 2021 Sep 1].
4. Bonasia DE, Rosso F, Cottino U, Rossi R. Exercise-induced leg pain. *Asia Pac J Sports Med Arthrosc Rehabil Technol*. 2015;2(3):73–84. <https://doi.org/10.1016/j.asmart.2015.03.003>.
5. Lohrer H, Malliaropoulos N, Korakakis V, Padhiar N. Exercise-induced leg pain in athletes: diagnostic, assessment, and management strategies. *Phys Sportsmed*. 2019;47(1):47–59.
6. Rajasekaran S, Finnoff JT. Exertional Leg Pain. *Phys Med Rehabil Clin N Am*. 2016;27(1):91–119.
7. Rosenthal MD, Hawkes N, Garbrecht JD. Popliteal Artery Entrapment Syndrome. *J Orthop Sports Phys Ther*. 2020;50(9):531.
8. Williams EH, Dellon AL. Intraseptal superficial peroneal nerve. *Microsurgery*. 2007;27(5):477–80. Available from: <https://www.scopus.com/inwards/record.uri?eid=2-s2.0-34547870185&doi=10.1002%2Fmicr.20390&partnerID=40&md5=d2173c41cd6442412d6f1f2ba283384a>.
9. Padhiar N, Malliaropoulos N, Lohrer H. Exercise-induced leg pain in sport. *Br J Sports Med*. 2015;49(24):1546–7.
10. Biber Brewer R, Gregory AJM. Chronic lower leg pain in athletes: a guide for the differential diagnosis, evaluation, and treatment. *Sports Health*. 2012;4(2):121–7.
11. Peters MDJ, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc*. 2015;13(3):141–6.
12. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Ann Intern Med*. 2018;169(7):467–73.
13. Peter M, Godfrey M, Christina, McInerney P, SB. The Joanna Briggs Institute Reviewers' Manual 2015: Methodology for JBI scoping reviews. Joanna Briggs Institute. 2015;(February 2016):1–24.
14. Arksey H, O'Malley L. Scoping studies: Towards a methodological framework. *Int J Soc Res Methodol*. 2005;8(1):19–32.
15. Levac D, Colquhoun H, OK. Scoping studies: advancing the methodology. *Implement Sci* 2010; 5:69. 2010;1–18.
16. Truong LK, Mosewich AD, Holt CJ, Le CY, Miciak M, Whittaker JL. Psychological, social and contextual factors across recovery stages following a sport-related knee injury: a scoping review. *Br J Sports Med*. 2020;54(19):1149–56.
17. Griffin SA, Panagodage Perera NK, Murray A, Hartley C, Fawcner SG, P T Kemp S, et al. The relationships between rugby union, and health and well-being: A scoping review. *Br J Sports Med*. 2021;55(6):319–26.
18. Grant MJ, Booth A. A typology of reviews: An analysis of 14 review types and associated methodologies. *Health Info Libr J*. 2009;26(2):91–108.
19. Zachary M, Pollock D, Khalil H, Alexander L, McInerney PM, Godfrey C, et al. What are scoping reviews? Providing a formal definition of scoping reviews as a type of evidence synthesis. 2022. p. 950–2.

20. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol*. 2018;18(1):143.
21. Bosnina F, Padhiar N, Miller S, Girotra K, Massoura C, Morrissey D. Developing a framework for the diagnostic criteria of patients with Exercise Induced Leg Pain (EILP): a scoping review. *Open Science Framework*. 2022;15(2):1–23. Available from: <https://osf.io/4k27b>
22. Bosnina F, Padhiar N, Miller S, Girotra K, Massoura C, Morrissey D. Developing a framework for the diagnostic criteria of patients with Exercise Induced Leg Pain (EILP): A scoping review protocol. 2022.
23. Peters MDJ, Marnie C, Tricco AC, Pollock D, Munn Z, Alexander L, et al. Updated methodological guidance for the conduct of scoping reviews. *JBI Evid Synth*. 2020;18(10):219–26.
24. de Bruijn JA, van Zantvoort APM, van Klaveren D, Winkes MB, van der Cruisjen-Raaijmakers M, Hoogeveen AR, et al. Factors predicting lower leg chronic exertional compartment syndrome in a large population. *Int J Sports Med*. 2018;39(1):58–66.
25. Vogels S, Ritchie ED, van Dongen TTFC, Scheltinga MRM, Zimmermann WO, Hoencamp R. Systematic review of outcome parameters following treatment of chronic exertional compartment syndrome in the lower leg. *Scand J Med Sci Sports*. 2020;30(10):1827–45.
26. Johnson N, Phillips M. Rayyan for systematic reviews. *J Electron Resour Librariansh*. 2018;30(1):46–8.
27. Peters MDJ, Godfrey C, Mclnerney P, Baldini Soares C, Khalil H PD. In: Aromataris E, Munn Z (Editors). *Joanna Briggs Institute Reviewer's Manual*, JBI, 2017. Chapter 11: Scoping reviews. Available from: <https://jbi-global-wiki.refined.site/space/MANUAL/3283910770/Chapter+11%3A+Scoping+reviews>.
28. Hong QN, Gonzalez-Reyes A, Pluye P. Improving the usefulness of a tool for appraising the quality of qualitative, quantitative and mixed methods studies, the Mixed Methods Appraisal Tool (MMAT). *J Eval Clin Pract*. 2018;24(3):459–67. <https://doi.org/10.1111/jep.12884>.
29. Pace R, Pluye P, Bartlett G, Macaulay AC, Salsberg J, Jagosh J, et al. Testing the reliability and efficiency of the pilot Mixed Methods Appraisal Tool (MMAT) for systematic mixed studies review. *Int J Nurs Stud*. 2012;49(1):47–53. <https://doi.org/10.1016/j.ijnurstu.2011.07.002>.
30. Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect Dis*. 2022;22(4):e102–7. [https://doi.org/10.1016/S1473-3099\(21\)00703-9](https://doi.org/10.1016/S1473-3099(21)00703-9).
31. Sargeant JM, O'Connor AM. Scoping reviews, systematic reviews, and meta-analysis: applications in veterinary medicine. *Front Vet Sci*. 2020;7(January):1–14.
32. Pollock D, Tricco AC, Peters MDJ, Mclnerney PA, Khalil H, Godfrey CM, et al. Methodological quality, guidance, and tools in scoping reviews: a scoping review protocol and 12 The Scottish Centre for Evidence-based Multiprofessional Practice: A JBI Centre of Excellence, Aberdeen, Great Britain. Vol. 20, *JBI Evid Synth*. 2022. Available from: <https://jbi.global/>
33. Miller TL, Backs R, Vaccaro PS. Popliteal artery entrapment syndrome: a diagnostic and treatment enigma for orthopaedic surgeons. *J Am Acad Orthop Surg*. 2021;29(17):e834–45.
34. Winkes M, van Eerten P, Scheltinga M. Deep posterior chronic exertional compartment syndrome as a cause of leg pain. *Unfallchirurg*. 2020;123(Suppl 1):3–7.
35. Velasco TO, Leggit JC. Chronic exertional compartment syndrome: a clinical update. *Curr Sports Med Rep*. 2020;19(9):347–52.
36. Nwakibu U, Schwarzman G, Zimmermann WO, Hutchinson MR. Chronic exertional compartment syndrome of the leg management is changing: where are we and where are we going? *Curr Sports Med Rep*. 2020;19(10):438–44.
37. de Bruijn J, Winkes M, van Eerten P, Scheltinga M. Chronic exertional compartment syndrome as a cause of anterolateral leg pain. *Unfallchirurg*. 2019;2020(123):8–14.
38. Feldman JJ, Bowman EN, Phillips BB, Weinklein JC. Tibial Stress Fractures in Athletes. *Orthop Clin North Am*. 2016;47(4):733–41. <https://doi.org/10.1016/j.joc.2016.05.015>.
39. Braver RT. Chronic exertional compartment syndrome. *Clin Podiatr Med Surg*. 2016;33(2):219–33.
40. Burrus MT, Werner BC, Starman JS, Gwathmey FW, Carson EW, Wilder RP, et al. Chronic leg pain in athletes. *Am J Sports Med*. 2015;43(6):1538–47.
41. Pegrum J, Crisp T, Padhiar N, Flynn J. The pathophysiology, diagnosis, and management of stress fractures in postmenopausal women. *Physician and Sportsmedicine*. 2012;40(3):32–42.
42. Van Boxem K, Cheng J, Patijn J, van Kleef M, Lataster A, Mekhail N, et al. 11. Lumbosacral radicular pain. *Pain Pract*. 2010;10(4):339–58.
43. Harrast MA, Colonno D. Stress fractures in runners. *Clin Sports Med*. 2010;29(3):399–416.
44. Fredericson M, Jennings F, Beaulieu C, Matheson GO. Stress fractures in athletes. *Top Magn Reson Imaging*. 2006;17(5):309–25.
45. Englund J. Chronic compartment syndrome: tips on recognising and treating. *J Fam Pract*. 2005;64(12):599–603.
46. Bong MR, Polatsch DB, Jazrawi LM, Rokito AS. Chronic exertional compartment syndrome: diagnosis and management. *Bull Hosp Jt Dis*. 2005;62(3–4):77–84.
47. Peris P. Stress fractures. *Best Pract Res Clin Rheumatol*. 2003;17(6):1043–61.
48. Couture CJ, Karlson KA. Tibial stress injuries: decisive diagnosis and treatment of “shin splints.” *Phys Sportsmed*. 2002;30(6):29–36.
49. Korkola M, Amendola A. Exercise-Induced Leg Pain: Sifting Through a Broad Differential. *Physician and Sportsmedicine* [Internet]. 2001;29(6):35–50.
50. Boden BP, Osbahr DC. High-risk stress fractures: evaluation and treatment. *J Am Acad Orthop Surg*. 2000;8(6):344–53.
51. Touliopolous S, Hershman EB. Lower leg pain. Diagnosis and treatment of compartment syndromes and other pain syndromes of the leg. *Sports Med*. 1999;27(3):193–204.
52. Schepis AA, Lynch G. Exertional compartment syndromes of the lower extremity. *Curr Opin Rheumatol*. 1996;8(2):143–7.
53. Reneman RS. The anterior and the lateral compartmental syndrome of the leg due to intensive use of muscles. *Clin Orthop Relat Res*. 1975;113:69–80.
54. Hutchinson MR, Ireland ML. Common compartment syndromes in athletes. *Treat Rehabil Sports Med*. 1994;17(3):200–8.
55. Brukner P, Bennell K. Stress fractures in female athletes. Diagnosis, management and rehabilitation. *Sports Med*. 1997;24(6):419–29.
56. Neslon BJ, Arciero RA. Stress fractures in the female athlete. *Curr Sports Med Rep*. 2002;4(6):323–8.
57. Lucia A, Nogales-Gadea G, Pérez M, Martín MA, Andreu AL, Arenas J. McArdle disease: What do neurologists need to know? *Nat Clin Pract Neurol*. 2008;4(10):568–77.
58. Brennan FH, Kane SF. Diagnosis, treatment options, and rehabilitation of chronic lower leg exertional compartment syndrome. *Curr Sports Med Rep*. 2003;2(5):247–50. <https://doi.org/10.1249/00149619-200310000-00003>.
59. Moeyersoons JP, Martens M. Chronic compartment syndrome: diagnosis and management. *Acta Orthop Belg*. 1992;58(1):23–7.
60. Hameed M, Coupland A, Davies AH. Popliteal artery entrapment syndrome: an approach to diagnosis and management. *Br J Sports Med*. 2018;52(16):1073–4.
61. Padhiar N, King JB. Exercise induced leg pain: chronic compartment syndrome. Is the increase in intracompartment pressure exercise specific? *Br J Sports Med*. 1997;31:353.
62. Mukish P, Reybet-Degat PY, Demangel A, Trouilloud P, Baulot E. Accessory soleus muscle: a difficult diagnosis: a case report and a review of the literature. *Eur J Orthop Surg Traumatol*. 1999;22(SUPPL. 1):610–2.
63. Young AJ, McAllister DR. Evaluation and treatment of tibial stress fractures. *Clin Sports Med*. 2006;25(1):117–28.
64. Ross DG. Chronic compartment syndrome. *Orthopaedic Nurs*. 1996;15(3):23–7.
65. Winters M. The diagnosis and management of medial tibial stress syndrome: an evidence update. *Unfallchirurg*. 2020;123(Suppl 1):15–9.
66. Wuellner JC, Nathe CD, Kreulen CD, Burnham KJ, Giza E. Chronic Exertional Compartment Syndrome: The Athlete's Claudication. *Oper Tech Sports Med*. 2017;25(2):52–8. <https://doi.org/10.1053/j.otsm.2017.03.004>.
67. Coscolluela PE, McCulloch PC. Anterior Tibial Stress Fractures: Intramedullary Nail vs Anterior Tension Band Plating. *Oper Tech Sports Med*. 2017;25(2):67–74. <https://doi.org/10.1053/j.otsm.2017.03.003>.
68. Joy SM, Raudales R. Popliteal Artery Entrapment Syndrome. *Curr Sports Med Rep*. 2015;14(5):364–7. Available from: <https://www.scopus.com/inwards/record.uri?eid=2-s2.0-84941286577&doi=10>.

- 1249%2FJSR.000000000000199&partnerID=40&md5=8f16ae3672f851e011322b4559180c24.
69. Flick D, Flick R. Chronic exertional compartment syndrome testing. *Curr Sports Med Rep.* 2015;14(5):380–5.
 70. Williamson BL, Arthur CHC. Shin-splints: common exercise-related syndromes affecting the lower leg. *J R Nav Med Serv.* 2014;100(3):272–6.
 71. Dunn JC, Waterman BR. Chronic exertional compartment syndrome of the leg in the military. *Clin Sports Med.* 2014;33(4):693–705.
 72. Craig A. Entrapment neuropathies of the lower extremity. *PM and R.* 2013;5(5 SUPPL.):31–40.
 73. Murray MC, Heckman MM. Chronic exertional compartment syndrome: diagnostic techniques and management. *Tech Orthop.* 2012;27(1):75–8.
 74. Murdock M, Murdoch MM. Compartment syndrome: a review of the literature. *Clin Podiatr Med Surg.* 2012;29(2):301–10, viii.
 75. Patel DS, Roth M, Kapil N. Stress fractures: Diagnosis, Treatment, and prevention. *Am Fam Physician.* 2011;83(1):39–46. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L361808068&from=export>.
 76. Wilder RP, Magrum E. Exertional compartment syndrome. *Clin Sports Med.* 2010;29(3):429–35.
 77. McDonald S, Bearcroft P. Compartment syndromes. *Semin Musculoskeletal Radiol.* 2010;14(2):236–44.
 78. Gill CS, Halstead ME, Matava MJ. Chronic exertional compartment syndrome of the leg in athletes: evaluation and management. *Phys Sportsmed.* 2010;38(2):126–32.
 79. Pell 4th. RF, Khanuja HS, Cooley GR. Leg pain in the running athlete. *J Am Acad Orthop Surg.* 2004;12(6):396–404. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L39701583&from=export>.
 80. Stager A, Clement D. Popliteal artery entrapment syndrome. *Sports Med.* 1999;28(1):61–70.
 81. Styf J. Chronic exercise-induced pain in the anterior aspect of the lower leg. An overview of diagnosis. *Sports Med.* 1989;7(5):331–9.
 82. Detmer DE. Chronic shin splints. Classification and management of medial tibial stress syndrome. *Sports Med.* 1986;3(6):436–46.
 83. Leach RE, Zohn DA, Stryker WS. Anterior tibial compartment syndrome due to strenuous exercise. *Mil Med.* 1964;129(7):610–3. <https://doi.org/10.1093/milmed/129.7.610>.
 84. George CA, Hutchinson MR. Chronic exertional compartment syndrome. *Clin Sports Med.* 2012;31(2):307–19.
 85. Al-Tayef TA, Rziki A, Rasras H, el Mahi O, Benzirar A. Popliteal artery entrapment syndrome: a case report with literature review. *Pan Afr Med J.* 2021;39:80.
 86. Finnoff JT, Johnson W. Ultrasound-guided fasciotomy for chronic exertional compartment syndrome: a case report. *Clin J Sport Med.* 2020;30(6):e231–3.
 87. Kokalj J, Majstorović M. Bilateral Distal Tibial Stress Fracture in a Triathlete. *Acta Chir Orthop Traumatol Cech.* 2019;86(2):156–8.
 88. Drigny J, Reboursière E, Desvergée A, Ruet A, Hulet C. Concurrent exertional compartment syndrome and functional popliteal artery entrapment syndrome: a case report. *PM R.* 2019;11(6):669–72.
 89. Henry HT, Szolomayer LK, Sumpio BE, Sutton KM. Popliteal artery entrapment syndrome: bilateral lower extremity involvement. *Orthopedics.* 2018;41(2):e295–8.
 90. Carneiro Júnior FCF, Carrizo ENDA, Araújo ST, Nakano LCU, de Amorim JE, Cacione DG. Popliteal artery entrapment syndrome: a case report and review of the literature. *Am J Case Rep.* 2018;19:29–34.
 91. Cormier DJ, Gellhorn AC, Singh JR. Soleus muscle herniation with magnetic resonance imaging and ultrasound correlation in a female long-distance runner: a case report. *PM R.* 2017;9(5):529–32.
 92. Myers RJ, Murdock EE, Farooqi M, van Ness G, Crawford DC. A unique case of common peroneal nerve entrapment. *Orthopedics.* 2015;38(7):e644–6.
 93. Tsilogianni Z, Grapatsas K, Papanikolaou Z, Kokkini-Paschou A, Tsantilas A, Tsiligiris V, et al. Popliteal artery entrapment syndrome: a common cause of a rare clinical entity—critical leg ischemia in the young. *Mil Med.* 2014;179(1):e124–6.
 94. Jovčić M, Jovčić V, Hrković M, Lazović M. Medial tibial stress syndrome: case report. *Med Pregl.* 2014;67(7–8):247–51.
 95. McAree BJ, O'Donnell ME, Davison GW, Boyd C, Lee B, Soong CV. Bilateral popliteal artery occlusion in a competitive bike rider: case report and clinical review. *Vasc Endovascular Surg.* 2008;42(4):380–5.
 96. Farr D, Selesnick H. Chronic exertional compartment syndrome in a collegiate soccer player: a case report and literature review. *Am J Orthop (Belle Mead NJ).* 2008;37(7):374–7.
 97. Schraml FV, Riego De Dios RL, Flemming DJ. Exercise-related longitudinal tibial stress fracture in a young person. *Ann Nucl Med.* 2006;20(6):441–4.
 98. Bryan BM, Lutz GE, O'Brien SJ. Sural nerve entrapment after injury to the gastrocnemius: a case report. *Arch Phys Med Rehabil.* 1999;80(5):604–6.
 99. Grimley RP, O'Garra JA, Cheate T, Mohan SR, Hickman P. Popliteal artery entrapment syndrome. *Int J Surg Case Rep.* 1990;1(2):13–5.
 100. Bell S. Intracompartmental pressures on exertion in a patient with a popliteal artery entrapment syndrome. *Am J Sports Med.* 1985;13(5):365–6.
 101. McAuliffe TB, Fiddian NJ, Browett JP. Entrapment neuropathy of the superficial peroneal nerve. A bilateral case. 2022.
 102. Rubio FJ, Franco L, Montero MJ, Ugarte P, Valero A. Accessory soleus muscle in an athlete. Presentation of a case and a literature review. *Apunts Medicina de l'Esport.* 2015;50(186):79–82.
 103. Allen GJ. Longitudinal stress fractures of the tibia: diagnosis with CT. *Radiology.* 1988;167(3):799–801.
 104. McDonald PT, Easterbrook JA, Rich NM, Collins GJJ, Kozloff L, Claggett GP, et al. Popliteal artery entrapment syndrome. Clinical, noninvasive and angiographic diagnosis. *Am J Surg.* 1980;139(3):318–25.
 105. Brodie JT, Dormans JP, Gregg JR, Davidson RS. Accessory soleus muscle. *Feuillets de Radiologie.* 1997;51(6):345–7.
 106. Lauder TD, Stuart MJ, Amrami KK, Felmlee JP. Exertional compartment syndrome and the role of magnetic resonance imaging. *Am J Phys Med Rehabil.* 2002;81(4):315–9.
 107. Yang LJS, Gala VC, McGillicuddy JE. Superficial peroneal nerve syndrome: an unusual nerve entrapment. Case report. *J Neurosurg.* 2006;104(5):820–3.
 108. Truex N, Menge TJ. Bilateral chronic exertional compartment syndrome of the leg: a rare but often debilitating condition in athletes. *S D Med.* 2018;71(7):310–4.
 109. Romanus B, Lindahl S, Stener B. Accessory soleus muscle. A clinical and radiographic presentation of eleven cases. *J Bone Joint Surg Series A.* 1986;68(5):731–4.
 110. Milgrom C, Zloczower E, Fleischmann C, Spitzer E, Landau R, Bader T, et al. Medial tibial stress fracture diagnosis and treatment guidelines. *J Sci Med Sport.* 2021;24(6):526–30. <https://doi.org/10.1016/j.jsams.2020.11.015>.
 111. van Zantvoort A, Setz M, Hoogeven A, van Eerten P, Scheltinga M. Chronic lower leg pain: entrapment of common peroneal nerve or tibial nerve. *Unfallchirurg.* 2020;123(Suppl 1):20–4.
 112. Drexler M, Rutenberg TF, Rozen N, Warschawski Y, Rath E, Chechik O, et al. Single minimal incision fasciotomy for the treatment of chronic exertional compartment syndrome: outcomes and complications. *Arch Orthop Trauma Surg.* 2017;137(1):73–9.
 113. Fabre T, Montero C, Gaujard E, Gervais-Dellion F, Durandea A. Chronic calf pain in athletes due to sural nerve entrapment: a report of 18 cases. *Am J Sports Med.* 2000;28(5):679–82.
 114. Van der Velde GM, Hsu WS. Posterior tibial stress fracture: a report of three cases. *J Manipulative Physiol Ther.* 1999;22(5):341–6.
 115. Clayer M, Krishnan J, Lee WK, Tamblin P. Longitudinal stress fracture of the tibia: two cases. *Clin Radiol.* 1992;46(6):401–4.
 116. Clement DB. Tibial stress syndrome in athletes. *Am J Sports Med.* 1974;2(2):81–5.
 117. Detmer DE, Sharpe K, Sufit RL, Girdley FM. Chronic compartment syndrome: diagnosis, management, and outcomes. *Am J Sports Med.* 1985;13(3):162–70.
 118. Fouasson-Chailloux A, Menu P, Alloquent J, Dauty M. Determination of the predictive clinical parameters to diagnose chronic exertional compartment syndrome. *Eur J Sport Sci.* 2018;18(2):279–85.
 119. Fronek J, Mubarak SJ, Hargens AR, Lee YF, Gershuni DH, Garfin SR, et al. Management of chronic exertional anterior compartment syndrome of the lower extremity. *Clin Orthop Relat Res.* 1987;220:217–27.
 120. Ha KI, Hahn SH, Chung MY, Yang BK, Yi SR. A clinical study of stress fractures in sports activities. *Orthopedics.* 1991;14(10):1089–95.

121. Lane R, Nguyen T, Cuzzilla M, Oomens D, Mohabbat W, Hazelton S. Functional popliteal entrapment syndrome in the sportsperson. *Eur J Vasc Endovasc Surg*. 2012;43(1):81–7. <https://doi.org/10.1016/j.ejvs.2011.10.013>.
122. Lindorsson S, Zhang Q, Brisby H, Rennerfelt K. Significantly lower intramuscular pressure in the posterior and lateral compartments compared with the anterior compartment suggests alterations of the diagnostic criteria for chronic exertional compartment syndrome in the lower leg. *Knee Surg Sports Traumatol Arthrosc*. 2021;29(4):1332–9.
123. Mubarak SJ, Gould RN, Lee YF, Schmidt DA, Hargens AR. The medial tibial stress syndrome. A cause of shin splints. *Am J Sports Med*. 1982;10(4):201–5.
124. Orava S, Huikko A. Stress fracture of the mid-tibial shaft. 1984. p. 35–7.
125. Puranen J. The medial tibial syndrome: exercise ischaemia in the medial fascial compartment of the leg. *J Bone Joint Surg Br*. 1974;56-B(4):712–5.
126. Styf J. Diagnosis of exercise-induced pain in the anterior aspect of the lower leg. *Am J Sports Med*. 1988;16(2):165–9.
127. Styf J, Lysell E. Chronic compartment syndrome in the erector spinae muscle. *Spine (Phila Pa 1976)*. 1987;12(7):680–2.
128. DiMauro S, Andreu A, Bruno C, Hadjigeorgiou G. Myophosphorylase Deficiency (Glycogenosis Type V McArdle Disease). *Curr Mol Med*. 2005;2(2):189–96.
129. Leach RE, Purnell MB, Saito A. Peroneal nerve entrapment in runners. *Am J Sports Med*. 1989;17(2):287–91.
130. Lovelock T, Claydon M, Dean A. Functional popliteal artery entrapment syndrome: an approach to diagnosis and management. *Int J Sports Med*. 2021;42(13):1159–66.
131. Vajapey S, Miller TL. Evaluation, diagnosis, and treatment of chronic exertional compartment syndrome: a review of current literature. *Phys Sportsmed*. 2017;45(4):391–8.
132. Moen MH, Tol JL, Weir A, Steunebrink M, de Winter TC. Medial tibial stress syndrome: a critical review. *Sports Med*. 2009;39(7):523–46.
133. Galbraith RM, Lavallee ME. Medial tibial stress syndrome: Conservative treatment options. *Curr Rev Musculoskelet Med*. 2009;2(3):127–33.
134. McInnis KC, Ramey LN. High-risk stress fractures: diagnosis and management. *PM R*. 2016;8(3 Suppl):S113–24.
135. Veith RG, Matsen FA, Newell SG. Recurrent anterior compartmental syndromes. *Phys Sportsmed*. 1980;8(11):80–8.
136. Wallensten R, Eriksson E. Intramuscular pressures in exercise-induced lower leg pain. *Int J Sports Med*. 1984;5(1):31–5.
137. Roscoe D, Roberts AJ, Hulse D. Intramuscular compartment pressure measurement in chronic exertional compartment syndrome: new and improved diagnostic criteria. *Am J Sports Med*. 2015;43(2):392–8.
138. Pedret C, Balius R, Blasi M, Dávila F, Aramendi JF, Masci L, et al. Ultrasound classification of medial gastrocnemius injuries. *Scand J Med Sci Sports*. 2020;30(12):2456–65.
139. Pedowitz RA, Hargens AR, Mubarak SJ, Gershuni DH. Modified criteria for the objective diagnosis of chronic compartment syndrome of the leg. *Am J Sports Med*. 1990;18(1):35–40.
140. Winters M. Medial tibial stress syndrome: Diagnosis, treatment and outcome assessment (PhD Academy Award). *Br J Sports Med*. 2018;52(18):1213–4.
141. Orlin JR, Lied IH, Stranden E, Irgens HU, Andersen JR. Prevalence of chronic compartment syndrome of the legs: Implications for clinical diagnostic criteria and therapy. *Scand J Pain*. 2016;12:7–12.
142. Tzortziou V, Maffulli N, Padhiar N. Diagnosis and management of chronic exertional compartment syndrome (CECS) in the United Kingdom. *Clin J Sport Med*. 2006;16(3):209–13.
143. Ghasabmahaleh SH, Rezasoltani Z, Dadarkhah A, Hamidipanah S, Mofrad RK, Najafi S. Spinal manipulation for subacute and chronic lumbar radiculopathy: a randomised controlled trial. *Am J Med*. 2021;134(1):135–41. <https://doi.org/10.1016/j.amjmed.2020.08.005>.
144. Thein R, Tilbor I, Rom E, Herman A, Haviv B, Burstein G, et al. Return to sports after chronic anterior exertional compartment syndrome of the leg: Conservative treatment versus surgery. *J Orthop Surg*. 2019;27(2). Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L626932279&from=export>
145. Satpute K, Hall T, Bisen R, Lokhande P. The effect of spinal mobilisation with leg movement in patients with lumbar radiculopathy—a double-blind randomised controlled trial. *Arch Phys Med Rehabil*. 2019;100(5):828–36.
146. Yates B, White S. The Incidence and Risk Factors in the Development of Medial Tibial Stress Syndrome among Naval Recruits. *Am J Sports Med*. 2004;32(3):772–80.
147. FA M, RG V. compartmental syndrome in children. *J Pediatr Orthop*. 1981;1:33–41. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7341650>.
148. Black KP, Taylor DE. Current concepts in the treatment of common compartment syndromes in athletes. *Sports Med*. 1993;15(6):408–18.
149. Carmont MR, Mei-Dan O, Bennell KL. Stress fracture management: current classification and new healing modalities. *Oper Tech Sports Med*. 2009;17(2):81–9. <https://doi.org/10.1053/j.otsm.2009.05.004>.
150. Garcia JC. Nerve entrapment. *Telemicrosurgery: Robot Assisted. Microsurgery*. 2013;104:109–18.
151. Tzika M, Paraskevas G, Natsis K. Entrapment of the superficial peroneal nerve: an anatomical insight. *J Am Podiatr Med Assoc*. 2015;105(2):150–9.
152. Balius R, Blasi M, Pedret C, Alomar X, Peña-Amaro J, Vega JA, et al. A histoarchitectural approach to skeletal muscle injury: searching for a common nomenclature. *Orthop J Sports Med*. 2020;8(3):1–8.
153. Balius R, Maestro A, Pedret C, Estruch A, Mota J, Rodríguez L, et al. Central aponeurosis tears of the rectus femoris: practical sonographic prognosis. *Br J Sports Med*. 2009;43(11):818–24.
154. Berry JA, Elia C, Saini HS, Miulli DE. A review of lumbar radiculopathy, diagnosis, and treatment. *Cureus*. 2019;11(10):e5934.
155. Plečko M, Knežević I, Dimnjaković D, Josipović M, Bojanić I. Accessory soleus muscle: two case reports with a completely different presentation caused by the same entity. *Case Rep Orthop*. 2020;2020:1–7.
156. Lecocq J, Parier J, Fischer M. Le muscle soléaire accessoire. 2005. p. 232–8.
157. Vicenzino B, De Vos RJ, Alfredson H, Bahr R, Cook JL, Coombes BK, et al. ICON 2019 - International Scientific Tendinopathy Symposium Consensus: there are nine core health-related domains for tendinopathy (CORE DOMAINS): Delphi study of healthcare professionals and patients. *Br J Sports Med*. 2020;54(8):444–51.
158. Scott A, Squier K, Alfredson H, Bahr R, Cook JL, Coombes B, et al. ICON 2019: International Scientific Tendinopathy Symposium Consensus: Clinical Terminology. *Br J Sports Med*. 2020;54(5):260–2.
159. Aweid O, del Buono A, Malliaras P, Iqbal H, Morrissey D, Maffulli N, et al. Systematic review and recommendations for intracompartmental pressure monitoring in diagnosing chronic exertional compartment syndrome of the leg. *Clin J Sport Med*. 2012;22(4):356–70.
160. Winters M, Bakker EWP, Moen MH, Barten CC, Teeuwen R, Weir A. Medial tibial stress syndrome can be diagnosed reliably using history and physical examination. *Br J Sports Med*. 2018;52(19):1267–72.
161. Mulvad B, Nielsen RO, Lind M, Ramskov D. Diagnoses and time to recovery among injured recreational runners in the RUN CLEVER trial. *PLoS One*. 2018;13(10):e0204742. [Cited 2021 Sep 9].
162. Winters M. The diagnosis and management of medial tibial stress syndrome: an evidence update [Diagnostik und Therapie des Schienbeinkantensyndroms: Update zur Studienlage]. *Unfallchirurg*. 2020;123:15–9.
163. Dash N, Kushwaha AS. Stress fractures—a prospective study amongst recruits. *Med J Armed Forces India*. 2012;68(2):118–22.
164. Xu C, Silder A, Zhang J, Reifman J, Unnikrishnan G. A cross-sectional study of the effects of load carriage on running characteristics and tibial mechanical stress: implications for stress-fracture injuries in women. *BMC Musculoskelet Disord*. 2017;18(1):125.
165. Sinha S, Houghton J, Holt PJ, Thompson MM, Loftus IM, Hinchliffe RJ. Popliteal entrapment syndrome. *J Vasc Surg*. 2012;55(1):252–262.e30. <https://doi.org/10.1016/j.jvs.2011.08.050>.
166. Garwood ER, Duarte A, Bencardino JT. MR imaging of entrapment neuropathies of the lower extremity. *Radiol Clin North Am*. 2018;56(6):997–1012.
167. Ghasabmahaleh SH, Rezasoltani Z, Dadarkhah A, Hamidipanah S, Mofrad RK, Najafi S. Spinal manipulation for subacute and chronic lumbar radiculopathy: a randomised controlled trial. *Am J Med*. 2021;134(1):135–41. <https://doi.org/10.1016/j.amjmed.2020.08.005>.
168. Nsitem V. Diagnosis and rehabilitation of gastrocnemius muscle tear: a case report. *J Can Chiropr Assoc*. 2013;57(4):327–33. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24302780>. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3845475>.

169. Flecca D, Tomei A, Ravazzolo N, Martinelli M, Giovagnorio F. US evaluation and diagnosis of rupture of the medial head of the gastrocnemius (tennis leg). *J Ultrasound*. 2007;10(4):194–8.
170. Mukish P, Reybet-Degat PY, Demangel A, Trouilloud P, Baulot E. Accessory soleus muscle: a difficult diagnosis: a case report and a review of the literature. *Eur J Orthop Surg Traumatol*. 2012;22(SUPPL. 1):610–2.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

