

CORRECTION

Open Access



Correction to: Intrauterine resuscitation during the second stage of term labour by maternal hyperoxygenation versus conventional care: study protocol for a randomised controlled trial (INTEREST O2)

Lauren M. Bullens^{1,2*}, Alexandra D. J. Hulsboom¹, Suzanne Moors¹, Rohan Joshi^{3,4}, Pieter J. van Runnard Heimeel¹, M. Beatrijs van der Hout-van der Jagt^{1,2}, Edwin R. van den Heuvel⁵ and S. Guid Oei^{1,2}

Correction

Following publication of the original article [1], the authors noticed that the sample size for the study group was incorrectly reported in the Methods section. The correct number of patients to participate in the study is 116.

Author details

¹Department of Obstetrics and Gynaecology, Máxima Medical Centre, PO Box 7777, 5500 MB Veldhoven, The Netherlands. ²Department of Electrical Engineering, Eindhoven University of Technology, PO Box 513, 5500 MB Eindhoven, The Netherlands. ³Department of Clinical Physics, Máxima Medical Centre, PO Box 7777, 5500 MB Veldhoven, The Netherlands. ⁴Department of Industrial Design, Eindhoven University of Technology, PO Box 513, 5500 MB Eindhoven, The Netherlands. ⁵Department of Mathematics and Computer Science, Eindhoven University of Technology, PO Box 513, 5500 MB Eindhoven, The Netherlands.

Received: 4 October 2018 Accepted: 4 October 2018

Published online: 23 October 2018

Reference

1. Bullens LM, et al. Intrauterine resuscitation during the second stage of term labour by maternal hyperoxygenation versus conventional care: study protocol for a randomised controlled trial (INTEREST O2). *Trials*. 2018;19:195. <https://doi.org/10.1186/s13063-018-2567-x>.

* Correspondence: laurenbullens@gmail.com

¹Department of Obstetrics and Gynaecology, Máxima Medical Centre, PO Box 7777, 5500 MB Veldhoven, The Netherlands

²Department of Electrical Engineering, Eindhoven University of Technology, PO Box 513, 5500 MB Eindhoven, The Netherlands

