


STUDY PROTOCOL

Open Access



# Preemptive analgesia for hemorrhoidectomy: study protocol for a prospective, randomized, double-blind trial

Ekaterina Kazachenko<sup>1</sup>, Tatiana Garmanova<sup>1</sup>, Alexander Derinov<sup>2</sup>, Daniil Markaryan<sup>1</sup>, Hanjoo Lee<sup>3</sup>, Sabrina Magbulova<sup>2\*</sup>  and Petr Tsarkov<sup>2</sup>

## Abstract

**Background:** Hemorrhoidectomy is associated with intense postoperative pain that requires multimodal analgesia. It includes nonsteroidal anti-inflammatory drugs (NSAIDs), acetaminophen, and local anesthetics to reach adequate pain control. There are data in literature preemptive analgesia could decrease postoperative pain after hemorrhoidectomy. The aim of this study is to assess the efficacy of preemptive analgesia with ketoprofen 100 mg 2 h before procedure per os with spinal anesthesia to decrease postoperative pain according to visual analog scale and to reduce the opioids and other analgesics consumption.

**Methods:** Patients of our clinic who meet the following inclusion criteria are included: hemorrhoids grade III–IV and the planned Milligan-Morgan hemorrhoidectomy. After signing the consent all participants are randomly divided into 2 groups: the first one gets a tablet with 100 mg ketoprofen, the second one gets a tablet containing starch per os 2 h before surgery (72 participants per arm). Patients of both arms receive spinal anesthesia and undergo open hemorrhoidectomy. Following the procedure the primary and secondary outcomes are evaluated: opioid administration intake, the pain at rest and during defecation, duration, and frequency of other analgesics intake, readmission rate, overall quality of life, time from the procedure to returning to work, and the complications rate.

**Discussion:** Multimodality pain management has been shown to improve pain control and decrease opioid intake in patients after hemorrhoidectomy in several studies. Gabapentin can be considered as an alternative approach to pain control as NSAIDs have limitative adverse effects. Systemic admission of ketorolac with local anesthetics also showed significant efficacy in patients undergoing anorectal surgery. We hope to prove the efficacy of multimodal analgesia including preemptive one for patients undergoing excisional hemorrhoidectomy that will help to hold postoperative pain levels no more than 3–4 points on VAS with minimal consumption of opioid analgesics.

**Trial registration:** [ClinicalTrials.gov NCT04361695](https://clinicaltrials.gov/ct2/show/study/NCT04361695). Registered on April 24, 2020, version 1.0.

**Keywords:** Analgesics, Anorectal surgery, Hemorrhoidectomy, Pain management, Preemptive analgesia

\* Correspondence: [magbulova@kkmx.ru](mailto:magbulova@kkmx.ru)

<sup>2</sup>Department: Clinic of Colorectal and Minimally Invasive Surgery, Sechenov First Moscow State Medical University, Moscow 101000, Russia  
Full list of author information is available at the end of the article



© The Author(s). 2022 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Anorectal diseases are mostly benign and do not affect life expectancy, so currently both patients and clinicians are extremely interested in performing day care surgery. Anorectal surgery itself causes certain discomfort, especially pain, long recovery period, and a decreasing quality of life for several months.

The pain appears as a result of endogenous and exogenous factors affecting on the peripheral nerves endings, and also the pathological central nervous system excitation. Any surgical intervention activates nociceptors with various stimuli including mechanical factors as a result of a prick or cut of the tissue, chemical factors as a result of exposure to inflammatory mediators, and thermal factors as a result of heating or cooling the tissue [1]. Thus, any surgery is always accompanied by a pain syndrome with various pain severities.

In anorectal surgery, almost every patient experiences moderate/severe pain in the postoperative period, 12% of patients have severe pain throughout the recovery period, and control of postoperative pain is still problematic in 5% of cases when a severe pain syndrome continues despite standard pain management. It leads to long hospital admission and to an opioid intake increase [2]. The problem of analgesia is still relevant for patients after anorectal surgery, because the diseases affect more than 50% of the population over 50 years [3]. Excisional hemorrhoidectomy is the most effective method of stage III–IV hemorrhoidal disease treatment. However it is associated with intense postoperative pain which decreases significantly the quality of life in the postoperative period and overall patients' satisfaction with treatment, increases time spent in hospital, and opioid analgesics consumption [2, 4]. The pain after hemorrhoidectomy (HE) and other anorectal surgery depends on anal sphincter and puborectal muscles spasm, the type of intra- and postoperative anesthesia, wound healing, surgical technique, stool type, and patient's subjective perception [5, 6]. According to pain management international guidelines, the target level of postoperative pain should be 3–4 or less visual analog score (VAS) points [7].

Preemptive analgesia is aimed to prevent pain after surgery and affects several points of the "pain" cascade [2]. The most common non-narcotic analgesics that block peripheral pain perception include NSAIDs, corticosteroids, and acetylsalicylic acid. Non-narcotic drugs that inhibit central sensitization include ketamine, acetaminophen, and some anticonvulsants (in particular, gabapentin). So multimodal analgesia could theoretically reduce pain to a minimum due to blocking all kinds of pain receptors. However, the preoperative use of even one type of analgesic contributes to pain relief and to the opioid consumption decrease after the intervention [8–11].

One of the most commonly prescribed painkillers is non-steroidal anti-inflammatory drugs (NSAIDs), which have an adequate analgesic effect and even in acute pain can reduce the opioid dose by 18.3% [12]. Regarding ketoprofen as an analgesic drug, none of the published studies describes anorectal surgery. However, ketoprofen is widely used in spinal, orthopedic, general, dental, and children surgery with successful outcomes [13–16]. It also showed significant efficacy as preemptive analgesia compared to the other drugs [17]. Ketoprofen is approved for use as an analgesic for the treatment of mild to severe pain in the postoperative period in total daily doses up to 300 mg; the recommended initial dose is 25 to 100 mg (the dosages are intended for people without acute or decompensated concomitant diseases or non-pregnant women) [18]. Comparing ketoprofen to acetaminophen, the first one showed a significantly better anti-inflammatory effect on the 3rd and 6th day after surgery and lower pain intensity [19].

Therefore adequate postoperative analgesia requires the multimodal approach including opioid analgesics, nonsteroidal anti-inflammatory drugs (NSAIDs), acetaminophen, and local anesthetics. But there is no current standard on preemptive analgesia for adult patients with symptomatic hemorrhoids grade III–IV undergoing Milligan-Morgan hemorrhoidectomy without previous history of anorectal interventions and decompensated somatic diseases, although several studies report some regimens that seem to be effective.

## Methods

### Objective

The aim is to assess the efficacy of preemptive analgesia with Ketoprofen 100 mg 2 h before procedure per os with spinal anesthesia to decrease postoperative pain according to visual analog scale and to reduce the opioids and other analgesics consumption.

### Study design and setting

This is a prospective, randomized, double-blind, unicenter, superiority, parallel-group 2-arm study with 1:1 allocation ratio conducted in the surgical department of Moscow Research Educational Center of the Lomonosov Moscow State University. It is in the recruitment stage. We are anticipating 144 patients of all genders from 18 to 75 years old in total who come to the clinic. Thus, in this case, a uni-center design can assure sufficient patient recruitment.

### Eligibility criteria

Every patient included in the study must meet the following criteria:

- Symptomatic hemorrhoids grade III–IV

- Planned surgery: Milligan-Morgan hemorrhoidectomy

Patients who had contraindication or technical inability to perform subarachnoid anesthesia or other somatic diseases (history of peptic ulcer diseases and gastrointestinal bleeding and other acute or decompensated organ pathology) or refused to participate and pregnant women are not included. The written voluntary informed consent to participate (Additional file 3) is obtained from all eligible patients before randomization.

## Interventions

### *Preoperative preparation*

Two hours before the procedure, every patient receives a medication. The research group receives ketoprofen 100 mg per os; the control group receives a placebo.

### *Surgical technique*

Under spinal anesthesia, the patient is placed in a modified lithotomy position on the back, with legs spread apart on supports. During anesthesia, the patient is sitting. The needle is inserted between L3 and L4 space or L4 and L5 space. We usually use a bupivacaine solution 5 mg/ml for spinal anesthesia. No adjuvant medicaments are used. The operative field is treated with an antiseptic solution twice and draped. A complex of external and internal hemorrhoids or internal hemorrhoids only is excised with monopolar electrocautery or bipolar electro-surgery device. Hemorrhoid pedicle is tied with an absorbable polyfilament suture, and then the pedicle is crossed by monopolar or bipolar electrocautery. The wound is not sutured. One, two, or three nodes can be removed per procedure depending on the number of the enlarged hemorrhoidal nodes.

### *Postoperative period*

After the procedure, all patients receive the standard analgesics scheme. It includes ketoprofen (the average dose, 100 mg per day; the maximum dose, 200 mg per day), paracetamol (at the maximum dose of 1000 mg per day), and local anesthetics with lidocaine before defecation. Patients are usually discharged on the 4–5th day after surgery when their pain level is below 5 points according to VAS. After discharge, a standard analgesics scheme mentioned above is recommended to all patients. If the pain level reaches 5–6 points according to VAS on the 5th day after the procedure, patients continue inpatient treatment and may receive an opioid analgesic.

### *Baseline characteristics*

The list of the baseline characteristics that we collect to evaluate the population:

- Gender
- Age
- Hemorrhoid stage
- The number of enlarged hemorrhoids
- Hemorrhoid symptoms

## The main outcome measures

The trial is conducted to evaluate the primary outcome: the quantity of the opioid administration intake per day during the first week postoperatively that is necessary to hold pain level no more than 3–4 VAS points in every patient. We use tramadol as an opioid analgesic in a single standard dose of 100 mg for patients with persistent severe pain syndrome in our clinic. The study also assesses the following secondary outcomes: (1) the pain severity before and after defecation according to VAS on the 6, 12, and 24 h after the procedure, then 2 times per day up to 7th postoperative day; (2) duration and (3) frequency of other analgesics intake (systemically and topically) during the first week postoperatively; (4) readmission rate and (5) overall quality of life on the 7th and 30th days; (6) time from the procedure to return to work; and (7) the complications rate (i.e. bleeding, retention of urine, infectious complications) in the early postoperative period (30 days after procedure). The pain severity, duration and frequency of other analgesics intake during the first week postoperatively will be assessed using “Postoperative protocol for pain management” (Additional file 2). The overall quality of life will be assessed with a patient-reported questionnaire Short Form 36 (SF-36). A total score in each of the 8 sections will be calculated and transformed into a 0–100 scale with a score of zero equivalents to maximum disability and a score of 100 equivalents to no disability. Other secondary outcomes will be updated during visits or by phone on the 7 and 30 days after the surgery.

All patients are scheduled to return to the ambulatory clinic on the 7 and 30 days after the surgery. During these visits, postoperative data is collected and digital rectal examination is performed. If a patient fails to follow up, the researcher may contact the patient by all means available (phone, email, or mail) to ascertain whether the patient has had any complications and/or adverse events that were treated at another hospital. If the researcher is unsuccessful in contacting the patient, the patient will be considered as lost to follow-up.

### *Participant timeline*

For the schedule of enrolment, interventions, and assessments, see Table 1.

### *Sample size*

Considering that this is a superiority study, the sample size was calculated using a 1-sided Blackwelder test.

**Table 1** Schedule of enrollment, interventions, and assessments

Timepoint	Enrollment	Randomization	Interventions	Follow-up (1)*	Follow-up (1)**	Follow-up (3)***
<b>Eligibility criteria</b>	●					
<b>Informed consent</b>	●					
<b>Physical examination</b>	●					
<b>Demographic characteristics</b>	●					
<b>Allocation</b>		●				
<b>Drug administration</b>			●			
<b>Procedure</b>			●			
<b>Primary outcome assessment</b>				●	●	
<b>Secondary outcomes assessment</b>				●	●	●

\*Follow-up (1)—the first postoperative day  
 \*\*Follow-up (2)—the first week after the procedure  
 \*\*\*Follow-up (3)—postoperative day 30

According to published data, the incidence of opioid intake after hemorrhoidectomy varies from 20 to 30% [7]. The expected incidence of opioid intake after hemorrhoidectomy with preemptive analgesia is not more than 10%. The purpose of this study is to show that the opioids intake in patients with preemptive analgesia is lower than without it. Considering that  $\alpha = 0.05$ , the statistical power of the study is 80%, the patients are randomized into 2 groups with 1:1 allocation ratio, the noninferiority margin  $D = 5\%$ , and the required sample size is 144 patients (72 patients in each of the 2 groups).

**Recruitment**

All patients diagnosed with HD II–III stage will be considered for this study.

**Assignment of interventions**


Participants will be randomly assigned to either control or experimental group with a 1:1 allocation ratio using cluster randomization with a computerized random number generator. All subjects will be allocated any

interventions. The experimental group receives a tablet with 100 mg ketoprofen, the control one receives a tablet containing starch per os 2 h before surgery (72 participants per arm) (see Fig. 1). The investigator who does not operate generates the allocation sequence, enrolls participants, obtains informed consent, and assigns participants to interventions. The surgeon and the anesthesiology team are blinded.

All relevant data from patient charts except patients' names will be transferred into an electronic case report form (eCRF). The eCRF should contain results of all the screening procedures, including patient history and demographics, imaging studies, filled-out questionnaires, operation notes, and postoperative rounds during a patient stay in the surgical ward.

**Data collection, management, and analysis**

All data will be collected prospectively using eCRFs designed for this trial. The reasons for withdrawal will be documented. The investigator will attempt to contact each participant at least 3 times during each follow-up



**Fig. 1** Trial flow diagram

window before declaring them lost for observation. The study exit form will be recorded in the eCRF. All prior data will be analyzed within the research.

All patients will receive clarifications of all the study procedures and will be able to discuss them with the primary investigator. All patient data will be handled according to the principles of doctor–patient confidentiality; the subjects will be anonymized and analyzed with individual identifier numbers transcribed into eCRF.

Data analysis will be performed by one of the researchers who is blind to the randomization by using Statistical Product and Service Solutions (SPSS) Statistics for Windows, version 20.0 (IBM Corp., Armonk, NY, USA). Continuous variables will be presented as mean and standard deviation, and categorical variables will be presented as frequency or percentage.

### Statistics

The ITT analysis included all randomized patients treated for at least 1 week, who were evaluated for efficacy at baseline and at least once at subsequent evaluation points during the study. The PP-analysis will exclude from the full analysis all the non-compliant patients, patients who received incomplete treatment, and patients who refused further treatment after randomization.

Quantitative variables are described as means with standard deviations, medians, range, or interquartile range as appropriate. Categorical variables are described in absolute numbers and percentages. The statistical analysis of the quantitative variables, with independent groups, is performed with the parametric Student's *t* test, providing that its conditions for application are met. Otherwise, the non-parametric Mann-Whitney *U* test is used. Statistical analysis for categorical variables is performed using the Pearson  $\chi^2$  test or the Fisher exact test. Specifically, the above methods are used to compare the two groups in terms of baseline characteristics in order to assess whether the randomization has been effective. Ninety-five percent confidence intervals for the ratio will be reported. In all cases, the level of significance will be the usual 5% ( $\alpha = 0.05$ ).

This study has a low risk of selection bias due to the use of a computer random number generator. Central allocation and the use of sequentially numbered drug containers of identical appearance allow to achieve a low risk of deviations. Performance bias may be considered as having an average risk due to blinding only participants of the study, the personnel is not blinding. Blinding of outcome assessment is ensured, so detection bias is unlikely. Attrition bias could probably appear because of the difficulty of collecting data from patients outside the hospital or the refusal of future treatment and visiting the doctor during follow-up.

### Data monitoring

There is no data monitoring committee designated for this trial. Any adverse and serious adverse events will be immediately reported to the principal investigator and the primary sponsor. Participants' names and collected data are subject to medical confidentiality. In cases of withdrawal, collection of data will cease but not be erased. A logistics database with a patient's complete ID will be used and kept within a separate password-protected system from the results database with all study information. At the end of the trial, the original peri-operative package and database will be archived by the principal investigator, who is responsible for providing data to the trial investigators.

### Ethical approval

This study is conducted in accordance to the principles of the Declaration of Helsinki (Fortaleza, 2013). The study protocol is approved by the Local Ethics committee of Sechenov University (see Additional file 1).

### Protocol amendments

Any protocol amendments that may influence the conduct of the study will be communicated to the local ethics committee and study director and will be uploaded to clinical trials.

### Consent or assent

A member of the research team will obtain the consent form. All participants will be able to address their questions about the study to one of the members of the research team.

### Confidentiality

All patient data will be secured at the study site. No one apart from the members of the research team will have access to any patient data, including anonymized eCRFs with a coded ID, as well as filled out questionnaires.

### Declaration of interests

The authors declare they have no competing interests.

### Access to data

No one apart from the members of the research team will have access to the final trial dataset.

### Ancillary and post-trial care

If a study participant receives a complication or suffers in any way, he will be provided with a full recovery in our clinical center free of charge.

### Dissemination policy

Trial results will be e-mailed to all participants of the trial. Trial results will be disseminated to healthcare



professionals via publication in a peer-reviewed scientific journal and by mass media, as well as conference papers to inform the public and stakeholders, and will be uploaded to the primary registry. We have no intention of granting public access to the full protocol, participant-level dataset, and statistical code.

## Discussion

Multimodality pain management for anorectal surgery has been shown to improve pain control and decrease opioid requirements in several studies. Preoperative oral acetaminophen and gabapentin admission followed by intravenous ketamine in the early postoperative period resulted to significantly less pain level postoperatively and decreased narcotics intake [2]. Place RJ [20] showed a significant decrease of postoperative analgesics requirement in addition to reducing voiding problems by systemic admission of ketorolac with local anesthetics in patients undergoing anorectal surgery.

Several studies showed that ketoprofen significantly reduced opioid consumption (by 33%) and improved postoperative analgesia after spinal and abdominal surgery [13–16]. Patients with acute pain who have undergone dental surgery get a meaningful pain relief, a faster onset of effect, the highest peak effect, and the longest duration of action using ketoprofen than other NSAIDs and analgesics [18, 21]. In all mentioned above studies, adverse effects related to ketoprofen were minor and infrequent.

We hope to prove the efficacy of multimodal analgesia including preemptive one for patients undergoing excisional hemorrhoidectomy that will help to hold postoperative pain levels no more than 3–4 points on VAS with minimal consumption of opioid analgesics.

## Trial status

Registered at [clinicaltrials.gov](https://clinicaltrials.gov) number ID NCT04361695, date of registration: April 24, 2020, version 1.0. Recruitment began in April 2020 and is expected to be completed by the end of December 2021.

## Abbreviations

VAS: Visual analog score; NSAIDs: Nonsteroidal anti-inflammatory drugs; eCRF: Electronic case report form

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13063-022-06107-0>.

**Additional file 1.** SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents.

**Additional file 2.** Postoperative protocol for pain management.

**Additional file 3.** Written form of informed consent of the patient.

## Acknowledgements

The research team expresses gratitude to Moscow State University, Sechenov University administration for supporting this study.

## Authors' contributions

Each author has made substantial contributions: TG, AD, and EK to the conception design of the work; HL to the acquisition; EK and SM to analysis; HL to the interpretation of data; AD and SM to the creation of new software used in the work; PT and TG have drafted the work or substantively revised it. Study Principal Investigator: PT. The authors have read and approved the final manuscript.

## Funding

The study has no funding.

## Availability of data and materials

Not applicable.

## Declarations

### Ethics approval and consent to participate

The study protocol is approved by the Local Ethics committee of Sechenov University. Prior to randomization, written informed consent is obtained from all patients.

### Consent for publication

Additional file 3.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>Surgical Department, Moscow Research Educational Center of the Lomonosov Moscow State University, Moscow 101000, Russia. <sup>2</sup>Department: Clinic of Colorectal and Minimally Invasive Surgery, Sechenov First Moscow State Medical University, Moscow 101000, Russia. <sup>3</sup>Section of Colorectal Surgery, Department of Surgery, New York Medical College, Westchester Medical Center, Taylor Pavilion, Suite D-361, 100 Woods Road, Valhalla, NY 10595, USA.

Received: 7 December 2020 Accepted: 15 February 2022

Published online: 27 June 2022

## References

1. Sneddon LU. Comparative physiology of nociception and pain. *Physiology* (Bethesda). 2018;33(1):63–73. <https://doi.org/10.1152/physiol.00022.2017> PMID: 29212893.
2. Van Backer JT, Jordan MR, Leahy DT, Moore JS, Callas P, Dominick T, et al. Preemptive analgesia decreases pain following anorectal surgery: a prospective, randomized, double-blinded, placebo-controlled trial. *Dis Colon Rectum*. 2018;61(7):824–9. <https://doi.org/10.1097/DCR.0000000000001069>.
3. Gallo G, Martellucci J, Sturiale A, Clerico G, Milito G, Marino F, et al. Consensus statement of the Italian Society of Colorectal Surgery (SICCR): management and treatment of hemorrhoidal disease. *Tech Coloproctol*. 2020;24(2):145–64. <https://doi.org/10.1007/s10151-020-02149-1>.
4. Penprase B, Brunetto E, Dahmani E, Forthoffer JJ, Kapoor S. The efficacy of preemptive analgesia for postoperative pain control: a systematic review of the literature. *AORN J*. 2015;101(1):94–105.e8. <https://doi.org/10.1016/j.aorn.2014.01.030>.
5. Huang YJ, Chen CY, Chen RJ, Kang YN, Wei PL. Topical diltiazem ointment in post-hemorrhoidectomy pain relief: a meta-analysis of randomized controlled trials. *Asian J Surg*. 2018;41(5):431–7. <https://doi.org/10.1016/j.asjsur.2017.06.002>.
6. Liu JW, Lin CC, Kiu KT, Wang CY, Tam KW. Effect of glyceryl trinitrate ointment on pain control after hemorrhoidectomy: a meta-analysis of randomized controlled trials. *World J Surg*. 2016;40(1):215–24. <https://doi.org/10.1007/s00268-015-3344-6>.
7. Frasco PE, Sprung J, Trentman TL. The impact of the joint commission for accreditation of healthcare organizations pain initiative on perioperative opiate consumption and recovery room length of stay. *Anesth Analg*. 2005; 100(1):162–8.
8. Poylin V, Quinn J, Messer K, Nagle D. Gabapentin significantly decreases posthemorrhoidectomy pain: a prospective study. *Int J Colorectal Dis*. 2014; 29(12):1565–9. <https://doi.org/10.1007/s00384-014-2018-4>.

9. Wu JI, Lu SF, Chia YY, Yang LC, Fong WP, Tan PH. Sevoflurane with or without antiemetic prophylaxis of dexamethasone in spontaneously breathing patients undergoing outpatient anorectal surgery. *J Clin Anesth.* 2009;21(7):469–73. <https://doi.org/10.1016/j.jclinane.2008.11.007>.
10. Khalili G, Janghorbani M, Saryazdi H, Emaminejad A. Effect of preemptive and preventive acetaminophen on postoperative pain score: a randomized, double-blind trial of patients undergoing lower extremity surgery. *J Clin Anesth.* 2013;25(3):188–92. <https://doi.org/10.1016/j.jclinane.2012.09.004>.
11. Richman IM. Use of Toradol in anorectal surgery. *Dis Colon Rectum.* 1993; 36(3):295–6. <https://doi.org/10.1007/BF02053515>.
12. Chen JY, Ko TL, Wen YR, Wu SC, Chou YH, Yien HW, et al. Opioid-sparing effects of ketorolac and its correlation with the recovery of postoperative bowel function in colorectal surgery patients: a prospective randomized double-blinded study. *Clin J Pain.* 2009;25(6):485–9. <https://doi.org/10.1097/AJP.0b013e31819a506b>.
13. Aubrun F, Langeron O, Heitz D, Coriat P, Riou B. Randomised, placebo-controlled study of the postoperative analgesic effects of ketoprofen after spinal fusion surgery. *Acta Anaesthesiol Scand.* 2000;44(8):934–9. <https://doi.org/10.1034/j.1399-6576.2000.440807.x>.
14. Fletcher D, Nègre I, Barbin C, François A, Carreres C, Falgueirettes C, et al. Postoperative analgesia with i.v. propacetamol and ketoprofen combination after disc surgery. *Can J Anaesth.* 1997;44(5 Pt 1):479–85. <https://doi.org/10.1007/BF03011934>.
15. Hiller A, Meretoja OA, Korpela R, Piiparinen S, Taivainen T. The analgesic efficacy of acetaminophen, ketoprofen, or their combination for pediatric surgical patients having soft tissue or orthopedic procedures. *Anesth Analg.* 2006;102(5):1365–71. <https://doi.org/10.1213/01.ane.0000204278.71548.bf>.
16. Oberhofer D, Skok J, Nesek-Adam V. Intravenous ketoprofen in postoperative pain treatment after major abdominal surgery. *World J Surg.* 2005;29(4):446–9. <https://doi.org/10.1007/s00268-004-7612-0>.
17. Velásquez GC, Santa Cruz LA, Espinoza MA. Ketoprofen is more effective than diclofenac after oral surgery when used as a preemptive analgesic: a pilot study. *J Oral Facial Pain Headache.* 2014;28(2):153–8. <https://doi.org/10.11607/ofph.1200>.
18. Gaskell H, Derry S, Wiffen PJ, Moore RA. Single dose oral ketoprofen or dextketoprofen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2017;5(5):CD007355. <https://doi.org/10.1002/14651858.CD007355.pub3> PMID: 28540716; PMCID: PMC6481461.
19. Björnsson GA, Haanaes HR, Skoglund LA. Ketoprofen 75 mg qid versus acetaminophen 1000 mg qid for 3 days on swelling, pain, and other postoperative events after third-molar surgery. *J Clin Pharmacol.* 2003;43(3): 305–14. <https://doi.org/10.1177/0091270002250603>.
20. Place RJ, Coloma M, White PF, Huber PJ, Van Vlymen J, Simmang CL. Ketorolac improves recovery after outpatient anorectal surgery. *Dis Colon Rectum.* 2000;43(6):804–8. <https://doi.org/10.1007/bf02238018>.
21. Cooper SA. Ketoprofen in oral surgery pain: a review. *J Clin Pharmacol.* 1988;28(5):540–6. <https://doi.org/10.1002/j.1552-4604.1988.tb05976.x>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

