

**CORRECTION**

**Open Access**




# Correction to: Comorbidity in gout at the time of first diagnosis: sex differences that may have implications for dosing of urate lowering therapy

Panagiota Drivelegka<sup>1\*</sup>, Valgerdur Sigurdardottir<sup>1,2</sup>, Anna Svärd<sup>2</sup>, Lennart T. H. Jacobsson<sup>1</sup> and Mats Dehlin<sup>1</sup>

**Correction**

Unfortunately, after publication of this article [1], it was noticed that Fig. 2 is incorrect. The correct Fig. 2 can be seen below.



\* Correspondence: [panagiota.drivelegka@vgregion.se](mailto:panagiota.drivelegka@vgregion.se)

<sup>1</sup>Department of Rheumatology and Inflammation Research, Sahlgrenska Academy, University of Gothenburg, Grona Straket 12, Sahlgrenska University Hospital, 413 45 Gothenburg, Sweden



**Author details**

<sup>1</sup>Department of Rheumatology and Inflammation Research, Sahlgrenska Academy, University of Gothenburg, Grona Straket 12, Sahlgrenska University Hospital, 413 45 Gothenburg, Sweden. <sup>2</sup>Centre of Clinical Research Dalarna, Falun, Sweden.

Received: 13 August 2018 Accepted: 15 August 2018

Published online: 10 September 2018

**Reference**

1. Drivelegka P, Sigurdardottir V, Svärd A, Jacobsson LTH, Dehlin M. Comorbidity in gout at the time of first diagnosis: sex differences that may have implications for dosing of urate lowering therapy. *Arthritis Res Ther.* 2018;20:108. <https://doi.org/10.1186/s13075-018-1596-x>