

RESEARCH

Open Access



Multiple positive solutions for a fractional elliptic system with critical nonlinearities

Haining Fan*

*Correspondence:
fanhaining888@163.com
School of Mathematics, China
University of Mining and
Technology, Xuzhou, 221116, China

Abstract

In this paper, we study the multiplicity results of positive solutions for a fractional elliptic system involving both concave-convex and critical growth terms. With the help of the Nehari manifold and the Ljusternik-Schnirelmann category, we prove that the problem admits at least $\text{cat}(\Omega) + 1$ distinct positive solutions.

MSC: 35J50; 35J57; 35J66

Keywords: Nehari manifold; fractional elliptic system; Ljusternik-Schnirelmann category; multiple positive solutions

1 Introduction and the main result

In this paper, we are concerned with the number of positive solutions of the fractional elliptic system:

$$(E_{\lambda, \mu}) \quad \begin{cases} (-\Delta)^{\frac{s}{2}} u = \lambda |u|^{q-2} u + \frac{\alpha}{\alpha+\beta} |u|^{\alpha-2} u |v|^{\beta} & \text{in } \Omega, \\ (-\Delta)^{\frac{s}{2}} v = \mu |v|^{q-2} v + \frac{\beta}{\alpha+\beta} |u|^{\alpha} |v|^{\beta-2} v & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded set in \mathbb{R}^N with smooth boundary, $N > s$ with $s \in (0, 2)$ fixed, $1 < q < 2$, $\lambda, \mu > 0$, $\alpha, \beta > 1$ satisfy $\alpha + \beta = 2_s^* = \frac{2N}{N-s}$, 2_s^* is the fractional Sobolev critical exponent, and $(-\Delta)^{\frac{s}{2}}$ is the fractional Laplacian. These types of operators are the infinitesimal generators of Lévy stable diffusion process and arise in anomalous diffusions in plasmas, flames propagation and chemical reactions in Liquids, population dynamics, geophysical fluid dynamics, and American options in finance; see [1, 2].

In recent years, a great deal of attention has been focused on studying of problems involving fractional Sobolev spaces and corresponding nonlocal equations, both from a pure mathematical point of view and for concrete applications. We refer to [3–7] for the subcritical case and to [8–12] for the critical case. In particular, set $\alpha + \beta = p \leq 2_s^*$, $\lambda = \mu$, and $u = v$, $(E_{\lambda, \mu})$ reduces to the following fractional elliptic equation with concave-convex nonlinearities:

$$(E_{\lambda}) \quad \begin{cases} (-\Delta)^{\frac{s}{2}} u = \lambda |u|^{q-2} u + |u|^{p-2} u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

Goyal and Sreenadh [13] studied the existence and multiplicity of non-negative solutions to (E_λ) . Moreover, by Nehari manifold and fibering maps, Chen and Deng [9] obtained the existence of multiple solutions to (E_λ) for subcritical case and critical case. For the fractional Laplacian system with concave-convex nonlinearities, He *et al.* [14] proved that $(E_{\lambda,\mu})$ permits at least two positive solutions when the pair of parameters (λ, μ) belongs to a certain subset of \mathbb{R}^2 . Similar results were taken by Chen and Deng [15]. The tool of them is the decomposition of the Nehari manifold.

Motivated by the results mentioned above, the purpose of this article is to get a better information on the number of positive solutions of $(E_{\lambda,\mu})$, for $\lambda, \mu > 0$ small enough, via the tools of the variational theory and the Ljusternik-Schnirelmann category theory. We refer the reader to [16–21] for similar results to $(E_{\lambda,\mu})$ for Laplacian operator. Our main result can be stated as follows.

Theorem 1.1 *There exists $\Lambda_* > 0$ such that if $\lambda, \mu \in (0, \Lambda_*)$, $(E_{\lambda,\mu})$ has at least $\text{cat}(\Omega) + 1$ distinct positive solutions. Here $\text{cat}(\Omega)$ denotes the Ljusternik-Schnirelmann category of Ω in itself.*

Remark 1.1 If Ω is a general domain, $\text{cat}(\Omega) \geq 1$, and Theorem 1.1 is the main result of [14, 15].

Remark 1.2 Concerning regularity, one can get an *a priori* estimate for the solutions to $(E_{\lambda,\mu})$ and hence obtain, as in [22], Proposition 5.2, $u, v \in C^\infty(\overline{\Omega})$ for $s = 1$, $u, v \in C^{0,s}(\overline{\Omega})$ if $0 < s < 1$ and $u, v \in C^{1,s-1}$ if $1 < s < 2$.

This paper is organized as follows: In Section 2, we introduce some notations and preliminaries. In Section 3, we give some technical results which are crucial to the proof of Theorem 1.1. In Section 4, we give the proof of Theorem 1.1.

2 Notations and preliminaries

In this section, we collect preliminary facts for future reference. First of all, let us write the standard notations which we will use in this paper. We denote the upper half-space in \mathbb{R}_+^{N+1} by

$$\mathbb{R}_+^{N+1} := \{(x, y); (x_1, x_2, \dots, x_N, y) \in \mathbb{R}^{N+1}, y > 0\}.$$

Denote the half cylinder with base Ω by $C_\Omega = \Omega \times (0, \infty) \subset \mathbb{R}_+^{N+1}$ and its lateral boundary by $\partial_L C_\Omega = \partial\Omega \times [0, \infty)$. We shall use $C(C_i, i = 1, 2, \dots)$ to denote any positive constant.

Let φ_j, λ_j be the eigenfunctions and eigenvectors of $-\Delta$ in Ω with zero Dirichlet boundary data. The fractional Laplacian $(-\Delta)^{\frac{s}{2}}$ is defined in the space of functions

$$H_0^{\frac{s}{2}}(\Omega) := \left\{ u = \sum_{j=1}^{\infty} a_j \varphi_j \in L^2(\Omega); \|u\|_{H_0^{\frac{s}{2}}(\Omega)} = \left(\sum_{j=1}^{\infty} a_j^2 \lambda_j^{\frac{s}{2}} \right)^{\frac{1}{2}} < \infty \right\},$$

and $\|u\|_{H_0^{\frac{s}{2}}(\Omega)} = \|(-\Delta)^{\frac{s}{4}} u\|_{L^2(\Omega)}$. The dual space $H^{-\frac{s}{2}}(\Omega)$ is defined in the standard way as well as the inverse operator $(-\Delta)^{-\frac{s}{2}}$.

Definition 2.1 We say that $(u, v) \in H_0^{\frac{s}{2}}(\Omega) \times H_0^{\frac{s}{2}}(\Omega)$ is a solution of $(E_{\lambda,\mu})$ if the identity

$$\begin{aligned} & \int_{\Omega} (-\Delta)^{\frac{s}{4}} u (-\Delta)^{\frac{s}{4}} \varphi_1 + (-\Delta)^{\frac{s}{4}} v (-\Delta)^{\frac{s}{4}} \varphi_2 \, dx \\ &= \int_{\Omega} (\lambda |u|^{q-2} u \varphi_1 + \mu |v|^{q-2} v \varphi_2) \, dx + \frac{\alpha}{\alpha + \beta} \int_{\Omega} |u|^{\alpha-2} u |v|^{\beta} \varphi_1 \, dx \\ & \quad + \frac{\beta}{\alpha + \beta} \int_{\Omega} |u|^{\alpha} |v|^{\beta-2} v \varphi_2 \, dx \end{aligned}$$

holds for all $(\varphi_1, \varphi_2) \in H_0^{\frac{s}{2}}(\Omega) \times H_0^{\frac{s}{2}}(\Omega)$.

Associated with $(E_{\lambda,\mu})$ we consider the energy functional

$$\begin{aligned} J_{\lambda,\mu}(u, v) &:= \frac{1}{2} \int_{\Omega} (|(-\Delta)^{\frac{s}{4}} u|^2 + |(-\Delta)^{\frac{s}{4}} v|^2) \, dx \\ & \quad - \frac{1}{q} \int_{\Omega} (\lambda |u|^q + \mu |v|^q) \, dx - \frac{1}{2_s^*} \int_{\Omega} |u|^{\alpha} |v|^{\beta} \, dx. \end{aligned}$$

This functional is well defined in $H_0^{\frac{s}{2}}(\Omega) \times H_0^{\frac{s}{2}}(\Omega)$, and, moreover, the critical points of $J_{\lambda,\mu}$ correspond to weak solutions of $(E_{\lambda,\mu})$.

To treat the nonlocal problem $(E_{\lambda,\mu})$, we will study a corresponding extension problem, which allows us to investigate $(E_{\lambda,\mu})$ by studying a local problem via classical variational methods. We define the extension operator and fractional Laplacian for functions in $H_0^{\frac{s}{2}}(\Omega)$.

Definition 2.2 Given a function $u \in H_0^{\frac{s}{2}}(\Omega)$, we define its s -harmonic extension $\omega = E_s(u)$ to the cylinder C_{Ω} as a solution to the problem

$$\begin{cases} \operatorname{div}(y^{1-s} \nabla \omega) = 0 & \text{in } C_{\Omega}, \\ \omega = 0 & \text{on } \partial_L C_{\Omega}, \\ \omega = u & \text{on } \Omega \times \{0\}, \end{cases}$$

and

$$(-\Delta)^{\frac{s}{2}} u(x) = -K_s \lim_{y \rightarrow 0^+} y^{1-s} \frac{\partial \omega}{\partial y}(x, y),$$

where K_s is a normalization constant.

The extension function $\omega(x, y)$ belongs to the space $H_{0,L}^s(C_{\Omega}) = \overline{C_0^{\infty}(\Omega \times [0, \infty))}$, with

$$\|\omega\|_{H_{0,L}^s(C_{\Omega})} = \left(K_s \int_{C_{\Omega}} y^{1-s} |\nabla \omega|^2 \, dx \, dy \right)^{\frac{1}{2}}.$$

The extension operator is an isometry between $H_0^{\frac{s}{2}}(\Omega)$ and $H_{0,L}^s(C_{\Omega})$, namely,

$$\|\omega\|_{H_{0,L}^s(C_{\Omega})} = \|u\|_{H_0^{\frac{s}{2}}(\Omega)}, \quad \forall u \in H_0^{\frac{s}{2}}(\Omega). \tag{2.1}$$

With this extension, we can transform $(E_{\lambda,\mu})$ into the following local problem:

$$(\widehat{E}_{\lambda,\mu}) \begin{cases} -\operatorname{div}(y^{1-s}\nabla\omega_1) = 0, & -\operatorname{div}(y^{1-s}\nabla\omega_2) = 0 & \text{in } C_\Omega, \\ \omega_1 = \omega_2 = 0 & & \text{on } \partial_L C_\Omega, \\ \frac{\partial\omega_1}{\partial\nu^s} = \lambda|\omega_1|^{q-2}\omega_1 + \frac{\alpha}{\alpha+\beta}|\omega_1|^{\alpha-2}\omega_1|\omega_2|^\beta & & \text{on } C_\Omega \times \{0\}, \\ \frac{\partial\omega_2}{\partial\nu^s} = \lambda|\omega_2|^{q-2}\omega_2 + \frac{\beta}{\alpha+\beta}|\omega_1|^\alpha|\omega_2|^{\beta-2}\omega_2 & & \text{on } C_\Omega \times \{0\}, \\ \omega_1 = u, \quad \omega_2 = v & & \text{on } C_\Omega \times \{0\}, \end{cases}$$

where

$$\frac{\partial\omega_i}{\partial\nu^s} := -K_s \lim_{y \rightarrow 0^+} y^{1-s} \frac{\partial\omega_i}{\partial y}, \quad i = 1, 2.$$

In the following, we will study $(\widehat{E}_{\lambda,\mu})$ in the framework of the Sobolev space $H = H_{0,L}^s(C_\Omega) \times H_{0,L}^s(C_\Omega)$ using the standard norm

$$\|(\omega_1, \omega_2)\|_H = \left(K_s \int_\Omega y^{1-s} (|\nabla\omega_1|^2 + |\nabla\omega_2|^2) \, dx \, dy \right)^{\frac{1}{2}}.$$

An energy solution to $(\widehat{E}_{\lambda,\mu})$ is a function $(\omega_1, \omega_2) \in H$ satisfying

$$\begin{aligned} & K_s \int_{C_\Omega} y^{1-s} \nabla\omega_1 \nabla\varphi_1 \, dx \, dy + K_s \int_{C_\Omega} y^{1-s} \nabla\omega_2 \nabla\varphi_2 \, dx \, dy \\ &= \int_{\Omega \times \{0\}} (\lambda|\omega_1|^{q-2}\omega_1\varphi_1 + \mu|\omega_2|^{q-2}\omega_2\varphi_2) \, dx \\ & \quad + \frac{\alpha}{\alpha+\beta} \int_{\Omega \times \{0\}} |\omega_1|^{\alpha-2}\omega_1|\omega_2|^\beta\varphi_1 \, dx + \frac{\beta}{\alpha+\beta} \int_{\Omega \times \{0\}} |\omega_1|^\alpha|\omega_2|^{\beta-2}\omega_2\varphi_2 \, dx \end{aligned}$$

for all $(\varphi_1, \varphi_2) \in H$.

If (ω_1, ω_2) satisfies $(\widehat{E}_{\lambda,\mu})$, then the trace $(u, v) = (\omega_1(\cdot, 0), \omega_2(\cdot, 0))$ is a solution of $(E_{\lambda,\mu})$. The converse is also true. Therefore, both formulations are equivalent.

The associated energy functional to $(\widehat{E}_{\lambda,\mu})$ is

$$\begin{aligned} I_{\lambda,\mu}(\omega_1, \omega_2) &= \frac{1}{2} \|(\omega_1, \omega_2)\|_H^2 - \frac{1}{q} \int_{\Omega \times \{0\}} (\lambda|\omega_1|^q + \mu|\omega_2|^q) \, dx \\ & \quad - \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta \, dx. \end{aligned}$$

Clearly, critical points of $I_{\lambda,\mu}$ in H correspond to critical points of $J_{\lambda,\mu}$ in $H_0^{\frac{s}{2}}(\Omega) \times H_0^{\frac{s}{2}}(\Omega)$.

In the following lemmas, we will list some relevant inequalities from [14, 15].

Lemma 2.1 *For every $1 \leq r \leq 2_s^*$, and every $\omega \in H_{0,L}^s(C_\Omega)$, we have*

$$\left(\int_{\Omega \times \{0\}} |\omega|^r \, dx \right)^{\frac{2}{r}} \leq C \int_{C_\Omega} y^{1-s} |\nabla\omega|^2 \, dx \, dy \tag{2.2}$$

for some positive constant C . Furthermore, the space $H_{0,L}^s(C_\Omega)$ is compactly embedded into $L^r(\Omega)$, for every $r < 2_s^*$.

Remark 2.1 When $r = 2_s^*$, the best constant is denoted by $S(s, N)$, that is,

$$S(s, N) := \inf_{\omega \in H_{0,L}^s(C_\Omega) \setminus \{0\}} \frac{\int_{C_\Omega} y^{1-s} |\nabla \omega|^2 dx dy}{\left(\int_{\Omega \times \{0\}} |\omega|^{2_s^*} dx\right)^{\frac{2}{2_s^*}}}. \tag{2.3}$$

It is not achieved in any bounded domain and, for all $\omega \in H^s(\mathbb{R}_+^{N+1})$,

$$S(s, N) \left(\int_{\mathbb{R}^N \times \{0\}} |\omega|^{2_s^*} dx\right)^{\frac{1}{2_s^*}} \leq \int_{\mathbb{R}_+^{N+1}} y^{1-s} |\nabla \omega|^2 dx dy, \tag{2.4}$$

$S(s, N)$ is achieved for $\Omega = \mathbb{R}^N$ by the functions ω_ε which are the s -harmonic extensions of

$$u_\varepsilon(x) := \frac{\varepsilon^{\frac{(N-s)}{2}}}{(\varepsilon^2 + |x|^2)^{\frac{(N-s)}{2}}}, \quad \varepsilon > 0, x \in \mathbb{R}^N. \tag{2.5}$$

The constant $S(s, N)$ given in (2.3) takes the exact value

$$S(s, N) = \frac{2\pi^{\frac{s}{2}} \Gamma(\frac{2-s}{2}) \Gamma(\frac{N+s}{2}) (\Gamma(\frac{N}{2}))^{\frac{s}{N}}}{\Gamma(\frac{s}{2}) \Gamma(\frac{N-s}{2}) (\Gamma(N))^{\frac{s}{N}}},$$

and it is achieved for $\Omega = \mathbb{R}^N$ by the functions $\omega_\varepsilon = E_s(u_\varepsilon)$.

We consider the following minimization problem:

$$S_{s,\alpha,\beta} := \inf_{(\omega_1, \omega_2) \in H \setminus \{(0,0)\}} \frac{\int_{C_\Omega} y^{1-s} (|\nabla \omega_1|^2 + |\nabla \omega_2|^2) dx dy}{\left(\int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx\right)^{\frac{2}{2_s^*}}}. \tag{2.6}$$

From [14], we establish a relationship between $S(s, N)$ and $S_{s,\alpha,\beta}$.

Lemma 2.2 For the constants $S(s, N)$ and $S_{s,\alpha,\beta}$ introduced in (2.3) and (2.6), we have

$$S_{s,\alpha,\beta} = \left(\left(\frac{\alpha}{\beta}\right)^{\frac{\beta}{\alpha+\beta}} + \left(\frac{\beta}{\alpha}\right)^{\frac{\alpha}{\alpha+\beta}} \right) S(s, N).$$

In particular, the constant $S_{s,\alpha,\beta}$ is achieved for $\Omega = \mathbb{R}^N$.

To proceed, we introduce the Nehari manifold of $I_{\lambda,\mu}$ by setting

$$N_{\lambda,\mu} = \{(\omega_1, \omega_2) \in H \setminus \{(0,0)\}; I'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) = 0\}.$$

This enables us to construct homotopies between Ω and certain levels of $I_{\lambda,\mu}$. Clearly, $(\omega_1, \omega_2) \in N_{\lambda,\mu}$ if and only if

$$\|(\omega_1, \omega_2)\|_H^2 = \int_{\Omega \times \{0\}} (\lambda |\omega_1|^q + \mu |\omega_2|^q) dx + \int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx.$$

On the Nehari manifold $N_{\lambda,\mu}$, from Lemma 2.1 and the Young inequality, we have

$$\begin{aligned}
 I_{\lambda,\mu}(\omega_1, \omega_2) &= \left(\frac{1}{2} - \frac{1}{2_s^*}\right) \|(\omega_1, \omega_2)\|_H^2 - \left(\frac{1}{q} - \frac{1}{2_s^*}\right) \int_{\Omega \times \{0\}} (\lambda|\omega_1|^q + \mu|\omega_2|^q) dx \\
 &\geq \left(\frac{1}{2} - \frac{1}{2_s^*}\right) \|(\omega_1, \omega_2)\|_H^2 - \left(\frac{1}{q} - \frac{1}{2_s^*}\right) (\lambda + \mu) C \|(\omega_1, \omega_2)\|_H^q \tag{2.7}
 \end{aligned}$$

$$\geq -(\lambda + \mu)^{2/(2-q)} C, \tag{2.8}$$

where C denotes positive constants (possibly different) independent of $(\omega_1, \omega_2) \in H$. Let

$$\begin{aligned}
 \psi_{\lambda,\mu}(\omega_1, \omega_2) &:= I'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) \\
 &= \|(\omega_1, \omega_2)\|_H^2 - \int_{\Omega \times \{0\}} (\lambda|\omega_1|^q + \mu|\omega_2|^q) dx - \int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx. \tag{2.9}
 \end{aligned}$$

Then, for $(\omega_1, \omega_2) \in N_{\lambda,\mu}$,

$$\psi'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) = (2 - q) \|(\omega_1, \omega_2)\|_H^2 - (2_s^* - q) \int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx \tag{2.10}$$

$$\begin{aligned}
 &= (2 - 2_s^*) \|(\omega_1, \omega_2)\|_H^2 \\
 &\quad + (2_s^* - q) \int_{\Omega \times \{0\}} (\lambda|\omega_1|^q + \mu|\omega_2|^q) dx. \tag{2.11}
 \end{aligned}$$

Similar to the method used in [14, 15], we split $N_{\lambda,\mu}$ into three parts:

$$\begin{aligned}
 N_{\lambda,\mu}^+ &= \{(\omega_1, \omega_2) \in N_{\lambda,\mu}; \psi'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) > 0\}; \\
 N_{\lambda,\mu}^0 &= \{(\omega_1, \omega_2) \in N_{\lambda,\mu}; \psi'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) = 0\}; \\
 N_{\lambda,\mu}^- &= \{(\omega_1, \omega_2) \in N_{\lambda,\mu}; \psi'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) < 0\}.
 \end{aligned}$$

In the sequel, we shall use Λ_* to denote different small parameters. Then we have the following results.

Lemma 2.3 *Suppose that (ω_1, ω_2) is a local minimizer for $I_{\lambda,\mu}$ on $N_{\lambda,\mu}$. Then, if $(\omega_1, \omega_2) \notin N_{\lambda,\mu}^0$, (ω_1, ω_2) is a critical point of $I_{\lambda,\mu}$.*

Lemma 2.4 *There exists $\Lambda_* > 0$ such that, for each $\lambda, \mu \in (0, \Lambda_*)$, we have $N_{\lambda,\mu}^0 = \emptyset$.*

By Lemma 2.4, for $\lambda, \mu \in (0, \Lambda_*)$, we write $N_{\lambda,\mu} = N_{\lambda,\mu}^+ \cup N_{\lambda,\mu}^-$ and define

$$\alpha_{\lambda,\mu}^+ = \inf_{(\omega_1, \omega_2) \in N_{\lambda,\mu}^+} I_{\lambda,\mu}(\omega_1, \omega_2); \quad \alpha_{\lambda,\mu}^- = \inf_{(\omega_1, \omega_2) \in N_{\lambda,\mu}^-} I_{\lambda,\mu}(\omega_1, \omega_2).$$

Set

$$t_{\max} = \left(\frac{(2 - q) \|(\omega_1, \omega_2)\|_H^2}{(2_s^* - q) \int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx} \right)^{\frac{1}{2_s^* - 2}} > 0.$$

Then we have the following result.

Lemma 2.5 For each $(\omega_1, \omega_2) \in H$ with $\int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx > 0$, there exist unique $0 < t^+ < t_{\max} < t^-$ such that $(t^+ \omega_1, t^+ \omega_2) \in N_{\lambda, \mu}^+$, $(t^+ \omega_1, t^+ \omega_2) \in N_{\lambda, \mu}^+$, $(t^- \omega_1, t^- \omega_2) \in N_{\lambda, \mu}^-$ and

$$I_{\lambda, \mu}(t^+ \omega_1, t^+ \omega_2) = \inf_{0 \leq t \leq t_{\max}} I_{\lambda, \mu}(t \omega_1, t \omega_2); \quad I_{\lambda, \mu}(t^- \omega_1, t^- \omega_2) = \sup_{t \geq 0} I_{\lambda, \mu}(t \omega_1, t \omega_2).$$

Lemma 2.6 If $\lambda, \mu \in (0, \Lambda_*)$, then

- (i) $\alpha_{\lambda, \mu}^+ < 0$,
- (ii) $\alpha_{\lambda, \mu}^- \geq \delta$ for some $\delta > 0$.

For the proofs of Lemmas 2.3-2.6, the reader is referred to [14, 15] for similar proofs.

Remark 2.2 From Lemmas 2.5 and 2.6, it is easy to know that if $(\omega_1, \omega_2) \in N_{\lambda, \mu}^-$,

$$\int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx > 0.$$

Next we establish that $I_{\lambda, \mu}$ satisfies the $(PS)_c$ -condition under some restriction on the level of $(PS)_c$ -sequences in the following.

Lemma 2.7 For each $\lambda, \mu \in (0, \Lambda_*)$, $I_{\lambda, \mu}$ satisfies the $(PS)_c$ -condition for $c \in (-\infty, \alpha_{\lambda, \mu}^+ + \frac{s}{2N}(K_s S_{s, \alpha, \beta})^{N/s})$.

Proof Let $\{(\omega_{1,n}, \omega_{2,n})\} \subset H$ be a $(PS)_c$ -sequence for $I_{\lambda, \mu}$ and $c \in (-\infty, \alpha_{\lambda, \mu}^+ + \frac{s}{2N}(K_s S_{s, \alpha, \beta})^{N/s})$. Note (2.7), it is easy to see that $\{(\omega_{1,n}, \omega_{2,n})\}$ is bounded in H . Thus, there exists a subsequence still denoted by $\{(\omega_{1,n}, \omega_{2,n})\}$ and $(\omega_1, \omega_2) \in H$ such that $(\omega_{1,n}, \omega_{2,n}) \rightharpoonup (\omega_1, \omega_2)$ weakly in H . Furthermore, we get

- $\int_{\Omega \times \{0\}} (\lambda |\omega_{1,n}|^q + \mu |\omega_{2,n}|^q) dx = \int_{\Omega \times \{0\}} (\lambda |\omega_1|^q + \mu |\omega_2|^q) dx + o(1)$;
- $\|(\omega_{1,n} - \omega_1, \omega_{2,n} - \omega_2)\|_H^2 = \|(\omega_{1,n}, \omega_{2,n})\|_H^2 - \|(\omega_1, \omega_2)\|_H^2 + o(1)$;
- $\int_{\Omega \times \{0\}} |\omega_{1,n} - \omega_1|^\alpha |\omega_{2,n} - \omega_2|^\beta dx$
 $= \int_{\Omega \times \{0\}} |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx - \int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx + o(1).$

Moreover, we can obtain $I'_{\lambda, \mu}(\omega_1, \omega_2) = 0$. Since $I_{\lambda, \mu}(\omega_{1,n}, \omega_{2,n}) = c + o(1)$ and $I'_{\lambda, \mu}(\omega_{1,n}, \omega_{2,n}) = o(1)$, we deduce that

$$\begin{aligned} & \frac{1}{2} \|(\omega_{1,n} - \omega_1, \omega_{2,n} - \omega_2)\|_H^2 - \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |\omega_{1,n} - \omega_1|^\alpha |\omega_{2,n} - \omega_2|^\beta dx \\ &= c - I_{\lambda, \mu}(\omega_1, \omega_2) + o(1) \end{aligned} \tag{2.12}$$

and

$$\begin{aligned} o(1) &= I'_{\lambda, \mu}(\omega_{1,n}, \omega_{2,n})(\omega_{1,n} - \omega_1, \omega_{2,n} - \omega_2) \\ &= (I'_{\lambda, \mu}(\omega_{1,n}, \omega_{2,n}) - I'_{\lambda, \mu}(\omega_1, \omega_2))(\omega_{1,n} - \omega_1, \omega_{2,n} - \omega_2) \\ &= \|(\omega_{1,n} - \omega_1, \omega_{2,n} - \omega_2)\|_H^2 - \int_{\Omega \times \{0\}} |\omega_{1,n} - \omega_1|^\alpha |\omega_{2,n} - \omega_2|^\beta dx + o(1). \end{aligned}$$

Now we may assume that

$$\begin{aligned} & \|(\omega_{1,n} - \omega_1, \omega_{2,n} - \omega_2)\|_H^2 \rightarrow l \quad \text{and} \\ & \int_{\Omega \times \{0\}} |\omega_{1,n} - \omega_1|^\alpha |\omega_{2,n} - \omega_2|^\beta dx \rightarrow l \quad \text{as } n \rightarrow \infty \end{aligned}$$

for some $l \in [0, +\infty)$.

Suppose $l \neq 0$. Using (2.6) and passing to the limit as $n \rightarrow \infty$, we have

$$l \geq K_s S_{s,\alpha,\beta} l^{\frac{2}{2^*}},$$

that is,

$$l \geq (K_s S_{s,\alpha,\beta})^{N/s}. \tag{2.13}$$

Then by (2.12)-(2.13) and $(\omega_1, \omega_2) \in N_{\lambda,\mu} \cup \{0\}$,

$$c = I_{\lambda,\mu}(\omega_1, \omega_2) + \left(\frac{1}{2} - \frac{1}{2^*}\right)l \geq \alpha_{\lambda,\mu}^+ + \frac{s}{2N}(K_s S_{s,\alpha,\beta})^{N/s},$$

which contradicts the definition of c . Hence $l = 0$, and the proof is completed. □

Lemma 2.8 *For $\lambda, \mu \in (0, \Lambda_*)$, the functional $I_{\lambda,\mu}$ has a minimizer $((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) \in N_{\lambda,\mu}^+$ and it satisfies:*

- (i) $I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) = \alpha_{\lambda,\mu}^+$;
- (ii) $((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+)$ is a positive solution of $(\widehat{E}_{\lambda,\mu})$;
- (iii) $I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) \rightarrow 0$ as $\lambda, \mu \rightarrow 0$;
- (iv) $\lim_{\lambda,\mu \rightarrow 0} \|((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+)\|_H = 0$.

Proof (i)-(ii) are consequences of [14]. It follows from (2.8) and Lemma 2.6 that

$$0 > I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) \geq -(\lambda + \mu)^{2/(2-q)} C.$$

We obtain $I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) \rightarrow 0$ as $\lambda, \mu \rightarrow 0$.

Now we show (iv). By (i)-(iii),

$$\begin{aligned} 0 &= \lim_{\lambda,\mu \rightarrow 0} I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) \\ &= \lim_{\lambda,\mu \rightarrow 0} \left(\frac{s}{2N} \|((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+)\|_H^2 \right. \\ &\quad \left. - \left(\frac{1}{q} - \frac{1}{2^*}\right) \int_{\Omega \times \{0\}} (\lambda |(\omega_1)_{\lambda,\mu}^+|^q + \mu |(\omega_2)_{\lambda,\mu}^+|^q) dx \right). \end{aligned} \tag{2.14}$$

Since $I_{\lambda,\mu}$ is coercive and bounded below on $N_{\lambda,\mu}$, $((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+)$ is bounded in H and so that

$$\lim_{\lambda,\mu \rightarrow 0} \int_{\Omega \times \{0\}} (\lambda |(\omega_1)_{\lambda,\mu}^+|^q + \mu |(\omega_2)_{\lambda,\mu}^+|^q) dx = 0. \tag{2.15}$$

Therefore, we obtain the desired result. □

3 Some technical results

In this section, we shall introduce some useful results which are crucial for the proof of Theorem 1.1.

Lemma 3.1 *Let $\{(\omega_{1,n}, \omega_{2,n})\} \subset H$ be a non-negative function sequence with*

$$\int_{\Omega \times \{0\}} |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx = 1 \quad \text{and} \quad \|(\omega_{1,n}, \omega_{2,n})\|_H^2 \rightarrow K_s S_{s,\alpha,\beta}.$$

Then there exists a sequence $\{(y_n, \varepsilon_n)\} \subset \mathbb{R}^N \times \mathbb{R}^+$ such that

$$(W_{1,n}(x), W_{2,n}(x)) := (E_s(\varepsilon_n^{\frac{N-s}{2}} \omega_{1,n}(\varepsilon_n x + y_n, 0)), E_s(\varepsilon_n^{\frac{N-s}{2}} \omega_{2,n}(\varepsilon_n x + y_n, 0)))$$

contains a convergent subsequence denoted again by $\{(W_{1,n}(x), W_{2,n}(x))\}$ such that

$$(W_{1,n}(x), W_{2,n}(x)) \rightarrow (W_1, W_2) \quad \text{in } H.$$

Moreover, we have $\varepsilon_n \rightarrow 0$ and $y_n \rightarrow y \in \overline{\Omega}$ as $n \rightarrow \infty$.

Proof Let $Z_{n,1}(x) = \omega_{1,n}(x, 0)$, $Z_{n,2}(x) = \omega_{2,n}(x, 0)$, we have

$$\int_{\Omega} |Z_{n,1}|^\alpha |Z_{n,2}|^\beta dx = 1 \quad \text{and} \quad \|Z_{n,1}\|_{H_0^s(\Omega)}^2 + \|Z_{n,2}\|_{H_0^s(\Omega)}^2 \rightarrow K_s S_{\alpha,\beta} \quad \text{as } n \rightarrow \infty.$$

By the proof of Lemma 2.2, we know that $\{Z_{n,1}\}$ and $\{Z_{n,2}\}$ are minimizing sequences for the critical Sobolev inequality in the form (2.3). Thus we deduce from [23], Theorem 3 and [23], Theorem 5, that there exist a sequence of points $\{y_n\} \subseteq \mathbb{R}^N$ and a sequence of numbers $\{\varepsilon_n\} \subset (0, \infty)$ such that $\widehat{Z}_{n,1}(x) = \varepsilon_n^{\frac{N-s}{2}} Z_{n,1}(\varepsilon_n x + y_n) \rightarrow \widehat{Z}_1(x)$ and $\widehat{Z}_{n,2}(x) = \varepsilon_n^{\frac{N-s}{2}} Z_{n,2}(\varepsilon_n x + y_n) \rightarrow \widehat{Z}_2(x)$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$. Moreover, we have $\varepsilon_n \rightarrow 0$ and $y_n \rightarrow y \in \overline{\Omega}$ as $n \rightarrow \infty$. Denote $W_{1,n} = E_s(\widehat{Z}_{n,1})$, $W_{2,n} = E_s(\widehat{Z}_{n,2})$ and $W_1 = E_s(\widehat{Z}_1)$, $W_2 = E_s(\widehat{Z}_2)$. Then we obtain the result. \square

Lemma 3.2 *Suppose that X is a Hilbert manifold and $F \in C^1(X, \mathbb{R})$. Assume that, for $c_0 \in \mathbb{R}$ and $K \in \mathbb{N}$:*

- (i) $F(x)$ satisfies the $(PS)_c$ condition for $c \leq c_0$,
- (ii) $\text{cat}(\{x \in X; F(x) \leq c_0\}) \geq K$.

Then $F(x)$ has at least K critical points in $\{x \in X; F(x) \leq c_0\}$.

Proof See [24], Theorem 2.3. \square

Up to translations, we may assume that $0 \in \Omega$. Moreover, in the following, we fix $r > 0$ such that $B_r = \{x \in \mathbb{R}^N; |x| < r\} \subset \Omega$ and the sets

$$\Omega_r^+ := \{x \in \mathbb{R}^N; \text{dist}(x, \Omega) < r\}, \quad \Omega_r^- := \{x \in \Omega; \text{dist}(x, \partial\Omega) > r\}$$

are both homotopically equivalent to Ω .

Noting Remark 2.2, below we can define the continuous map $\Phi : N_{\lambda,\mu}^- \rightarrow \mathbb{R}^N$ by setting

$$\Phi(\omega_1, \omega_2) := \frac{\int_{\Omega \times \{0\}} x |\omega_1|^\alpha |\omega_2|^\beta dx}{\int_{\Omega \times \{0\}} |\omega_1|^\alpha |\omega_2|^\beta dx}.$$

Denote

$$c_{\lambda,\mu} := \alpha_{\lambda,\mu}^+ + \frac{S}{2N} (K_s S_{s,\alpha,\beta})^{N/s}$$

and

$$N_{\lambda,\mu}^-(c_{\lambda,\mu}) := \{(\omega_1, \omega_2) \in N_{\lambda,\mu}^-; I_{\lambda,\mu}(\omega_1, \omega_2) \leq c_{\lambda,\mu}\}.$$

Lemma 3.3 *There exists $\Lambda_* > 0$ such that if $\lambda, \mu \in (0, \Lambda_*)$ and $(\omega_1, \omega_2) \in N_{\lambda,\mu}^-(c_{\lambda,\mu})$,*

$$\Phi(\omega_1, \omega_2) \in \Omega_r^+.$$

Proof By way of contradiction, let $\lambda_n, \mu_n \rightarrow 0$, $\{(\omega_{1,n}, \omega_{2,n})\} \subset N_{\lambda_n, \mu_n}^-(c_{\lambda_n, \mu_n})$ and $\Phi(\omega_{1,n}, \omega_{2,n}) \notin \Omega_r^+$. From (2.7), we see that $\{(\omega_{1,n}, \omega_{2,n})\}$ is bounded and $\int_{\Omega \times \{0\}} (\lambda_n |\omega_{1,n}|^q + \mu_n |\omega_{2,n}|^q) dx \rightarrow 0$. Thus,

$$\begin{aligned} \lim_{n \rightarrow \infty} I_{\lambda_n, \mu_n}(\omega_{1,n}, \omega_{2,n}) &= \lim_{n \rightarrow \infty} \frac{S}{2N} \left\| (\omega_{1,n}, \omega_{2,n}) \right\|_H^2 \\ &= \lim_{n \rightarrow \infty} \frac{S}{2N} \int_{\Omega \times \{0\}} |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx \\ &\leq \frac{S}{2N} (K_s S_{s,\alpha,\beta})^{N/s}. \end{aligned} \tag{3.1}$$

Defining

$$(W_{1,n}, W_{2,n}) = \left(\frac{\omega_{1,n}}{\left(\int_{\Omega \times \{0\}} |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx\right)^{1/(\alpha+\beta)}}, \frac{\omega_{2,n}}{\left(\int_{\Omega \times \{0\}} |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx\right)^{1/(\alpha+\beta)}} \right),$$

we see that $\int_{\Omega \times \{0\}} |W_{1,n}|^\alpha |W_{2,n}|^\beta dx = 1$. By (3.1) and the definition of $S_{s,\alpha,\beta}$, we obtain

$$\lim_{n \rightarrow \infty} \left\| (W_{1,n}, W_{2,n}) \right\|_H^2 = K_s S_{s,\alpha,\beta}.$$

By Lemma 3.1, there is a sequence $\{(y_n, \varepsilon_n)\} \in \mathbb{R}^N \times \mathbb{R}^+$ such that $\varepsilon_n \rightarrow 0$, $y_n \rightarrow y \in \bar{\Omega}$ and $(E_s(\varepsilon_n^{-\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n)), E_s(\varepsilon_n^{-\frac{N-s}{2}} W_{2,n}(\varepsilon_n x + y_n))) \rightarrow (W_1, W_2)$ in H as $n \rightarrow \infty$.

Considering $\varphi \in C_0^\infty(\mathbb{R}^N)$ such that $\varphi(x) = x$ in Ω , we infer

$$\begin{aligned} &\Phi(\omega_{1,n}, \omega_{2,n}) \\ &= \frac{\int_{\Omega \times \{0\}} x |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx}{\int_{\Omega \times \{0\}} |\omega_{1,n}|^\alpha |\omega_{2,n}|^\beta dx} = \frac{\int_{\mathbb{R}^N \times \{0\}} \varphi(x) |W_{1,n}|^\alpha |W_{2,n}|^\beta dx}{\int_{\mathbb{R}^N \times \{0\}} |W_{1,n}|^\alpha |W_{2,n}|^\beta dx} \\ &= \frac{\int_{\mathbb{R}^N \times \{0\}} \varphi(\varepsilon_n x + y_n) |E_s(\varepsilon_n^{-\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n))|^\alpha |E_s(\varepsilon_n^{-\frac{N-s}{2}} W_{2,n}(\varepsilon_n x + y_n))|^\beta dx}{\int_{\mathbb{R}^N \times \{0\}} |E_s(\varepsilon_n^{-\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n))|^\alpha |E_s(\varepsilon_n^{-\frac{N-s}{2}} W_{2,n}(\varepsilon_n x + y_n))|^\beta dx}. \end{aligned}$$

Moreover, by the Lebesgue theorem, we have

$$\frac{\int_{\mathbb{R}^N \times \{0\}} \varphi(\varepsilon_n x + y_n) |E_s(\varepsilon_n^{\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n))|^\alpha |E_s(\varepsilon_n^{\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n))|^\beta dx}{\int_{\mathbb{R}^N \times \{0\}} |E_s(\varepsilon_n^{\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n))|^\alpha |E_s(\varepsilon_n^{\frac{N-s}{2}} W_{1,n}(\varepsilon_n x + y_n))|^\beta dx} \rightarrow y \in \overline{\Omega},$$

as $n \rightarrow \infty$, so that $\lim_{n \rightarrow \infty} \Phi(\omega_{1,n}, \omega_{2,n}) = y \in \overline{\Omega}$, in contradiction with $\Phi(\omega_{1,n}, \omega_{2,n}) \notin \Omega_r^+$. \square

Next, we will use $\omega_\varepsilon = E_s(u_\varepsilon)$, the family of minimizers to the inequality (2.4), where u_ε is given in (2.5). Let $\eta \in C^\infty(C_\Omega)$, $0 \leq \eta(x, y) \leq 1$ and for small fixed ρ ,

$$\eta(x, y) = \begin{cases} 1, & (x, y) \in B_\rho^+ := \{(x_1, x_2, \dots, x_N, y); \sqrt{x_1^2 + x_2^2 + \dots + x_N^2 + y^2} < \frac{\rho}{2}, y > 0\}, \\ 0, & (x, y) \notin B_\rho^+ := \{(x_1, x_2, \dots, x_N, y); \sqrt{x_1^2 + x_2^2 + \dots + x_N^2 + y^2} < \rho, y > 0\}. \end{cases}$$

We take $\rho < \frac{r}{4}$ small enough such that

$$\overline{B_\rho^+(x - z, y)} \subset \overline{C_\Omega}$$

for all $z \in \Omega_r^-$, where

$$\overline{B_\rho^+(x - z, y)} := \{(x_1, x_2, \dots, x_N, y); \sqrt{(x_1 - z_1)^2 + (x_2 - z_2)^2 + \dots + (x_N - z_N)^2 + y^2} \leq \rho, y \geq 0\}.$$

Assume

$$v_{\varepsilon, z} = \eta(x - z, y) \omega_\varepsilon(x - z, y) = \eta(x - z, y) E_s(u_\varepsilon(x - z)), \quad z \in \Omega_r^-,$$

where u_ε is defined in (2.5). We obtain from [11]

$$\|v_{\varepsilon, z}\|_{H_{0,L}^s(C_\Omega)}^2 = K_s \int_{\mathbb{R}_+^{N+1}} y^{1-s} |\nabla \omega_\varepsilon|^2 dx dy + O(\varepsilon^{N-s}), \tag{3.2}$$

$$\int_{\Omega \times \{0\}} |v_{\varepsilon, z}|^{2_s^*} dx = \int_{\mathbb{R}^N \times \{0\}} |\omega_\varepsilon|^{2_s^*} dx + O(\varepsilon^N) = \int_{\mathbb{R}^N} \left(\frac{\varepsilon}{\varepsilon^2 + |x|^2} \right)^N dx + O(\varepsilon^N). \tag{3.3}$$

Then we have the following.

Lemma 3.4 *There exist $\varepsilon_0, \sigma(\varepsilon) > 0$ such that, for $\varepsilon \in (0, \varepsilon_0)$ and $\sigma \in (0, \sigma(\varepsilon))$, we have*

$$\sup_{t \geq 0} I_{\lambda, \mu}((\omega_1)_{\lambda, \mu}^+ + t\sqrt{\alpha}v_{\varepsilon, z}, (\omega_2)_{\lambda, \mu}^+ + t\sqrt{\beta}v_{\varepsilon, z}) < c_{\lambda, \mu} - \sigma \quad \text{uniformly in } z \in \Omega_r^-,$$

where $((\omega_1)_{\lambda, \mu}^+, (\omega_2)_{\lambda, \mu}^+)$ is a local minimum in Lemma 2.8. Furthermore, there exists $t_{(\lambda, \mu, \varepsilon, z)}^- > 0$ such that

$$((\omega_1)_{\lambda, \mu}^+ + t_{(\lambda, \mu, \varepsilon, z)}^- \sqrt{\alpha}v_{\varepsilon, z}, (\omega_2)_{\lambda, \mu}^+ + t_{(\lambda, \mu, \varepsilon, z)}^- \sqrt{\beta}v_{\varepsilon, z}) \in N_{\lambda, \mu}^-(c_{\lambda, \mu} - \sigma)$$

and

$$\Phi((\omega_1)_{\lambda,\mu}^+ + t^-(\omega_1)_{\lambda,\mu,\varepsilon,z}^-, (\omega_2)_{\lambda,\mu}^+ + t^-(\omega_2)_{\lambda,\mu,\varepsilon,z}^-) \in \Omega_r^+.$$

Proof Since

$$\begin{aligned} & I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}v_{\varepsilon,z}, (\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}v_{\varepsilon,z}) \\ &= \frac{K_s}{2} \int_{C_\Omega} y^{1-s} (|\nabla((\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}v_{\varepsilon,z})|^2 + |\nabla((\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}v_{\varepsilon,z})|^2) dx dy \\ &\quad - \frac{1}{q} \int_{\Omega \times \{0\}} (\lambda |(\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)|^q \\ &\quad + \mu |(\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}\eta(x-z, 0)u_\varepsilon(x-z)|^q) dx \\ &\quad - \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |(\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)|^\alpha \\ &\quad \times |(\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}\eta(x-z, 0)u_\varepsilon(x-z)|^\beta dx \\ &= \frac{1}{2} \|((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+)\|_H^2 + t^2 \frac{\alpha + \beta}{2} \|v_{\varepsilon,z}\|_{H_{0,L}^s(C_\Omega)}^2 \\ &\quad + tK_s \left(\int_{C_\Omega} \nabla(\omega_1)_{\lambda,\mu}^+ \nabla(\sqrt{\alpha}v_{\varepsilon,z}) dx dy + \int_{C_\Omega} \nabla(\omega_2)_{\lambda,\mu}^+ \nabla(\sqrt{\beta}v_{\varepsilon,z}) dx dy \right) \\ &\quad - \frac{1}{q} \int_{\Omega \times \{0\}} (\lambda |(\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)|^q \\ &\quad + \mu |(\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}\eta(x-z, 0)u_\varepsilon(x-z)|^q) dx \\ &\quad - \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |(\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)|^\alpha \\ &\quad \times |(\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}\eta(x-z, 0)u_\varepsilon(x-z)|^\beta dx \\ &\leq I_{\lambda,\mu}((\omega_1)_{\lambda,\mu}^+, (\omega_2)_{\lambda,\mu}^+) + t^2 \frac{\alpha + \beta}{2} \|v_{\varepsilon,z}\|_{H_{0,L}^s(C_\Omega)}^2 \\ &\quad - \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |(\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)|^\alpha \\ &\quad \times |(\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}\eta(x-z, 0)u_\varepsilon(x-z)|^\beta dx + \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |(\omega_1)_{\lambda,\mu}^+|^\alpha |(\omega_2)_{\lambda,\mu}^+|^\beta dx \\ &\quad + \frac{1}{2_s^*} \int_{\Omega \times \{0\}} \alpha |(\omega_1)_{\lambda,\mu}^+|^{\alpha-1} |(\omega_2)_{\lambda,\mu}^+|^\beta (t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)) dx \\ &\quad + \frac{1}{2_s^*} \int_{\Omega \times \{0\}} \beta |(\omega_1)_{\lambda,\mu}^+|^\alpha |(\omega_2)_{\lambda,\mu}^+|^{\beta-1} (t\sqrt{\beta}\eta(x-z, 0)u_\varepsilon(x-z)) dx \\ &= \alpha_{\lambda,\mu}^+ + K(t), \end{aligned} \tag{3.4}$$

where

$$\begin{aligned} K(t) &= t^2 \frac{\alpha + \beta}{2} \|v_{\varepsilon,z}\|_{H_{0,L}^s(C_\Omega)}^2 \\ &\quad - \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |(\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}\eta(x-z, 0)u_\varepsilon(x-z)|^\alpha \end{aligned}$$

$$\begin{aligned} & \times |(\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}\eta(x-z,0)u_\varepsilon(x-z)|^\beta dx \\ & + \frac{1}{2_s^*} \int_{\Omega \times \{0\}} |(\omega_1)_{\lambda,\mu}^+|^\alpha |(\omega_2)_{\lambda,\mu}^+|^\beta dx \\ & + \frac{1}{2_s^*} \int_{\Omega \times \{0\}} \alpha |(\omega_1)_{\lambda,\mu}^+|^{\alpha-1} |(\omega_2)_{\lambda,\mu}^+|^\beta (t\sqrt{\alpha}\eta(x-z,0)u_\varepsilon(x-z)) dx \\ & + \frac{1}{2_s^*} \int_{\Omega \times \{0\}} \beta |(\omega_1)_{\lambda,\mu}^+|^\alpha |(\omega_2)_{\lambda,\mu}^+|^{\beta-1} (t\sqrt{\beta}\eta(x-z,0)u_\varepsilon(x-z)) dx. \end{aligned}$$

In the following we shall show that

$$\sup_{t \geq 0} K(t) < \frac{S}{2N} (K_s S_{s,\alpha,\beta})^{N/s} \quad \text{for } \varepsilon > 0 \text{ small enough.}$$

It is easy to see that

$$\lim_{t \rightarrow 0} K(t) = 0.$$

Thus, for all ε sufficiently small, there exists $t_0 > 0$ such that

$$K(t) < \frac{S}{2N} (K_s S_{s,\alpha,\beta})^{N/s} \quad \text{for all } t \in (0, t_0]. \tag{3.5}$$

From [25], Lemma 4.1, we see that there exist $C_1(\alpha), C_2(\beta) > 0$ such that

$$\begin{aligned} (a+b)^\alpha (c+d)^\beta & \geq a^\alpha c^\beta + a^\alpha d^\beta + b^\alpha c^\beta + b^\alpha d^\beta + C_1(\alpha) a^{\alpha-1} b c^\beta + C_1(\alpha) a^{\alpha-1} b d^\beta \\ & \quad + C_2(\beta) b^\alpha c^{\beta-1} d + C_2(\beta) a^\alpha c^{\beta-1} d + C_1(\alpha) C_2(\beta) a^{\alpha-1} b c^{\beta-1} d \end{aligned}$$

for any $a, b, c, d > 0$. Consequently,

$$K(t) \leq t^2 \frac{2_s^*}{2} \|v_{\varepsilon,z}\|_{H_{0,L}^s(C_\Omega)}^2 - \frac{\alpha^{\frac{\alpha}{2}} \beta^{\frac{\beta}{2}}}{2_s^*} t^{2_s^*} \int_{\Omega \times \{0\}} |v_{\varepsilon,z}|^{2_s^*} dx - C_1 \int_{\Omega \times \{0\}} |v_{\varepsilon,z}|^{2_s^*-1} dx \tag{3.6}$$

with some constant $C_1 > 0$. Note that

$$\begin{aligned} & \int_{\Omega \times \{0\}} |v_{\varepsilon,z}|^{2_s^*-1} dx \\ & = \int_{\Omega \times \{0\}} |\eta(x-z,0)u_\varepsilon(x-z)|^{2_s^*-1} dx \\ & = \int_{B_{2\rho}} \left[\frac{\eta(x,0)\varepsilon^{\frac{N-s}{2}}}{(\varepsilon^2 + |x|^2)^{\frac{N-s}{2}}} \right]^{\frac{N+s}{N-s}} dx \\ & \geq \int_{B_\rho} \frac{\varepsilon^{\frac{N+s}{2}}}{\varepsilon^{N+s}(\varepsilon^2 + |x|^2)^{\frac{N+s}{2}}} \varepsilon^N dx \\ & = C_2 \varepsilon^{\frac{N-s}{2}} \int_0^\rho \frac{r^{N-1}}{(1+r^2)^{\frac{N+s}{2}}} dr \\ & = C_3 \varepsilon^{\frac{N-s}{2}} \end{aligned} \tag{3.7}$$

for some $C_2, C_3 > 0$. It follows from (3.2)-(3.7) that

$$\begin{aligned}
 K(t) &\leq t^2 \frac{2^*}{2} \|v_{\varepsilon,z}\|_{H_{0,L}^s(C\Omega)}^2 - \frac{\alpha^{\frac{\alpha}{2}} \beta^{\frac{\beta}{2}}}{2_s^*} t^{2_s^*} \int_{\Omega \times \{0\}} |v_{\varepsilon,z}|^{2_s^*} dx - C_4 \varepsilon^{\frac{N-s}{2}} \\
 &\leq \frac{s}{2N} \left(\frac{(\alpha + \beta) \|v_{\varepsilon,z}\|_{H_{0,L}^s(C\Omega)}^2}{\left(\int_{\Omega \times \{0\}} \alpha^{\frac{\alpha}{2}} \beta^{\frac{\beta}{2}} |v_{\varepsilon,z}|^{2_s^*} dx \right)^{\frac{2}{2_s^*}}} \right)^{\frac{N}{s}} - C_4 \varepsilon^{\frac{N-s}{2}} \\
 &= \frac{s}{2N} \left(\left(\left(\frac{\alpha}{\beta} \right)^{\frac{\beta}{\alpha+\beta}} + \left(\frac{\beta}{\alpha} \right)^{\frac{\alpha}{\alpha+\beta}} \right) \frac{K_s \int_{\mathbb{R}_+^{N+1}} y^{1-s} |\nabla \omega_\varepsilon|^2 dx dy + O(\varepsilon^{N-s})}{\left(\int_{\mathbb{R}^N} \left(\frac{\varepsilon}{\varepsilon^2 + |x|^2} \right)^N dx + O(\varepsilon^N) \right)^{\frac{2}{2_s^*}}} \right)^{\frac{N}{s}} - C_4 \varepsilon^{\frac{N-s}{2}} \\
 &= \frac{s}{2N} (K_s S_{s,\alpha,\beta})^{\frac{N}{s}} + O(\varepsilon^{N-s}) - C_4 \varepsilon^{\frac{N-s}{2}} \\
 &< \frac{s}{2N} (K_s S_{s,\alpha,\beta})^{\frac{N}{s}} \tag{3.8}
 \end{aligned}$$

for ε sufficiently small and $t \in [t_0, +\infty)$. Noting the compactness of $\overline{\Omega_r^-}$, it follows from (3.4)-(3.5) and (3.8) that there exist $\varepsilon_0, \sigma(\varepsilon) > 0$ such that, for $\varepsilon \in (0, \varepsilon_0)$ and $\sigma \in (0, \sigma(\varepsilon))$, we have

$$\sup_{t \geq 0} I_{\lambda,\mu} \left((\omega_1)_{\lambda,\mu}^+ + t\sqrt{\alpha}v_{\varepsilon,z}, (\omega_2)_{\lambda,\mu}^+ + t\sqrt{\beta}v_{\varepsilon,z} \right) < c_{\lambda,\mu} - \sigma \quad \text{uniformly in } z \in \Omega_r^-.$$

Arguing as the proof of [11], Lemma 4.4, we conclude that there exists $t_{(\lambda,\mu,\varepsilon,z)}^- > 0$ such that

$$\left((\omega_1)_{\lambda,\mu}^+ + t_{(\lambda,\mu,\varepsilon,z)}^- \sqrt{\alpha}v_{\varepsilon,z}, (\omega_2)_{\lambda,\mu}^+ + t_{(\lambda,\mu,\varepsilon,z)}^- \sqrt{\beta}v_{\varepsilon,z} \right) \in N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma).$$

Moreover, we obtain from Lemma 3.3

$$\Phi \left((\omega_1)_{\lambda,\mu}^+ + t_{(\lambda,\mu,\varepsilon,z)}^- \sqrt{\alpha}v_{\varepsilon,z}, (\omega_2)_{\lambda,\mu}^+ + t_{(\lambda,\mu,\varepsilon,z)}^- \sqrt{\beta}v_{\varepsilon,z} \right) \in \Omega_r^+$$

for $\lambda, \mu \in (0, \Lambda_*)$. □

From Lemma 3.4, we can define the map $\gamma : \Omega_r^- \rightarrow N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma)$ defined by

$$\gamma(z) := \left((\omega_1)_{\lambda,\mu}^+ + t_{(\lambda,\mu,\varepsilon,z)}^- \sqrt{\alpha}v_{\varepsilon,z}, (\omega_2)_{\lambda,\mu}^+ + t_{(\lambda,\mu,\varepsilon,z)}^- \sqrt{\beta}v_{\varepsilon,z} \right).$$

Furthermore, by Lemma 2.6 and Lemma 2.8(iv), we can define the map $\Phi_{\lambda,\mu} : N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma) \rightarrow \mathbb{R}^N$ by setting

$$\Phi_{\lambda,\mu}(\omega_1, \omega_2) := \frac{\int_{\Omega \times \{0\}} x |\omega_1 - (\omega_1)_{\lambda,\mu}^+|^\alpha |\omega_2 - (\omega_2)_{\lambda,\mu}^+|^\beta dx}{\int_{\Omega \times \{0\}} |\omega_1 - (\omega_1)_{\lambda,\mu}^+|^\alpha |\omega_2 - (\omega_2)_{\lambda,\mu}^+|^\beta dx}.$$

Then, for each $z \in \Omega_r^-$, note that $u_\varepsilon(x)$ is radial, we have

$$(\Phi_{\lambda,\mu} \circ \gamma)(z) = z.$$

Next we define the map $H_{\lambda,\mu} : [0, 1] \times N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma) \rightarrow \mathbb{R}^N$ by

$$H_{\lambda,\mu}(t, (\omega_1, \omega_2)) = t\Phi_{\lambda,\mu}(\omega_1, \omega_2) + (1 - t)\Phi_{\lambda,\mu}(\omega_1, \omega_2).$$

Lemma 3.5 *For each $\varepsilon \in (0, \varepsilon_0)$, there exists $\Lambda_* > 0$ such that if $\lambda, \mu, \sigma \in (0, \Lambda_*)$,*

$$H_{\lambda,\mu}([0, 1] \times N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma)) \subset \Omega_r^+.$$

Proof Suppose by contradiction that there exist $t_n \in [0, 1]$, $\lambda_n, \mu_n, \sigma_n \rightarrow 0$, and $(\omega_{1,n}, \omega_{2,n}) \in N_{\lambda_n, \mu_n}^-(c_{\lambda_n, \mu_n} - \sigma_n)$ such that

$$H_{\lambda_n, \mu_n}(t_n, (\omega_{1,n}, \omega_{2,n})) \notin \Omega_r^+ \quad \text{for all } n.$$

Furthermore, we can assume that $t_n \rightarrow t_0 \in [0, 1]$. Then by Lemma 2.8(iv) and argue as in the proof of Lemma 3.3, we have

$$H_{\lambda_n, \mu_n}(t_n, (\omega_{1,n}, \omega_{2,n})) \rightarrow z \in \overline{\Omega} \quad \text{as } n \rightarrow \infty,$$

which is a contradiction. □

4 Proof of Theorem 1.1

We begin with the following lemma.

Lemma 4.1 *If (ω_1, ω_2) is a critical point of $I_{\lambda,\mu}$ on $N_{\lambda,\mu}^-$, then it is a critical point of $I_{\lambda,\mu}$ in H .*

Proof Assume $(\omega_1, \omega_2) \in N_{\lambda,\mu}^-$, then $I'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) = 0$. On the other hand,

$$I'_{\lambda,\mu}(\omega_1, \omega_2) = \theta \psi'_{\lambda,\mu}(\omega_1, \omega_2) \tag{4.1}$$

for some $\theta \in \mathbb{R}$, where $\psi_{\lambda,\mu}$ is defined in (2.9).

Remark that $(\omega_1, \omega_2) \in N_{\lambda,\mu}^-$, and so $\psi'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2) < 0$. Thus by (4.1),

$$0 = \theta \psi'_{\lambda,\mu}(\omega_1, \omega_2)(\omega_1, \omega_2),$$

which implies that $\theta = 0$, consequently $I'_{\lambda,\mu}(\omega_1, \omega_2) = 0$. □

Below we denote by $I_{N_{\lambda,\mu}^-}$ the restriction of $I_{\lambda,\mu}$ on $N_{\lambda,\mu}^-$.

Lemma 4.2 *Any sequence $\{(\omega_{1,n}, \omega_{2,n})\} \subset N_{\lambda,\mu}^-$ such that $I_{N_{\lambda,\mu}^-}(\omega_{1,n}, \omega_{2,n}) \rightarrow c \in (-\infty, c_{\lambda,\mu})$ and $I'_{N_{\lambda,\mu}^-}(\omega_{1,n}, \omega_{2,n}) \rightarrow 0$ contains a convergent subsequence for all $\lambda, \mu \in (0, \Lambda_*)$.*

Proof By hypothesis there exists a sequence $\{\theta_n\} \subset \mathbb{R}$ such that

$$I'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n}) = \theta_n \psi'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n}) + o(1).$$

Recall that $(\omega_{1,n}, \omega_{2,n}) \in N_{\lambda,\mu}^-$ and so

$$\psi'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n})(\omega_{1,n}, \omega_{2,n}) < 0.$$

If $\psi'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n})(\omega_{1,n}, \omega_{2,n}) \rightarrow 0$, we see from (2.10) and (2.11) that there are two positive numbers C_1, C_2 independent of $(\omega_{1,n}, \omega_{2,n})$ and λ, μ such that

$$\begin{aligned} \|\omega_{1,n}, \omega_{2,n}\|_H^2 &\leq C_1 \|\omega_{1,n}, \omega_{2,n}\|_H^{2_s^*} + o(1) \quad \text{and} \\ \|\omega_{1,n}, \omega_{2,n}\|_H^2 &\leq (\lambda + \mu) C_2 \|\omega_{1,n}, \omega_{2,n}\|_H^q + o(1) \end{aligned}$$

or

$$\begin{aligned} \|\omega_{1,n}, \omega_{2,n}\|_H^2 &\geq C_1^{-\frac{2}{2_s^*-2}} + o(1) \quad \text{and} \\ \|\omega_{1,n}, \omega_{2,n}\|_H^2 &\leq (\lambda + \mu)^{\frac{2}{2-q}} c_2^{\frac{2}{2-q}} + o(1). \end{aligned}$$

If $\lambda, \mu > 0$ is sufficiently small, this is impossible. Thus we may assume that $\psi'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n})(\omega_{1,n}, \omega_{2,n}) \rightarrow l < 0$ as $n \rightarrow \infty$. Since $I'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n})(\omega_{1,n}, \omega_{2,n}) = 0$, we conclude that $\theta_n \rightarrow 0$ and, consequently, $I'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n}) \rightarrow 0$. Using this information we have

$$I_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n}) \rightarrow c \in (-\infty, c_{\lambda,\mu}) \quad \text{and} \quad I'_{\lambda,\mu}(\omega_{1,n}, \omega_{2,n}) \rightarrow 0,$$

so by Lemma 2.7 the proof is complete. □

Lemma 4.3 *If $\lambda, \mu, \sigma \in (0, \Lambda_*)$, then*

$$\text{cat}(N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma)) \geq \text{cat}(\Omega).$$

Proof Suppose that

$$N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma) = A_1 \cup \dots \cup A_n,$$

where $A_j, j = 1, \dots, n$, is closed and contractible in $N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma)$, i.e., there exists $h_j \in C([0, 1] \times A_j, N_{\lambda,\mu}^-(c_{\lambda,\mu} - \sigma))$ such that

$$h_j(0, z) = z \quad \text{and} \quad h_j(1, z) = \omega \quad \text{for all } z \in A_j,$$

where $\omega \in A_j$ is fixed. Consider $B_j := \gamma^{-1}(A_j), 1 \leq j \leq n$. The sets B_j are closed and

$$\Omega_r^- = B_1 \cup \dots \cup B_n.$$

Noting Lemma 3.5, we define the deformation $g_j : [0, 1] \times B_j \rightarrow \Omega_r^+$ by setting

$$g_j(t, y) := H_{\lambda,\mu}(t, h_j(t, \gamma(y)))$$

for $\lambda, \mu, \sigma \in (0, \Lambda_*)$. Note that

$$g_j(0, y) := H_{\lambda,\mu}(0, h_j(0, \gamma(y))) = y \quad \text{for all } y \in B_j$$

and

$$g_j(1, y) := H_{\lambda, \mu}(1, h_j(1, \gamma(y))) = \Phi_{\lambda, \mu}(\omega) \in \Omega_r^+.$$

Thus the sets B_j are contractible in Ω_r^+ . It follows that

$$\text{cat}(\Omega) = \text{cat}_{\Omega_r^+}(\Omega_r^-) \leq n. \quad \square$$

Now, we can give the proof of Theorem 1.1.

Proof of Theorem 1.1 Applying Lemmas 2.7 and 4.2, $I_{N_{\lambda, \mu}^-}$ satisfies $(PS)_c$ condition for all $c \in (-\infty, c_{\lambda, \mu})$. Then, by Lemmas 3.2 and 4.3, $I_{N_{\lambda, \mu}^-}$ contains at least $\text{cat}(\Omega)$ critical points in $N_{\lambda, \mu}^-(c_{\lambda, \mu} - \sigma)$. Hence, we deduce from Lemma 4.1 that $I_{\lambda, \mu}$ has at least $\text{cat}(\Omega)$ critical points in $N_{\lambda, \mu}^-$. Moreover, by Lemma 2.6 and $N_{\lambda, \mu}^+ \cap N_{\lambda, \mu}^- = \emptyset$, $I_{\lambda, \mu}$ has at least $\text{cat}(\Omega) + 1$ critical points in H . If we change the definition of $J_{\lambda, \mu}$ as follows:

$$J_{\lambda, \mu}(u, v) := \frac{1}{2} \int_{\Omega} (|(-\Delta)^{\frac{s}{4}} u|^2 + |(-\Delta)^{\frac{s}{4}} v|^2) dx - \frac{1}{q} \int_{\Omega} (\lambda u_+^q + \mu v_+^q) dx - \frac{1}{2_s^*} \int_{\Omega} u_+^\alpha v_+^\beta dx, \quad (4.2)$$

where $u_+ = \max\{u, 0\}$ and $v_+ = \max\{v, 0\}$. Then all the steps of our paper for (4.2). Thus we see that $J_{\lambda, \mu}$ has at least $\text{cat}(\Omega) + 1$ non-negative critical points. By the maximum principle [26], we complete the proof. \square

Competing interests

The authors declare that they have no competing interests.

Author's contributions

HF carried out the proofs of the theorems and the check of the manuscript. The author read and approved the final manuscript.

Acknowledgements

The authors are very grateful to the referees for their careful reading, comments, and suggestions, which improved the presentation of this paper. This work is supported by the Natural Science Foundation of Jiangsu Province (Grant No. BK20150168).

Received: 13 July 2016 Accepted: 26 October 2016 Published online: 08 November 2016

References

1. Applebaum, D: Lévy process-from probability to finance and quantum groups. *Not. Am. Math. Soc.* **51**, 1336-1347 (2004)
2. Vlahos, L, Isliker, H, Komninos, Y, Hizonidis, K: Normal and anomalous diffusion: a tutorial. In: Bountis, T (ed.) *Order and Chaos*, vol. 10. Patras University Press, Patras (2008)
3. Servadei, R, Valdinoci, E: Mountain pass solutions for nonlinear elliptic operators. *J. Math. Anal. Appl.* **389**, 887-898 (2012)
4. Servadei, R, Valdinoci, E: Lewy-Stampacchia type estimates for variational inequalities driven by nonlocal operators. *Rev. Mat. Iberoam.* **29**, 1091-1126 (2013)
5. Serra, J, Ros-Oton, X: The Dirichlet problem for the fractional Laplacian: regularity up to boundary. *J. Math. Pures Appl.* **101**, 275-302 (2014)
6. Yu, X: The Nehari manifold for elliptic equation involving the square root of the Laplacian. *J. Differ. Equ.* **252**, 1283-1308 (2012)
7. Wei, Y, Su, X: Multiplicity of solutions for non-local elliptic equations driven by the fractional Laplacian. *Calc. Var. Partial Differ. Equ.* **52**, 95-124 (2015)
8. Barrios, B, Colorado, E, Servadei, R, Soria, F: A critical fractional equation with concave-convex power nonlinearities. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **32**, 875-900 (2015)
9. Chen, W, Deng, S: The Nehari manifold for a nonlinear elliptic operators involving concave-convex nonlinearities. *Z. Angew. Math. Phys.* **66**, 1387-1400 (2015)

10. Colorado, E, de Pablo, A, Sánchez, U: Perturbation of a critical fractional equations. *Pac. J. Math.* **271**, 65-85 (2014)
11. Shang, X, Zhang, J, Yang, Y: Positive solutions of nonhomogeneous fractional Laplacian problem with critical exponent. *Commun. Pure Appl. Anal.* **13**, 567-584 (2014)
12. Servadei, R, Valdinoci, E: The Brezis-Nirenberg result for the fractional Laplacian. *Trans. Am. Math. Soc.* **367**, 67-102 (2015)
13. Goyal, S, Sreenadh, K: A Nehari manifold for non-local elliptic operator with concave-convex non-linearities and sign-changing weight function (2016). arXiv:1307.5149
14. He, X, Squassina, M, Zou, W: The Nehari manifold for fractional systems involving critical nonlinearities (2016). arXiv:1509.02713
15. Chen, W, Deng, S: The Nehari manifold for a fractional p -Laplacian system involving concave-convex nonlinearities. *Nonlinear Anal., Real World Appl.* **27**, 80-92 (2016)
16. Benci, V, Cerami, G: The effect of the domain topology on the number of positive solutions of nonlinear elliptic problems. *Arch. Ration. Mech. Anal.* **114**, 79-93 (1991)
17. Han, P: The effect of the domain topology on the number of positive solutions of elliptic systems involving critical Sobolev exponents. *Houst. J. Math.* **32**, 1241-1257 (2006)
18. Benmouloud, S, Echarchaoui, R, Sbaj, SM: Multiplicity of positive solutions for a critical quasilinear elliptic system with concave and convex nonlinearities. *J. Math. Anal. Appl.* **396**, 375-385 (2012)
19. Alves, CO, Ding, YH: Multiplicity of positive solutions for the p -Laplacian equation involving critical nonlinearity. *J. Math. Anal. Appl.* **279**, 508-521 (2003)
20. Ding, L, Xiao, SW: Multiple positive solutions for a critical quasilinear elliptic system. *Nonlinear Anal.* **72**, 2592-2607 (2010)
21. Fan, H: Multiple positive solutions for a critical elliptic system with concave and convex nonlinearities. *Nonlinear Anal., Real World Appl.* **18**, 14-22 (2014)
22. Barrios, B, Colorado, E, de Pablo, A, Sánchez, U: On some critical problems for the fractional Laplacian. *J. Differ. Equ.* **252**, 6133-6162 (2012)
23. Palatucci, G, Pisante, A: Improved Sobolev embeddings, profile decomposition, and concentration-compactness for fractional Sobolev spaces. *Calc. Var. Partial Differ. Equ.* (2014). doi:10.1007/s00526-013-0656-y
24. Alves, CO, de Morais Filho, DC, Souto, MAS: On systems of elliptic equations involving subcritical or critical Sobolev exponents. *Nonlinear Anal.* **42**, 771-787 (2000)
25. Chen, CY, Wu, TF: The Nehari manifold for indefinite semilinear elliptic systems involving critical exponent. *Appl. Math. Comput.* **218**, 10817-10828 (2012)
26. Silvestre, L: Regularity of the obstacle problem for a fractional power of the Laplace operator. *Commun. Pure Appl. Math.* **60**, 67-112 (2007)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
