

RESEARCH

Open Access



Targeted single-cell genomics reveals novel host adaptation strategies of the symbiotic bacteria *Endozoicomonas* in *Acropora tenuis* coral

Keigo Ide^{1,2,3†}, Yohei Nishikawa^{2,3†}, Toru Maruyama^{1†}, Yuko Tsukada^{1,2}, Masato Kogawa^{1,2,3}, Hiroki Takeda¹, Haruka Ito¹, Ryota Wagatsuma^{1,2}, Rimi Miyaoka¹, Yoshikatsu Nakano^{4,5}, Koji Kinjo⁶, Michihiro Ito⁴, Masahito Hosokawa^{1,2,3,7}, Kei Yura^{1,2,3,8}, Shoichiro Suda⁹ and Haruko Takeyama^{1,2,3,7*}

Abstract

Background: *Endozoicomonas* bacteria symbiosis with various marine organisms is hypothesized as a potential indicator of health in corals. Although many amplicon analyses using 16S rRNA gene have suggested the diversity of *Endozoicomonas* species, genome analysis has been limited due to contamination of host-derived sequences and difficulties in culture and metagenomic analysis. Therefore, the evolutionary and functional potential of individual *Endozoicomonas* species symbiotic with the same coral species remains unresolved.

Results: In this study, we applied a novel single-cell genomics technique using droplet microfluidics to obtain single-cell amplified genomes (SAGs) for uncultured coral-associated *Endozoicomonas* spp. We obtained seven novel *Endozoicomonas* genomes and quantitative bacterial composition from *Acropora tenuis* corals at four sites in Japan. Our quantitative 16S rRNA gene and comparative genomic analysis revealed that these *Endozoicomonas* spp. belong to different lineages (Clade A and Clade B), with widely varying abundance among individual corals. Furthermore, each *Endozoicomonas* species possessed various eukaryotic-like genes in clade-specific genes. It was suggested that these eukaryotic-like genes might have a potential ability of different functions in each clade, such as infection of the host coral or suppression of host immune pathways. These *Endozoicomonas* species may have adopted different host adaptation strategies despite living symbiotically on the same coral.

Conclusions: This study suggests that coral-associated *Endozoicomonas* spp. on the same species of coral have different evolutionary strategies and functional potentials in each species and emphasizes the need to analyze the genome of each uncultured strain in future coral-*Endozoicomonas* relationships studies.

Background

In recent years, coral bleaching has been known to have several impacts on marine ecosystems. Corals have symbionts, including intracellular symbiotic dinoflagellates, coral-associated bacteria, and viruses, which have been reported to play an important role in coral health and survival [1, 2]. The importance of consideration for the function of the coral itself as well as the interactions among the holobionts in the coral ecology has been

[†]Keigo Ide, Yohei Nishikawa and Toru Maruyama contributed equally to this work.

*Correspondence: haruko-takeyama@waseda.jp

¹ Department of Life Science and Medical Bioscience, Waseda University, Tokyo, Japan

Full list of author information is available at the end of the article



globally recognized. In the functional analysis of coral symbionts, symbiotic dinoflagellates have been the main target, but recently, the functions of coral-associated bacteria have also been gradually clarified, including the degradation of dead symbiotic dinoflagellates [3], provision of nitrogen sources [4], and protection from pathogenic bacteria via antibiotic activities [5].

The genus *Endozoicomonas* has been found to be abundant in corals worldwide and is associated with an extensive range of marine organisms, including shellfish [6, 7], sea slugs [8], sponges [9, 10], and sea anemones [11, 12]. *Endozoicomonas* has been the most widely studied coral-associated bacterium, and it has been shown that phylogenetically different *Endozoicomonas* species colocalize within a single individual [3, 13]. In addition, *Endozoicomonas* has been reported to play several functional roles in coral survival, such as suppressing mitochondrial degradation, supplying sugars, and contributing to the coral sulfur cycle through dimethylsulphoniopropionate (DMSP) [14, 15]. However, little is known about the mechanisms by which *Endozoicomonas* establishes symbiotic relationships with its host corals. Recently, Ding et al. investigated the host cell entry mechanism of *Endozoicomonas montiporae* in *Montipora* coral. They hypothesized that this bacterium used eukaryotic-like genes, such as ephrin ligands, to enter the host cell [14]. This adaptation strategy is consistent with the fact that *Endozoicomonas* is not an obligate symbiont [14, 16]. *Endozoicomonas* diversity has been studied at the 16S rRNA gene level, but not yet at the genome level. Therefore, there are method-dependent gaps in our understanding of *Endozoicomonas* diversity. Furthermore, comparative genome analysis of *Endozoicomonas* spp. from the same host coral species can reveal the evolutionary and adaptive strategies of each *Endozoicomonas*. However, these understandings require genomic information on many *Endozoicomonas* from the same host coral species.

The lack of genomic information is due to the difficulty in culturing host-associated bacteria [16]. Thus, culture-independent approaches are needed for studying genome data to understand the relationship between *Endozoicomonas* and its host. Currently, shotgun metagenomics with binning is the mainstream method for microbial genome sequence analysis, but this method is greatly influenced by microbiome composition and diversity [17]. In most cases, especially in genome analysis of closely related species with highly similar 16S rRNA gene sequences, it is impossible to identify individual strain-level draft genome sequences by metagenomic binning [18]. Furthermore, detailed genome sequence analysis of host-associated bacteria is complicated because contamination by host also significantly affects binning efficiency [16, 19–21]. As

an alternative to metagenomics, single-cell genomics is very useful, as it can reveal the genome diversity of individual cells [22–24] from a mixed cell population. On the other hand, in the current single-cell genomics, single-cells are randomly sorted for subsequent analysis. Therefore, when host-derived sequences are contaminated as in this case, the efficiency of acquiring target microbe is greatly reduced. Deep sequencing was required to obtain multiple genome sequences to reveal genome diversity, which required wasteful cost and effort.


Here, we suggest a novel approach for targeted single-cell genomics by detecting the target gene using droplet microfluidics. In this study, we identified seven draft genomes of *Endozoicomonas* spp. from *A. tenuis* sampled at four different sites in Okinawa Prefecture, Japan. We then investigated the genomic differences between different clades of *Endozoicomonas* symbiotic with *A. tenuis* coral and examined the role of eukaryotic-like genes. Finally, we proposed a symbiotic strategy model for *Endozoicomonas* bacteria in *A. tenuis* corals, which revealed a diverse adaptation process.

The results of single-cell genome analysis indicate the existence of at least two strategies for microbe-host interactions, indicating the possibility of microbial regulation of host metabolic systems. These findings may provide insights into the robustness of coral reefs.

Materials and methods

Coral branch sampling

A. tenuis samples were collected at four sites, namely Sesoko Minami (26° 53.05' N, 127° 85.77' E), Ishikawabaru (26° 67.51' N, 127° 87.00' E), Onna Village (26° 50.36' N, 127° 85.39' E), and the coral aquaculture center of Yomitan Village (26° 40.90' N, 127° 71.55' E) (Fig. 1A). Official permission to collect coral samples was obtained from Okinawa Prefecture in accordance with Okinawa Prefecture Fishing Regulation. The level of coral bleaching was evaluated on a 3-point scale (0: no bleaching, 1: slight bleaching, 2: complete bleaching). Small pieces of branches (~3 cm) were collected from *A. tenuis* corals and kept in a sterile zipper bag (140×100×0.04 mm; SEISANNIPPONSHA Ltd., Japan) filled with seawater from each sampling site. For 16S rRNA gene amplicon sequencing, coral branches were immediately transferred into a 5-mL tube (Eppendorf Ltd., Germany) containing 4 mL of RNAlater (Thermo Fisher Scientific, Waltham, MA, USA) and then frozen with liquid nitrogen. The frozen samples were kept in a –80°C deep freezer until analysis. For single-cell



genome sequencing, the branches were kept on ice until just before the subsequent process.

DNA extraction and library preparation

The frozen coral branch was thawed on ice. The coral branch was then picked with tweezers and washed with artificial seawater. Then, in a sterilized zipper bag, the coral tissue was blown off using a Waterpik (EW-DJ61-W, Panasonic Corp., Japan), and the suspensions were collected into a 50-mL tube (Nippon Genetics Co., Ltd., Tokyo, Japan). After centrifugation at 10,000×g, 4°C for 30 min, the supernatant was removed, and the pellet was resuspended in 500 µL of artificial seawater. The suspension was transferred into a 1.5-mL tube (SSIBio, Lodi, CA, USA) and centrifuged at 15,000×g, 4°C for 15 min. After the supernatant was removed, the pellet was stored at -80°C until DNA extraction. DNeasy Plant Mini Kit (69104; QIAGEN, Hilden, Germany) was used for DNA extraction according to the manufacturer’s instruction, with a small modification. The thawed pellet was mixed

with 400 µL of Buffer AP1 and transferred into 2-mL screw cap tubes with stabilized 0.1 mm zirconia beads (Yasui Kikai Corp., Osaka, Japan). The tubes were homogenized three times at 2500 rpm for 60 s at 60-s intervals. After centrifugation, 4 µL of RNase A was added, and the sample was incubated at 65°C for 10 min before DNA extraction was performed following the kit protocol.

16S rRNA gene sequencing and analysis

The bacterial community was identified by targeting the variable region V1-V2 of the 16S rRNA gene using the primer set 27F and 338R (27F: 5’-GAG TTT GAT CCT GGC TCA G-3’, 338R: 5’-GCT GCC TCC CGT AGG AGT-3’). PCR amplification of the 16S rRNA gene amplicon (300 bp) was performed in a 25-µL mixture, and the amplicons were sequenced using Ion Torrent PGM with 318 Chip v2 (Thermo Fisher Scientific).

16S rRNA gene analysis was performed using QIIME2 2020.2 [25]. For quality control, amplicon sequence variants (ASVs) were constructed using

DADA2 (via q2-dada2) [26]. All ASVs were aligned using MAFFT (via q2-alignment) [27]. Phylogenetic tree was constructed based on masked aligned ASVs using FastTree2 (via q2-mask and q2-phylogeny) [28]. Taxonomic assignment was performed using the q2-feature-classifier [29] classify-sklearn naïve Bayes taxonomy classifier against the Silva 138 99% OTUs full-length sequences [30]. The downstream analysis was performed using the R bioconductor phyloseq [31] and ggtree [32] in R version 3.5.2.

Droplet digital PCR for quantification of 16S rRNA gene copy number

The PCR mixture contained 10 µL of 2× ddPCR Supermix for Probes (Bio-Rad, Hercules, CA, USA), 1.0 µL of 10 µM forward and reverse primers, 1.0 µL of 10 µM Taqman probe, 1.0 µL of 1 mM Dextran fluorescein (FD10S; Merck KGaA, Darmstadt, Germany), 1.0 µL of extracted DNA, and 5.0 µL of nuclease-free water (Sigma-Aldrich, St. Louis, MO, USA). A total of 20 µL of the PCR mixture was introduced into microfluidic devices (Supplemental Figure 1), and 40 µm of droplets was generated with droplet generation oil for probes (Bio-Rad). The primers and Taqman probes were synthesized by Integrated DNA Technologies (IDT, Inc., Coralville, IA). The sequence of the forward primer, reverse primer, and Taqman probe was 5'-AGA GTT TGA TCM TGG CTC AG-3', 5'-GCT GCC TCC CGT AGG AGT-3', and [6-FAM]-5'-CAG GCC TAA-[ZEN]-CAC ATG CAA GTC-[IBFQ]-3', respectively. The droplets were collected into 0.2-mL tube and then thermal cycled using a T100™ Thermal Cycler (Bio-Rad) with the following protocol: 95°C for 10 min, followed by 40 cycles of 94°C for 30 s and 60°C for 1 min, 95°C for 10 min, and maintenance at 4°C. Subsequently, the droplets were reintroduced into microfluidic devices for digital counting. We have developed a system for calculating the rate of fluorescence-positive droplets using a 488-nm laser (Lambda mini, Tokyo Instruments, Tokyo, Japan) (Supplementary Fig. S1). Fluorescent signals from the droplets were detected by a photosensor module (H11902-01; Hamamatsu Photonics KK, Hamamatsu, Japan) and recorded by an oscilloscope (PICOSCOPE 2204A; Pico Technology Ltd., St. Neots, Cambridgeshire, UK). Combined with the information of droplet diameter measured with ImageJ, 16S rRNA gene copy number per 1 ng of extracted DNA was calculated [33].

Measurement of surface area of coral skeletons by CT scan

The coral branches with surface tissue removed by water picking were used for calculating the surface area. 3D images of the coral branches were obtained using an X-Ray CT scanner (TDM1300-IS; Yamato Scientific,

Tokyo, Japan). The surface area was calculated by the Mimics software (Materialise, Leuven, Belgium). The 16S rRNA gene copy number per 1 cm² of surface area was calculated.

Preparation of bacterial suspension for single-cell genome sequencing

A small piece of coral branch was collected and kept in a 25-mL tube (Iwaki, Tokyo, Japan) filled with seawater from each sampling site. The tube was kept on ice immediately. Each branch was transferred into 5 mL of 0.22-µL-filtered UV-treated seawater of each sampling site and crushed thoroughly by a scalpel (Feather Safety Razor Co. Ltd., Japan). After 5 min of standing on ice, the supernatant was collected into three 1.5-mL tubes. The tube was subsequently centrifuged at 300×g for 5 min, and the supernatant was transferred into a new 1.5-mL tube to remove symbiotic dinoflagellates and other larger particles. The collected supernatant was centrifuged at 8000×g for 5 min. The supernatant was removed, and 500 µL of 0.22-µL-filtered UV-treated seawater was added to resuspend the pellet. The centrifugation at 300×g was repeated until most of the symbiotic dinoflagellates were removed. The centrifugation at 8000×g and the washing steps were repeated three times. Next, the pellet was resuspended in 500 µL of 1x SYBR Green (Thermo Fisher Scientific, Hampton, NH) for 5 min for DNA staining. After centrifugation at 8000×g, the pellet was resuspended in 50 µL of 0.22-µL-filtered UV-treated seawater, and the cell concentration was calculated with a bacterial counter (SLGC, Tokyo, Japan) under a fluorescent microscope (CKX53; Olympus Optical Co Ltd., Tokyo, Japan).

Whole genome amplification and detection of *Endozoicomonas* in gel beads

Cells were encapsulated into 40-µm microfluidic droplets at a concentration of 0.3 cell/droplet using a microfabricated microfluidic device reported previously [22]. The theoretical number of encapsulated cells in a droplet was derived by the Poisson distribution (Formula 1). The percentage of doublets would be less than 3.7% of the total droplets when the cell concentration was 0.3 cell/droplet.

$$p(k, \lambda) = \frac{\lambda^k e^{-\lambda}}{k!} \quad (1)$$

Next, gel-bead-based whole genome amplification (WGA) was conducted according to the previously reported WGA method (single-cell amplified genomes (SAGs) in the gel are referred to as SAG-gel) [21, 34]. After WGA, the gel beads were stained with DAPI, and amplification of the DNA was confirmed with microscopic observation (CKX53; Olympus). Subsequently,

the gel beads were washed and suspended in UV-treated Dulbecco's phosphate-buffered saline (-) (Thermo Fisher Scientific). To detect the target gene sequence, 20 μ L of PCR mixture consisting of 10 μ L of PrimeTime Gene Expression Master Mix (Integrated DNA Technologies, Coralville, IA, USA), 1.0 μ L of each 10 μ M primer (forward and reverse, final concentration 500 nM), 0.5 μ L of 10 μ M Taqman probe (final concentration 250 nM), and 7.5 μ L of gel beads solution was prepared. The sequence of each primer and probe is listed in Supplementary Table 1. On the basis of the 16S rRNA gene amplicon sequencing results, we designed two sets of primers and probes (sets A and B) using the Geneious software (Biomatters Ltd., Auckland, New Zealand). The primer and probe sequences were evaluated with Primer BLAST [35] to confirm that they did not exhibit any specificity to the mitochondrial genome of *A. tenuis*. To detect the *Endozoicomonas* sequence amplified in gel beads, either set A or set B was used. The PCR mixture was reintroduced into microfluidic devices, and 50 μ m of gel beads-containing microfluidic droplets were generated with carrier oil (HFE-7500 oil with 2% surfactant; RAN Biotechnologies Inc., Beverly, MA, USA). The concentration of the beads was adjusted at 0.3 bead/droplet. The droplets were collected into a 0.2-mL tube, and PCR amplification (95°C for 3 min, 25 cycles of 95°C for 5 s and 50°C for 40 s, and maintenance at 10°C) was conducted. After thermal cycling, the fluorescence of the droplets was observed with a microscope (CKX53; Olympus). The droplets exhibiting both DAPI and FAM fluorescence were manually picked with a micro dispenser (Drummond Science Company Broomall, PA, USA) and transferred into a 0.2-mL tube. Next, second-round WGA was performed using a REPLI-g Single Cell Kit (QIAGEN). Buffer D2 (0.6 μ L) was added to each well, followed by incubation at 65°C for 10 min. After that, 8.6 μ L of WGA mixture (0.6 μ L of stop solution, 1.8 μ L of H₂O, 5.8 μ L of reaction buffer, and 0.4 μ L of DNA polymerase) was added, and the mixture was incubated at 30°C for 120 min. The WGA reaction was terminated by heating at 65°C for 3 min. The amplicon yields were quantified by a Qubit dsDNA HS assay kit (Thermo Fisher Scientific). PCR amplification over the 16S rRNA gene was also performed against second-round WGA products. Primer pair sequences for the V3-V4 region were used according to Illumina's MiSeq system protocols (Forward: 5'-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CCT ACG GGN GGC WGC AG-3', Reverse: 5'-GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GGA CTA CHV GGG TAT CTA ATC C-3'). PCR amplification was confirmed with agarose electrophoresis (100 V, 15 min), and the amplicon sequences were obtained via Sanger sequencing (Fasmac, Kanagawa, Japan) to

determine whether the target *Endozoicomonas* sequence was acquired.

Endozoicomonas single-cell genome sequencing and genome assembly

Second-round WGA products from single-cell samples were used for next-generation sequencing library preparation with Nextera XT DNA sample prep kit (Illumina, San Diego, CA, USA) according to the manufacturer's instructions. Each SAG library was sequenced using an Illumina MiSeq 2 \times 75bp or 150bp (Supplementary Table 2).

All raw reads were trimmed to remove adapter sequences by fastp v0.20.0 with default parameters [36]. Trimmed reads were assembled using SPAdes v3.12.0 with "-k auto --sc --careful" parameters [37]. All SAGs were concatenated using ccSAG [38]. SAGs with an average similarity of > 99.9% in single-copy marker genes were co-assembled and used for subsequent analyses. Contigs less than 1000 bp in the SAG were removed. We removed host- and human-derived contaminations, which were annotated as eukaryote by BLAST against the NCBI nt database, from the SAG [39]. The quality of the SAGs was checked using CheckM v1.0.13 [40]. For the subsequent analyses, only high-quality SAG data were used (85% > completeness, < 5% contamination).

Phylogenetic and comparative genome analysis

Gene prediction was performed to estimate protein coding genes (Prodigal v. 2.6.3) [41], tRNAs (ARAGORN v. 1.2.38) [42], and rRNAs (Barrnap v. 0.9) [43] by Prokka v.1.14.5 for the SAGs [44]. Predicted proteins were annotated by diamond v0.9.14 [45] against the NCBI nr database [46]. We also ran InterProScan v.5.40-77.0 [47] and eggNOG-mapper v.1.0.3 [48] to annotate domain information and GO terms and KEGG orthology. Enrichment analysis was performed using the clusterProfiler package [49].

The orthogroup was estimated using OrthoFinder v2.3.0 with the default parameters [50]. Single-copy marker genes were extracted based on the OrthoFinder results. Each gene was aligned by MAFFT v7.407 and trimmed by trimAl v1.4 [27, 51]. A phylogenetic tree was constructed using the single-copy marker genes by IQ-TREE v1.6.7 [52] with ultrafast bootstrap [53] and ModelFinder [54]. Average Nucleotide Identity (ANI) and Average Amino Acid Identity (AAI) were calculated by the all-against-all ANI/AAI matrix calculator [55]. Synteny plots were created by gggenomes [56].

Eukaryotic-like protein analysis

All the Pfam IDs with a Z-score of 10,000 in EffectiveELD version 5.2 were extracted from the InterProScan results [57]. For extracting eukaryotic-like genes based on

sequence homology, genes were selected by taxonomic annotation using eggNOG-mapper v.1.0.3 [48]. In addition, we performed a taxonomy annotation of eukaryotic-like genes using MMseqs2 with “--tax-lineage 1 --vote-mode 0” options, using the eukaryotic-only NCBI nr database [58]. We also checked that these genes are not annotated with other bacteria to confirm that *Endozoicomonas* acquired the genes from eukaryotes during host adaptation using the NCBI nr database without the *Endozoicomonas* genus. Major hits of annotation result were segmented annelid worms *Capitella teleta*. These genes were considered to be derived from contamination and removed from the results.

Results

Diversity analysis of the *Endozoicomonas* genus in *Acropora tenuis*

A. tenuis were collected at four locations (Ishikawabaru, Sesoko Minami, Yomitan, and Onna) in Okinawa in May and August 2016 (Fig. 1A). These four locations are placed within a 15-km radius. Through 16S rRNA gene sequencing, 1418 ASVs were detected as coral-associated bacteria, and 209 ASVs were classified as *Endozoicomonas* spp. In Ishikawabaru, Yomitan, and Onna, *Endozoicomonas* spp. was dominant, representing 70.7–99.7% of the detected bacteria (Fig. 1B). However, in Sesoko Minami, the proportion of *Endozoicomonas* spp. was only 1.8–50.3%, and order Rickettsiales was dominant instead (Supplementary Figure 2A), including Rickettsiales bacteria strain SESOKO1, which was previously reported by our group [59]. The results of principal coordinate analysis (PCoA) based on the Bray-Curtis distance of *Endozoicomonas* composition showed spatial specificity for *A. tenuis* (Supplementary Figure 2B and C).

Phylogenetic analysis of the 16S rRNA gene revealed that the *Endozoicomonas* species can be divided into three major clades: Clade A1 (Red) and Clade A2 (Green) constituted the Onna and Yomitan samples, whereas Clade B (Blue) comprised the Ishikawabaru and Sesoko Minami samples (Fig. 1C, phylogenetic tree and heatmap). Absolute abundance of 16S rRNA gene was calculated by the 16S rRNA gene copy number measured by ddPCR and the surface area value obtained by CT scan. The average (\pm sd) number of *Endozoicomonas* was $37,952 \pm 75,836$ copies/mm² (Ishikawabaru), 2306.8 ± 4241.6 copies/mm² (Sesoko Minami), $30,180 \pm 30,165$ copies/mm² (Onna), and $37,287 \pm 29,083$ copies/mm² (Yomitan), respectively (Fig. 1C, bar plot). These results suggest that the cell numbers of *Endozoicomonas* spp. vary drastically from site to site. Especially at Sesoko Minami, where the copy number of 16S rRNA gene was exceedingly low, the absolute number of *Endozoicomonas*


was significantly lower than that at other sites. In August 2016, large-scale coral bleaching was observed around Okinawa Prefecture; however, the bleaching level of coral colonies at Sesoko Minami was lower than that at Ishikawabaru [60]. In particular, the single coral colony, in which *Endozoicomonas* Clade A1 was dominant, did not show any bleaching in August 2016 (Fig. 1C; star). In our previous study, the density per surface area of zooxanthellae in corals was greatly reduced during coral bleaching in August 2016 [60]. However, in the same coral, the absolute number of *Endozoicomonas* did not decrease significantly from May to August in 2016 (Fig. 1C).

Single-cell genome sequencing of *Endozoicomonas* bacteria

To selectively obtain genomic information of target bacteria from crushed coral tissue suspensions, we have developed a method for targeted single-cell genomics using droplet microfluidics. Bacterial suspensions prepared from coral tissues were encapsulated into agarose gel droplets and processed to gel-bead-based WGA to obtain SAGs [24] (Fig. 2A). Microscopic observation revealed that 24.4–26.3% of the gel beads exhibited DAPI fluorescence, which corresponded to the Poisson distribution value. Subsequently, PCR-based detection of target genes with FAM probes was conducted, which enabled the identification of gel beads containing target SAGs (Fig. 2B). After PCR amplification, there were droplets displaying both DAPI and FAM fluorescence (Fig. 2C). The percentage of DAPI fluorescence-positive droplets is 0.8–1.2%, which represents the rate of droplets encapsulating the target bacterium.

The results of 16S rRNA gene sequencing followed by second-round WGA showed that 64% (28 out of 44) of the samples corresponded to the *Endozoicomonas* sequence. Among the remaining 16 samples, 13 exhibited insufficient DNA yield (< 2 ng/ μ L) after the second-round WGA. The remaining three were confirmed to be derived from host mitochondria. When droplets that were positive only for DAPI were isolated, all the sequence corresponded to the sequence of host mitochondria ($n = 30$), suggesting that our targeted single-cell genomics has a low false-negative rate. In order to obtain 28 target bacterial SAGs by random sampling, 2333–3500 droplets need to be isolated according to the calculation, which means that our method showed 53.0–79.5 times higher screening efficiency than the random sampling.

As a result, seven draft SAGs ranging from 2.1 to 6.4 Mbp were acquired from four sampling sites, three of which were classified as high-quality and three of which were classified as medium-quality according to the criteria by the Genomic Standards Consortium (GSC) (Table 1).



Comparative genome analysis of *Endozoicomonas* genus

We conducted a comparative genome analysis using our five draft SAGs, which include four from this study and one from our previous report [60], collected from *A. tenuis*, and 10 public *Endozoicomonas* isolates collected from various kinds of marine organisms. Phylogenetic trees constructed with single-copy marker genes showed that *Endozoicomonas* spp. represented single clades that corresponded to symbiotic host organisms. However, *Endozoicomonas* spp. collected from *Acropora* coral (*Endozoicomonas* sp. ISH11, SESOKO1, ONNA1, ONNA2, and YOMI1) were divided into two clades (clade A1 and B) (Fig. 3A), which was also confirmed by the ANI and average AAI (Supplementary Figure 3).


A similar trend was observed in the ortholog analysis. Out of a total of 9414 orthogroups, the number of orthogroups commonly detected in all *Endozoicomonas* spp. was 1148. We confirmed that the bacterial secretion systems are evolutionarily conserved in all *Endozoicomonas* symbiotic relationships with marine organisms (Supplementary Table 3), which are important

for *Endozoicomonas* to secrete proteins related to eukaryotic pathways in the host. The prediction of effector proteins showed that 12.5–20.4% (T3SS) and 2.97–4.22% (T4SS) of proteins were released by bacterial secretion systems (Supplementary Table 4). On the other hand, the remaining 8266 orthogroups and 8061 unassigned genes were clade- or strain-specific, suggesting that *Endozoicomonas* spp. possess a diverse orthogroup profile. It appears that the profile of the orthogroups depends on the host of each *Endozoicomonas* sp. However, clade A1 and clade B, which were symbiotic with the same host (*A. tenuis*), shared no unique orthogroups.

Enrichment analysis of accessory genes to the KEGG pathway revealed the functional characteristics and differences of each clade (clade A1 and clade B). Of the top ten enriched pathways (nine pathway in Clade B) of each clade, eight pathways (except bacterial chemotaxis and flagellar assembly) in clade A1 as well as seven pathways (except sphingolipid metabolism and terpenoid backbone biosynthesis) in clade B were assigned as eukaryotic pathways (Fig. 3B, C). Primarily,

Table 1 Overview of *Endozoicomonas*

Strains	The number of seqs	Sum length (nt)	GC content (%)	Competeness	Contamination	Stran heterogeneity (>90%)	Method	Host	References
<i>Endozoicomonas acroporae</i> AcI-14T	309	6,048,850	49.35	98.56	1.39	0	Cultured	<i>Acropora</i> sp.	(Tandon, K et al, 2018) [61]
<i>Endozoicomonas arenosclerae</i> ab112	328	6,453,554	46.45	99.14	2.05	16.67	Cultured	<i>Arenosclera brasiliensis</i>	(Applinario LR et al, 2016) [62]
<i>Endozoicomonas ascidicola</i> AVMART05	36	6,130,497	38.38	98.28	1.81	0	Cultured	<i>Ascidifella</i> sp.	(Schreiber L et al, 2016) [63]
<i>Endozoicomonas ascidicola</i> KASP37	34	6,512,467	40.18	97.63	2.02	0	Cultured	<i>Ascidifella</i> sp.	(Schreiber L et al, 2016) [63]
<i>Endozoicomonas atrinae</i> WP70	980	6,687,418	48.86	98.92	3.43	0	Cultured	<i>Atrina pectinata</i>	(Hyun DW et al, 2014) [7]
<i>Endozoicomonas elysicola</i> DSM 22380	2	5,606,375	44.64	98.98	3.43	0	Cultured	<i>Elysia ornata</i>	(Neave NJ et al, 2014) [64]
<i>Endozoicomonas montiporae</i> CL-33	1	5,430,256	48.37	98.99	1.72	0	Cultured	<i>Montipora aequituberculata</i>	(Ding J-Y et al, 2016) [14]
<i>Endozoicomonas montiporae</i> LMG 24815	20	5,602,297	47.26	98.99	1.72	0	Cultured	<i>Montipora aequituberculata</i>	(Neave NJ et al, 2014) [64]
<i>Endozoicomonas numazuensis</i> DSM 25634	31	6,342,227	43.12	99.14	1.66	12.5	Cultured	Sponge (order Haplosclerida)	(Neave NJ et al, 2014) [64]
<i>Endozoicomonas</i> sp.OPT23	30	4,938,102	45.63	99.14	1.94	0	Cultured	<i>Ophilitaspongia papilla</i>	(Alex A et al, 2019) [10]
<i>Endozoicomonas</i> sp.YOM11	603	4,656,793	47.99	98.52	1.3	0	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. SESOKO1	996	5,532,550	47.14	90.73	1.49	11.11	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. ONNA1	1,217	4,681,217	48.3	87.97	1.73	0	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. ONNA2	415	4,220,889	49.63	95.02	1.79	11.11	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. SESOKO2	821	4,388,010	47.44	81.65	1.59	33.33	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. SESOKO3	808	3,194,186	47.44	59.41	2.05	16.67	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. SESOKO4	687	2,947,897	47.29	49.22	0.26	33.33	Single cell	<i>Acropora tenuis</i>	(This study)
<i>Endozoicomonas</i> sp. ISH11	144	6,454,849	48.58	98.49	1.06	0	Cultured	<i>Acropora tenuis</i>	(Maruyama T et al, 2021) [60]



in *Endozoicomonas* clade A1, immune-related pathways were enriched, such as the TRAF signaling pathways and NF- κ B pathways (Fig. 3B), which are generally not conserved in bacteria. Conversely, in clade

B, these immune-related pathways were not enriched, but proteasomes and proteolysis pathways including eukaryotic-like E3 ubiquitin ligase gene were enriched (Fig. 3C).




Fig. 4 Eukaryotic-like genes distribution and horizontal gene transfer in *Endozoicomonas*. **A** Properties of eukaryotic-like domains predicted using EffectiveDB in *Endozoicomonas* (Z-score = 10,000). The phylogenetic tree based on the single-copy marker gene is the same as that in Fig. 3A. The hierarchical clustering at the top is based on the profiles of eukaryotic-like genes. **B** Insertion of the inhibitor of apoptosis domain-containing genes of *Endozoicomonas* sp. YOMI1 and *Endozoicomonas acroporae* Acr-14 into the genome. The inserted eukaryotic-like gene is marked by a yellow square. The region where the GC content of *Endozoicomonas ascidiella* DSM22380 is flat is the gap region (N)

Different symbiotic strategies utilizing eukaryotic-like genes
 We further conducted a comparative genome analysis focusing on eukaryotic-like domains and found that all *Endozoicomonas* spp. used in this study had at least one eukaryotic-like domain (Fig. 4A). The types of eukaryotic-like domains were different between clade A1 (YOMI1, ONNA1, and ONNA2) and clade B (ISH1 and SESEKO1) *Endozoicomonas*; namely, the eukaryotic-like

domain found in common between clade A1 and clade B was only RING-type zinc finger (RING-type zinc-finger, Zinc-finger double domain, Zinc finger, and C3HC4 type were grouped), and the remaining six domains (SOCS box, Inhibitor of Apoptosis domain, ephrin ligand, Ets domain, MATH domain, and FANCL C-terminal domain) were found only in either species (Fig. 4A, Supplementary Table 5). We have previously reported that

Endozoicomonas sp. ISH11 has coral-like ephrin ligand genes [60]. In addition, there were other eukaryotic-like genes (such as the SNARE domain, Vps4 C terminal oligomerization domain-containing genes). In contrast, *Endozoicomonas* sp. YOM11, ONNA1, and ONNA2, which belong to clade A1, did not possess ephrin ligand, but had eukaryotic-like genes, including MATH/TRAF, SOCS box, an Ets-domain, and an inhibitor of apoptosis domain (Fig. 4A).

Furthermore, we found 532 genes that did not contain eukaryotic-like domains but were highly homologous against eukaryote in *A. tenuis*-symbiotic *Endozoicomonas* (YOM11: 125 genes, ONNA1: 134 genes, ONNA2: 47 genes, SESOKO1: 90 genes, ISH11: 136 genes) (Supplementary Table 6). As genes closely related to host *A. tenuis* coral genes, TRAF/MATH domain-containing genes (match: 26.0–62.2%), low-density lipoprotein receptor gene (match: 30.9–67.6%), and transcription factor 15-like gene (match: 27.5–30.4%) were detected in *Endozoicomonas* clade A1. In contrast, the genes for ephrin ligand (match: 31.3–46.0%) and RING finger and CHY zinc finger domain-containing gene (match: 46.1–64.3%) were detected in clade B. Eukaryotic-like E3 ubiquitin-protein ligase was detected as a common gene in both clades (Supplementary Table 5). Some of these eukaryotic-like genes were inserted into the genome (e.g., survivin gene, ephrin gene) (Fig. 4B). Furthermore, these genes did not contain introns, despite their similarity to the intron-containing host genes (Supplementary Figure 4).

Discussion

The absolute abundance of *Endozoicomonas* varies greatly among coral habitats

Previous studies have focused solely on bacterial composition and failed to establish a relationship between the amount of coral symbiotic bacteria and their host conditions. To overcome this limitation, this study quantified the absolute 16S rRNA gene copy number of bacteria and revealed the abundance of *Endozoicomonas* spp., which varied by more than 72-fold among sampling sites. Several researchers have considered that the abundance of *Endozoicomonas* is related to health and robustness of corals [65, 66]. However, in August 2016, corals from Ishikawabaru, which had a higher abundance of *Endozoicomonas* than Sesoko Minami, showed far extensive bleaching, which was not consistent with their findings [67]. Our previous studies also showed that inflammation-related genes are upregulated in *A. tenuis*, which is occupied by clade B *Endozoicomonas* spp. In this study, we measured quantitative *Endozoicomonas* abundance, which may be involved in the host side response at the gene expression level. The individual

adaptive capacity of *Endozoicomonas* may also have an effect on the host [68]. For a deeper understanding of *Endozoicomonas* function in *Acropora* coral, the difference of the clade or absolute amount of *Endozoicomonas* should be considered.

Targeted single-cell genomics enables selective accumulation of low-abundance bacterial genomes

In addition to the 16S rRNA gene-based phylogenetic analysis, we conducted single-cell whole genome sequencing of target bacteria and collected seven SAGs belonging to clade A1 (*Endozoicomonas* sp. YOM11, ONNA1, and ONNA2) and clade B (*Endozoicomonas* sp. SESOKO1 and ISH11). Although SAGs consisted of fragmented contigs, and the total length of some SAGs were shorter compared to the genome size of reference *Endozoicomonas*, the completeness was comparable to cultured strains reported previously [15], providing evidence that this method can expand genomic information of uncultured bacteria. It is noteworthy that our method enabled selective acquisition of draft SAGs even when the abundance ratio of target bacteria was approximately 1%, thereby overcoming the disadvantage of metagenomics. Moreover, even in the conventional single-cell genomics, the composition of sequenced bacterial taxa is highly dependent on the proportion of bacteria in the target environment because single cells are randomly sorted [69]. In contrast, in our proposed method, WGA was followed by detection of target gene sequence (*Endozoicomonas* 16S rRNA gene) in gel beads to achieve selective isolation and subsequent genome analysis. To the best of our knowledge, this is the first report of targeted single-cell genomics on environmental microbes. Our method showed > 50 times higher screening efficiency of target bacteria than the random sampling, thus drastically reducing the amount of reagents and labor wasted on non-target samples. Our targeted single-cell genomics can also be used to detect specific gene sequences obtained from metagenomic shotgun analysis or 16S rRNA gene sequencing by designing specific primers and probes. As a further application, specific single-cell genomics with unique characteristics will be possible by targeting the sequences of biosynthetic gene clusters and drug-resistant genes. As the potential of rare bacteria to exert important functions in microbial populations is gaining increasing attention [70], our method can be a technological innovation and breakthrough for the analysis of single-cell genomics. In the single-cell genomics of environmental bacteria, it is often important to process the samples immediately after collection to obtain high-quality draft SAGs. Because the microfluidic system used in this study is portable, on-site single-cell isolation and genome amplification can be performed with standard

biological laboratory equipment, which is critical for obtaining high-quality SAGs.

Diversity of host adaptation strategies in *Endozoicomonas*


Several 16S rRNA gene amplicon and shotgun metagenomic analyses have suggested that coral-associated *Endozoicomonas* spp. may improve host coral health [65, 66]. However, owing to the lack of reference genomes, detailed analysis of clade-specific functions in *Endozoicomonas* in the same coral species has not been performed. Therefore, we used our SAG data and compared the gene functions detected in each clade of *Endozoicomonas*. Comparative genome analysis revealed that the orthogroups of *Endozoicomonas* were conserved in each clade. Notably, even when the different clades of *Endozoicomonas* coexisted with *A. tenuis*, the shared orthogroups were only the core gene between clade A1 (*Endozoicomonas* sp. YOMI1, ONNA1, and ONNA2) and clade B (*Endozoicomonas* sp. ISHI1 and SESOKO1). These results suggest that *Endozoicomonas* independently adapts to its host during the process of evolution.

A comprehensive search for eukaryotic-like genes in *Endozoicomonas* revealed that various eukaryotic-like genes were enriched in a lineage-specific manner. Symbiotic bacteria are known to harbor eukaryotic-like domains, such as ankyrin repeat and WD40 repeat, which are involved in protein-protein interactions and are thought to be necessary for interaction with the host [71, 72]. For example, in plants and sponges, many symbiotic bacteria and pathogens intervene in host signaling pathways via eukaryotic-like genes that mimic host genes, thereby influencing host conditions [73, 74]. There have been few reports on how symbiotic bacteria acquire these eukaryotic-like genes, but in our study, some of these genes (e.g., ephrin ligand and survivin) were found to be inserted with spliced host coral genes (Supplementary Figure 4). This result suggests the bacterial-eukaryotic horizontal gene transfer is observed between host coral and coral-associated *Endozoicomonas* spp. One of many possible hypotheses is that the insertion of these genes may be mediated by host mRNA. Horizontal gene transfer between eukaryotes and prokaryotes has been studied in recent years, and more detailed studies are needed in the future.

A recent study reported that the copy number of ankyrin repeats is higher in *E. acroporae* genomes than in other marine organism-associated *Endozoicomonas* genomes [10], implying that some eukaryotic-like genes play essential roles in the symbiosis of *Endozoicomonas* with *Acropora* coral. On the basis of this idea, we hypothesized the functions and host adaptation

strategies of each *Endozoicomonas* clade based on the functions of enriched eukaryotic-like domains. Recently, it was hypothesized that ephrin ligands are required for *E. montiporae* to transition the symbiotic mechanisms [14]. In our previous report, we have also confirmed a gene expansion of ephrin ligand in *Endozoicomonas* Clade B (*Endozoicomonas* sp. ISHI1 and SESOKO1) with a higher degree than that in *E. montiporae* [60]. Interestingly, this is consistent with the fact that the number of ephrin-like receptor genes is expanded in host *Acropora* coral compared with that in *Montipora* coral [75]. These results reinforce the hypothesis that *Endozoicomonas* uses ephrin ligand gene as a part of its adaptation strategy to its host coral. The consistency of ephrin ligand and host ephrin receptor gene expansion suggests the co-evolution between *Endozoicomonas* and host coral. Furthermore, *Endozoicomonas* clade B encodes genes associated with intracellular invasion, such as SNARE domains, which contribute to the inhibition of degradation by phagosomes in *Chlamydia* and *Legionella* [76]. However, *Endozoicomonas* clade B does not encode immune-related genes. This result is consistent with our recent finding that *Endozoicomonas* sp. ISHI1 stimulates immune-related pathways in corals through infection (Fig. 5A) [60]. In addition, genes containing the MATH/TRAF, SOCS box, and inhibitor of apoptosis domain were present in clade A1, but not in clade B. These genes suppress apoptosis induced by the innate immune pathway, such as the NF- κ B and JAK-STAT pathways [77]. Recently, several studies reported that the innate immune pathways are activated in host corals as stress responses to bleaching [78]. In particular, the NF- κ B pathway is conserved in cnidarians and has attracted much attention in the analysis of bleaching [79–81]. Therefore, we speculate that a group of eukaryotic-like genes that are significantly enriched in *Endozoicomonas* clade A1 may affect host innate immune pathways and increase tolerance to environmental stress.

Homology-based analysis also revealed several eukaryotic-like genes with unknown functions in *Endozoicomonas* clade A1 and B. These genes are phylogenetically conserved and may have some function in host adaptation and symbiosis. One example is the low-density lipoprotein receptor (LDLR) gene conserved in clade A1. Recently, the function of LDLR during oogenesis in *A. tenuis* has been reported; it has been shown that the transcript of AtLDLR is present in the mesentery, mesenteric membrane, and mesenteric filaments surrounding an oocyte, but not in the membrane of a developing oocyte. AtLDLR can bind to major yolk lipoproteins and may be transported into the cytoplasm of an oocyte as it develops [82].



This indicates that the LDLR-like gene of *Endozoicomonas* clade A1 has the binding function to host egg lipoprotein. Therefore, we speculate that the LDLR-like gene, which is involved in egg development, may contribute to vertical transfer of *Endozoicomonas*, as other infection-related genes such as ephrin ligand gene and SNARE gene have not been identified in clade A1. The speculation can be verified by future experiments.

Conclusions

We succeeded in the targeted single-cell genomics of *Endozoicomonas* from *A. tenuis* corals with droplet microfluidics. Thus, we accomplished obtaining high-quality *Endozoicomonas* SAGs (clade A1 and B) from the same coral species, which were proved to be phylogenetically and functionally diverse. Comparative genome analysis revealed the difference in the eukaryotic-like gene profiles, suggesting that the symbiotic strategy of *Endozoicomonas* in *A. tenuis* is divided into two types. Our findings suggest that even in the same bacterial genus, each bacterial lineage has different gene profiles and functions. Therefore, it is necessary to conduct strain-level genome sequencing to analyze the functions and strategies for

host adaptation. Although in vitro studies in the field environment are required to prove this hypothesis, our targeted single-cell genomics could be a powerful tool for acquiring SAGs of uncultured bacterium to infer its function. Analysis of various coral-associated *Endozoicomonas* SAGs will clarify their host adaptation strategies in more detail and shed light on some aspects of the symbiotic mechanisms.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40168-022-01395-9>.

Additional file 1: Supplementary Figure 1. Droplet Digital PCR to quantify 16S rRNA gene copy number of bacteria. **Supplementary Figure 2.** Order level barplot and PCoA plot based on UniFrac distance of *Endozoicomonas* genus in four sampling points. **Supplementary Figure 3.** Average Nucleotide Identity (ANI) and Average Amino Acid Identity (AAI) of *Endozoicomonas* genomes. **Supplementary Figure 4.** Comparison of gene structures of Coral-like ephrin ligand genes.

Additional file 2: Table S1. List of Primer sets used to identify *Endozoicomonas*. **Table S2.** Raw read statistics of our acquired *Endozoicomonas* genus genome. **Table S3.** List of bacterial secretion systems conserved in *Endozoicomonas*. **Table S4.** Percentage of proteins released by the bacterial secretion system. **Table S5.** List of eukaryotic-like domains in *Endozoicomonas*. **Table S6.** List of all eukaryotic-like genes in *Endozoicomonas*.

Acknowledgements

We would like to thank Dr. Mitsuo Umezu for CT scan analysis, Naoko Okada for 16S rRNA gene sequencing, and Dr. Hisashi Anbutsu and Eisuke Iwamoto for useful suggestions. The super-computing resource was provided by the Human Genome Center (University of Tokyo). This study was supported by the Collaborative Research of Tropical Biosphere Research Center, University of the Ryukyus.

Authors' contributions

M.I., M.H., S.S., and H.T. designed the experiments. K.I., Y.Ni., Y.T., K.Y., and H.T. wrote all the manuscripts. M.I., K.I., Y.Ni., H.I., K.K., and Y.Na. collected the data. Y.Ni. and M.K. conducted the single-cell genomics experiments for obtaining *E. sp.* YOMI1, SESOKO1-4, and ONNA1-2 SAGs. H.T., R.M., and H.I. conducted the ddPCR analysis. H.I. conducted CT scan analysis. K.I., M.K., R.W., and T.M. performed the bioinformatics analysis. All authors have read and approved the final manuscript.

Funding

This work was supported by the "Construction of the environmental risk mathematical model by the meta-omics analyses of marine unculturable microbes based on single cell genome information" grant (JPMJCR12A4) from JST-CREST and by JSPS KAKENHI grant number JP17H06158.

Availability of data and materials

The assembled genomes of seven *Endozoicomonas* were deposited in DDBJ/ENA/GenBank under BioProject number PRJNA721663. The raw read data of 16S rRNA gene are available under BioProject number PRJNA722798.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

M.H. and H.T. are shareholders in bitBiome, Inc., which provides single-cell genomics services using the SAG-gel workflow as bit-MAP. K.I., Y.Ni., K.M., M.H., and H.T. are inventors on patent applications covering the technique for single-cell sequencing.

Author details

¹Department of Life Science and Medical Bioscience, Waseda University, Tokyo, Japan. ²Computational Bio Big-Data Open Innovation Laboratory (CBDD-OIL), National Institute of Advanced Industrial Science and Technology, Tokyo, Japan. ³Research Organization for Nano and Life Innovation, Waseda University, Tokyo, Japan. ⁴Tropical Biosphere Research Center, University of the Ryukyus, Okinawa, Japan. ⁵Marine Science Section, Research Support Division, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan. ⁶Umino-Tane Co., Ltd., Okinawa, Japan. ⁷Institute for Advanced Research of Biosystem Dynamics, Waseda Research Institute for Science and Engineering, Tokyo, Japan. ⁸Graduate School of Humanities and Sciences, Ochanomizu University, Tokyo, Japan. ⁹Faculty of Science, University of the Ryukyus, Okinawa, Japan.

Received: 11 May 2022 Accepted: 13 October 2022

Published online: 12 December 2022

References

- Muscatine L, Porter JW. Reef corals: mutualistic symbioses adapted to nutrient-poor environments. *Bioscience*. 1977;27:454–60.
- Liu H, Stephens TG, González-Pech RA, Beltran VH, Lapeyre B, Bongaerts P, et al. Symbiodinium genomes reveal adaptive evolution of functions related to coral-dinoflagellate symbiosis. *Commun Biol*. 2018;1:95.
- Neave MJ, Rachmawati R, Xun L, Michell CT, Bourne DG, Apprill A, et al. Differential specificity between closely related corals and abundant *Endozoicomonas* endosymbionts across global scales. *Isme J*. 2016;11:186–200.
- Lema KA, Willis BL, Bourne DG. Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Appl Environ Microb*. 2012;78:3136–44.
- Raina J-B, Tapiolas D, Motti CA, Foret S, Seemann T, Tebben J, et al. Isolation of an antimicrobial compound produced by bacteria associated with reef-building corals. *Peerj*. 2016;4:e2275.
- Cano I, van Aerle R, Ross S, Verner-Jeffreys DW, Paley RK, Rimmer GSE, et al. Molecular characterization of an *Endozoicomonas*-like organism causing infection in the king scallop (*Pecten maximus* L.). *Appl Environ Microb*. 2018;84:e00952–17.
- Hyun DW, Shin NR, Kim MS, Oh SJ, Kim PS, Whon TW, et al. *Endozoicomonas atrinae* sp. nov., isolated from the intestine of a comb pen shell *Atrina pectinata*. *Int J Syst Evol Micr*. 2014;64:2312–8.
- Kurahashi M, Yokota A. *Endozoicomonas elysicola* gen. nov., sp. nov., a γ-proteobacterium isolated from the sea slug *Elysia ornata*. *Syst Appl Microbiol*. 2007;30:202–6.
- Nishijima M, Adachi K, Katsuta A, Shizuri Y, Yamasato K. *Endozoicomonas numazuensis* sp. nov., a gammaproteobacterium isolated from marine sponges, and emended description of the genus *Endozoicomonas* Kurahashi and Yokota 2007. *Int J Syst Evol Micr*. 2013;63:709–14.
- Alex A, Antunes A. Comparative genomics reveals metabolic specificity of *Endozoicomonas* isolated from a marine sponge and the genomic repertoire for host-bacteria symbioses. *Microorg*. 2019;7:635.
- Du Z, Zhang W, Xia H, Lü G, Chen G. Isolation and diversity analysis of heterotrophic bacteria associated with sea anemones. *Acta Oceanol Sin*. 2010;29:62–9.
- Schuett C, Doepke H, Grathoff A, Gedde M. Bacterial aggregates in the tentacles of the sea anemone *Metridium senile*. *Helgoland Mar Res*. 2007;61:211–6.
- Glasl B, Smith CE, Bourne DG, Webster NS. Disentangling the effect of host-genotype and environment on the microbiome of the coral *Acropora tenuis*. *Peerj*. 2019;7:e6377.
- Ding J-Y, Shiu J-H, Chen W-M, Chiang Y-R, Tang S-L. Genomic insight into the host–endosymbiont relationship of *Endozoicomonas montiporae* CL-33T with its coral host. *Front Microbiol*. 2016;7(e1000376):e1000376–15.
- Tandon K, Lu C-Y, Chiang P-W, Wada N, Yang S-H, Chan Y-F, et al. Comparative genomics: Dominant coral-bacterium *Endozoicomonas acroporae* metabolizes dimethylsulfoniopropionate (DMSP). *Isme J*. 2020;14:1290–303.
- Neave MJ, Michell CT, Apprill A, Voolstra CR. *Endozoicomonas* genomes reveal functional adaptation and plasticity in bacterial strains symbiotically associated with diverse marine hosts. *Sci Rep-uk*. 2017;7(1):1–12.
- Yue Y, Huang H, Qi Z, Dou H-M, Liu X-Y, Han T-F, et al. Evaluating metagenomics tools for genome binning with real metagenomic datasets and CAMI datasets. *BMC Bioinformatics*. 2020;21:334.
- Quince C, Delmont TO, Raguideau S, Alneberg J, Darling AE, Collins G, et al. DESMAN: a new tool for de novo extraction of strains from metagenomes. *Genome Biol*. 2017;18:181.
- van de Water JAJM, Allemand D, Ferrier-Pagès C. Host-microbe interactions in octocoral holobionts - recent advances and perspectives. *Microbiome*. 2018;6:64.
- Feehery GR, Yigit E, Oyola SO, Langhorst BW, Schmidt VT, Stewart FJ, et al. A method for selectively enriching microbial DNA from contaminating vertebrate host DNA. *Plos One*. 2013;8:e76096.
- Schmieder R, Edwards R. Fast identification and removal of sequence contamination from genomic and metagenomic datasets. *Plos One*. 2011;6:e17288.
- Nishikawa Y, Hosokawa M, Maruyama T, Yamagishi K, Mori T, Takeyama H. Monodisperse picoliter droplets for low-bias and contamination-free reactions in single-cell whole genome amplification. *Plos One*. 2015;10:e0138733.
- Hosokawa M, Nishikawa Y, Kogawa M, Takeyama H. Massively parallel whole genome amplification for single-cell sequencing using droplet microfluidics. *Sci Rep-uk*. 2017;7:5199.
- Chijiwa R, Hosokawa M, Kogawa M, Nishikawa Y, Ide K, Sakanashi C, et al. Single-cell genomics of uncultured bacteria reveals dietary fiber responders in the mouse gut microbiota. *Microbiome*. 2020;8:5.
- Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol*. 2019;37:852–7.

26. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods*. 2016;13:581–3.
27. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res*. 1911;30:3059–66.
28. Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for large alignments. *Plos One*. 2010;5:e9490.
29. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome*. 2018;6:90.
30. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res*. 2013;41:D590–6.
31. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *Plos One*. 2013;8:e61217.
32. Yu G, Smith DK, Zhu H, Guan Y, Lam TT. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol Evol*. 2017;8:28–36.
33. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9:671–5.
34. Nishikawa Y, Kogawa M, Hosokawa M, Wagatsuma R, Mineta K, Takahashi K, Ide K, Yura K, Behzad H, Gojobori T, Takeyama H. Validation of the application of gel beads-based single-cell genome sequencing platform to soil and seawater. *ISME commun*. 2022;2:92.
35. Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*. 2012;13:134.
36. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*. 2018;34:i884–90.
37. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biology J Comput Mol Cell Biol*. 2012;19:455–77.
38. Kogawa M, Hosokawa M, Nishikawa Y, Mori K, Takeyama H. Obtaining high-quality draft genomes from uncultured microbes by cleaning and co-assembly of single-cell amplified genomes. *Sci Rep-uk*. 2018;8:2059.
39. McGinnis S, Madden TL. BLAST: at the core of a powerful and diverse set of sequence analysis tools. *Nucleic Acids Res*. 2004;32:W20–5.
40. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res*. 2015;25:1043–55.
41. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*. 2010;11:119–11.
42. Laslett D. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res*. 2004;32:11–6.
43. Seemann T. barrnap 0.9: rapid ribosomal RNA prediction. <https://github.com/tseemann/barrnap>.
44. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. 2014;30:2068–9.
45. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat Methods*. 2015;12:59–60.
46. Coordinators NR, Agarwala R, Barrett T, Beck J, Benson DA, Bollin C, et al. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res*. 2017;46:D8–D13.
47. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, et al. InterPro-Scan 5: genome-scale protein function classification. *Bioinformatics*. 2014;30:1236–40.
48. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, et al. Fast genome-wide functional annotation through orthology assignment by eggNOG-Mapper. *Mol Biol Evol*. 2017;34:2115–22.
49. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics J Integr Biology*. 2012;16:284–7.
50. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol*. 2019;20:238.
51. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. 2009;25:1972–3.
52. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol Biol Evol*. 2020;37:1530–4.
53. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBBoot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol*. 2017;35:518–22.
54. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermini LS. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*. 2017;14:587–9.
55. Rodriguez-RLM, Konstantinidis KT. The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *PeerJ Preprints*. 2016;4:e1900v1.
56. Thomas Hackl and Markus J. Ankenbrand (2020). gggenomes: a grammar of graphics for comparative genomics. R package version 0.0.0.9000. <https://github.com/thackl/gggenomes>
57. Eichinger V, Nussbaumer T, Platzer A, Jehl M-A, Arnold R, Rattei T. EffectiveDB—updates and novel features for a better annotation of bacterial secreted proteins and Type III, IV, VI secretion systems. *Nucleic Acids Res*. 2016;44:D669–74.
58. Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol*. 2017;35:1026–8.
59. Ide K, Nishikawa Y, Kogawa M, Iwamoto E, Samuel AZ, Nakano Y, et al. High-quality draft genome sequence of a Rickettsiales bacterium found in *Acropora tenuis* coral from Okinawa, Japan. *Microbiol Resour Announc*. 2020;9:e00848–20.
60. Maruyama T, Ito M, Wakaoji S, Okubo Y, Ide K, Fujimura H, et al. Multi-omics analysis highlights cross-organism interactions in coral holobiont. *bioRxiv* 2021:10.25.465660. <https://doi.org/10.1101/2021.10.25.465660>.
61. Tandon K, Chiang P-W, Chen W-M, Cang S-L. Draft Genome Sequence of *Endozoicomonas acroporae* Strain Acr-14T, Isolated from *Acropora* Coral. *Genome Announc*. 2018;6:e01576–17.
62. Appolinario LR, Tschoeke DA, Rua CPJ, Venas T, Campeão ME, Amaral GRS, et al. Description of *Endozoicomonas arenosclerae* sp. nov. using a genomic taxonomy approach. *Antonie Van Leeuwenhoek*. 2016;109:431–8.
63. Schreiber L, Kjeldsen KU, Obst M, Funch P, Schramm A. Description of *Endozoicomonas ascidiicola* sp. nov., isolated from Scandinavian ascidians. *Syst Appl Microbiol*. 2016;39:313–8.
64. Neave MJ, Michell CT, Apprill A, Voolstra CR. Whole-Genome Sequences of Three Symbiotic *Endozoicomonas* Strains. *Genome Announc*. 2014;2:e00802–14.
65. Bourne D, Iida Y, Uthicke S, Smith-Keune C. Changes in coral-associated microbial communities during a bleaching event. *Isme J*. 2008;2:350–63.
66. Pogoreutz C, Rädicker N, Cárdenas A, et al. Dominance of *Endozoicomonas* bacteria throughout coral bleaching and mortality suggests structural inflexibility of the Pocillopora verrucosa microbiome. *Ecol Evol*. 2018;8:2240–52.
67. Neave MJ, Apprill A, Ferrier-Pagès C, Voolstra CR. Diversity and function of prevalent symbiotic marine bacteria in the genus *Endozoicomonas*. *Appl Microbiol Biot*. 2016;100:8315–24.
68. Tandon K, Chiou Y-J, Yu S-P, Hsieh HJ, Lu C-Y, Hsu M-T, et al. Microbiome restructuring: dominant coral bacterium *Endozoicomonas* species respond differentially to environmental changes. *Msystems*. 2022;7:e00359-22.
69. Lan F, Demaree B, Ahmed N, Abate AR. Single-cell genome sequencing at ultra-high-throughput with microfluidic droplet barcoding. *Nat Biotechnol*. 2017;35:640–6.
70. Lynch MDJ, Neufeld JD. Ecology and exploration of the rare biosphere. *Nat Rev Microbiol*. 2015;13:217–29.
71. Mosavi LK, Cammett TJ, Desrosiers DC, Peng Z. The ankyrin repeat as molecular architecture for protein recognition. *Protein Sci*. 2004;13:1435–48.
72. Jain BP, Pandey S. WD40 repeat proteins: signalling scaffold with diverse functions. *Protein J*. 2018;37:391–406.
73. Diez-Vives C, Moitinho-Silva L, Nielsen S, Reynolds D, Thomas T. Expression of eukaryotic-like protein in the microbiome of sponges. *Mol Ecol*. 2017;26:1432–51.
74. Levy A, Gonzalez IS, Mittelviehhaus M, Clingenpeel S, Paredes SH, Miao J, et al. Genomic features of bacterial adaptation to plants. *Nat Genet*. 2018;50:138–50.

75. Cuning R, Bay RA, Gillette P, Baker AC, Traylor-Knowles N. Comparative analysis of the Pocillopora damicornis genome highlights role of immune system in coral evolution. *Sci Rep-uk*. 2018;8:16134.
76. Paumet F, Wesolowski J, Garcia-Diaz A, Delevoye C, Aulner N, Shuman HA, et al. Intracellular bacteria encode inhibitory SNARE-like proteins. *Plos One*. 2009;4:e7375–6.
77. Wong BR, Josien R, Lee SY, Vologodskaja M, Steinman RM, Choi Y. The TRAF family of signal transducers mediates NF- κ B activation by the TRANCE receptor*. *J Biol Chem*. 1998;273:28355–9.
78. Pinzón JH, Kamel B, Burge CA, Harvell CD, Medina M, Weil E, et al. Whole transcriptome analysis reveals changes in expression of immune-related genes during and after bleaching in a reef-building coral. *Roy Soc Open Sci*. 2015;2:140214.
79. Williams LM, Fuess LE, Brennan JJ, Mansfield KM, Salas-Rodríguez E, Welsh J, et al. A conserved Toll-like receptor-to-NF- κ B signaling pathway in the endangered coral *Orbicella faveolata*. *Dev Comp Immunol*. 2018;79:128–36.
80. Mansfield KM, Carter NM, Nguyen L, Cleves PA, Alshanbayeva A, Williams LM, et al. Transcription factor NF- κ B is modulated by symbiotic status in a sea anemone model of cnidarian bleaching. *Sci Rep-uk*. 2017;7:16025.
81. Mansfield KM, Cleves PA, Vlack EV, Kriefall NG, Benson BE, Camacho DJ, et al. Varied effects of algal symbionts on transcription factor NF- κ B in a sea anemone and a coral: possible roles in symbiosis and thermotolerance. *bioRxiv*. 2019:640177. <https://doi.org/10.1101/640177>.
82. Tan ES, Izumi R, Takeuchi Y, Isomura N, Takemura A. Molecular approaches underlying the oogenic cycle of the scleractinian coral, *Acropora tenuis*. *Sci Rep-uk*. 2020;10:9914.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

