CORRECTION Open Access

Check for updates

Correction to: Seismicity distribution in the Tonankai and Nankai seismogenic zones and its spatiotemporal relationship with interplate coupling and slow earthquakes

Yojiro Yamamoto^{1*}, Shuichiro Yada¹, Keisuke Ariyoshi¹, Takane Hori¹ and Narumi Takahashi²

Correction to: Progress in Earth and Planetary Science (2022) 9:32

https://doi.org/10.1186/s40645-022-00493-4

Following publication of the original article (Yamamoto et al. 2022), the below minor errors were found in the article:

On page 12, the citation information in the caption of Fig. 10 should be Arnulf et al. (2022);

On page 12, the color in Figure 11 is not correctly reflected, and correct Fig. 11 is shown as below:

On page 13, the citation Takemura et al. 20192019 should be Takemura et al. 2019;

On page 18, the author name of Kimura KH should be Kimura T, Kimura H; the correct reference is: Hirose H, Matsuzawa T, Kimura T, Kimura H (2014) The Boso slow slip events in 2007.

The original paper has been updated.

Author details

¹Agency for Marine-Earth Science and Technology, Kanazawa-ku, Yokohama, Kanagawa, Japan. ²National Research Institute for Earth Science and Disaster Resilience, 3-1 Tennodai, Tsukuba, Ibaraki 305-0006, Japan.

Published online: 22 June 2022

The original article can be found online at https://doi.org/10.1186/s40645-022-00493-4.

*Correspondence: yamamotoy@jamstec.go.jp

Full list of author information is available at the end of the article



Reference

Yamamoto et al (2022) Seismicity distribution in the Tonankai and Nankai seismogenic zones and its spatiotemporal relationship with interplate coupling and slow earthquakes. Prog Earth Planet Sci 9:32. https://doi.org/10.1186/s40645-022-00493-4

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

¹ Agency for Marine-Earth Science and Technology, Kanazawa-ku, Yokohama, Kanagawa, Japan




Fig. 11 The location of the active areas for interplate seismicity and very low-frequency earthquakes (VLFEs). Relocated hypocenters within the 1 km from the plate interface are shown by pink circles. a Numbers from 1 to 14 reflect area code. VLFEs estimated by using the offshore seismic network are shown by green circles (Sugioka et al. 2012; Nakano et al. 2018a; Toh et al. 2020; Yamamoto et al. 2022). Location of the subducted seamount of Muroto and the Paleo-Zenith Ridge is shown by light-blue areas. b VLFEs estimated by using onshore seismic network are shown by yellow circles (Takemura et al. 2019). Other symbols are the same as in Fig. 9