

CASE REPORT

Open Access



A novel missense mutation in *PLEKHG5* gene causing an intermediate form of autosomal-recessive Charcot–Marie–Tooth disease in an Iraqi family

Mostafa Neissi^{1,2,3*} , Hadideh Mabudi⁴, Adnan Issa Al-Badran⁵, Javad Mohammadi-Asl^{3,6} and Raed Abdulelah Al-Badran⁷

Abstract

Background Charcot–Marie–Tooth disease comprises a large spectrum of clinically heterogeneous disorders. *PLEKHG5* variants have shown an intermediate form of autosomal-recessive Charcot–Marie–Tooth disease C and distal spinal muscular atrophy IV. The purpose of this case study is to report a causative genetic defect associated with intermediate form of autosomal-recessive Charcot–Marie–Tooth disease C in an Iraqi consanguineous family.

Case presentation Whole-exome and Sanger sequencing was used to identify probable gene defects in a 9-year-old male affected by CMTRIC. We found a new single mutation (c.1844C > A; p.T615N) in the *PLEKHG5* gene, located in exon 17 (NM_020631.6), causing a missense mutation that has been changed one amino acid. The mutation was homozygous in the patient and heterozygous in his parents.

Conclusion Our results expand the *PLEKHG5* pathogenic mutation spectrum related to intermediate form of autosomal-recessive Charcot–Marie–Tooth disease C which is vital for screening and genetic diagnosis of the disease.

Keywords *PLEKHG5* gene, Charcot–Marie–Tooth disease, Mutation

Background

Charcot–Marie–Tooth (CMT) disease, also known as hereditary motor and sensory neuropathy (HMSN), encompasses a genetically and clinically heterogeneous group of disorders characterized by weakness, muscular wasting, and sensory loss usually most severe distally. At least 1 in 2,500 people are affected by this inherited neuromuscular disorder [1].

There are now >100 genetic mutations identified as causative for CMT. Based on neurophysiology and nerve biopsy, CMT is classified into two subtypes, demyelinating forms (CMT1) and axonal forms (CMT2). An intermediate group also exists with nerve conduction velocities, which overlaps the two main groups, and is subdivided into recessive intermediate CMT (RI-CMT)

*Correspondence:

Mostafa Neissi
iammostafaneissi@gmail.com

¹ Department of Genetics, Khuzestan Science and Research Branch, Islamic Azad University, Ahvaz, Iran

² Department of Genetics, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran

³ Noor Gene Genetic Laboratory, Ahvaz, Iran

⁴ Department of Biotechnology, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran

⁵ Department of Biology, College of Science, University of Basrah, Basrah, Iraq

⁶ Cancer, Petroleum and Environmental Pollutants Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁷ Department of Genetics, Science and Research Branch, Islamic Azad University, Tehran, Iran

and dominant intermediate (DI-CMT) by its pattern of inheritance [2].

Pleckstrin homology domain-containing, family G member 5 (*PLEKHG5*) gene encodes a guanine exchange factor specifically expressed in motor neurons, where it regulates the autophagy of synaptic vesicles. *PLEKHG5* gene plays essential role in the activation and regulation of RhoA signaling pathways, thereby controlling neuronal cell differentiation and maintaining stable cell–cell contacts. Moreover, mutations in the *PLEKHG5* gene are causing different forms of motor neuron diseases such as distal spinal muscular atrophy IV (DSMA4, also known as lower motor neuron diseases [LMND]) and intermediate Charcot–Marie–Tooth disease type C (CMTRIC), both of which inherited in an autosomal-recessive pattern [3, 4]. Autosomal-recessive CMTRIC is a very rarely seen neurogenetic disorder [5]. Whole-exome sequencing (WES) technique is a particularly useful tool for understanding the molecular genetic causes of the CMT [6]. According to this evidence, we employed WES to identify the likely disease-causing genetic variants in a consanguineous Iraqi family affected by CMT. Here, we present the first consanguineous CMTRIC case due to a *PLEKHG5* substitution mutation not previously described.

Case presentation

In the present study, there is an Iraqi family with a 9-year-old male patient (Fig. 1), who was referred to the pediatric neurologists due to frequent outdoor falls and

difficulty in climbing up stairs. The proband was determined to have a clinical diagnosis consistent with CMT. His parents mentioned that he had normal motor and mental development compared to his peers during infancy and childhood. In addition, the patient had not a family history of neuromuscular diseases. Neurologic examination showed limb–girdle muscular weakness. His parents also noted that he started experiencing difficulty in getting up from squatting position. Physical examination revealed scapula winging and absent deep reflexes. Creatine kinase (CPK) levels were slightly elevated, and metabolic screening of urine, lactate, and pyruvate levels was all within normal limits. During follow-up, motor regression was noted with a positive Gowers’ sign with a longer duration. In addition, he could not stand up from a sitting position without support. Neurological examination revealed atrophy of bilateral distal muscles and muscle weakness, predominantly in the lower limbs. He had a 25-degree thoracolumbar scoliosis. Ophthalmic examination and hearing tests were in normal ranges, and also, his cardiac evaluation, including an echocardiogram and electrocardiogram, was within normal limits.

At his electrophysiological examination, motor nerve conduction studies (NCSs) and F-wave studies were performed for the right tibial, ulnar, and peroneal nerves, and sensory NCSs were performed for the right median, ulnar, and bilateral sural nerves using standard techniques. F-wave latencies were significantly prolonged in the median and tibial motor nerves. The ulnar and median motor and sensory nerve conduction velocities




Fig. 1 Pedigree of family. Patient 1 was a 9-year-old affected male in the presented study. The parents of the affected son (3,4) are first cousins

were slow. The ulnar and median motor and ulnar sensory nerve conduction velocities were within normal ranges, while the median sensory nerve conduction velocity was slow. Bilateral sural nerve sensory action potentials could be obtained, but with very low amplitudes. Fibrillation potentials and positive sharp waves were both significant active and chronic neurogenic findings for the tibialis anterior, biceps, iliopsoas, and first dorsal interosseous muscles with needle electromyography (EMG) examination. Myokymic discharges were found in the biceps muscle of the proband. The electrophysiological study suggested a predominantly axonal sensory and motor polyneuropathy and some demyelinating features. Finally, he was diagnosed as having CMT and we applied WES technique to identify the likely disease-causing mutations in the patient's DNA that may be inherited from the patient's ancestors.

We solely performed WES for the proband. For genomic DNA (gDNA) extraction from peripheral blood, a standard salting out method was used. The patient's DNA was captured using SureSelect Human All Exome Kit V6 (Agilent Technologies Inc., Santa Clara, CA, USA) and sequenced those regions using the Illumina HiSeq 4000 machine (San Diego, CA, USA) as stated by the manufacturer's protocols. The average read depth was more than 100×, and depth of 20× or greater was achieved in 98.0% of the targeted genomic sequence. We detected a novel missense substitution mutation in exon 17 of *PLEKHG5* gene (NM_020631.6: c.1844C>A; p.Thr615Asn), which indicates that this mutation is associated with CMTRIC. This mutation predicts an alteration in codon translation, and finally a change in one amino acid (threonine converted to asparagine).

Furthermore, the c.1844C>A; p.Thr615Asn mutation was classified as likely pathogenic according to the American College of Medical Genetics and Genomics (ACMG) guidelines. Bioinformatics software such as MutationTaster, PolyPhen-2, SIFT, and CADD predicted that this mutation will be deleterious and disease-causing.

To co-segregate and confirm the presence of the pathogenic *PLEKHG5* mutation, direct Sanger sequencing method for the patient and his parents was done. So, we have designed a specific set of primer pairs (forward primer: GGCAGCCATATTCCACAAGT and the reverse primer: GGCATTGTAAATGGTGTCC) to amplify the mutated site in the genome by PCR method. After amplification of *PLEKHG5* sequences, we sequenced the PCR products directly on the automated genetic analyzer (Applied Biosystems 3130xl; USA) and the results are represented in Fig. 2. The sequences were blasted in NCBI website (<http://www.ncbi.nlm.nih.gov/blast>) and compared with normal sequences. Interestingly, this finding has not been reported in the other patients.

Figure 2D shows the location of the novel variant of *PLEKHG5* protein. The mutant amino acids are located in the highly conserved PH domain of *PLEKHG5* gene, which includes a domain in 584–684 [3].

Discussion

In this study, WES method was applied to detect the causative gene defects related to CMT in an Iraqi family. The index patient was a homozygous carrier for *PLEKHG5* T615N mutation, and the parents of the patient were both heterozygous. Therefore, the NM_000249 (*PLEKHG5*): c.1844C>A mutation is surely responsible for CMTRIC phenotype in the proband. This is the first Iraqi case of a CMTRIC phenotype caused by *PLEKHG5* gene mutation. So far, eight different *PLEKHG5* mutations have been recognized based on the Human Gene Mutation Database (HGMD; <http://www.hgmd.cf.ac.uk/ac/index.php>), one of which presented childhood-onset LMND, whereas the other seven presented intermediate CMT, amyotrophic lateral sclerosis (ALS), and myopathy.

Hence, in this study, we investigated the disease-causing mutation in a 9-year-old affected male, who had limb-girdle muscle weakness, and this condition was reported in almost all cases [7]. Distal sensory loss was reported in almost 1/3 of the patients. All of the patients had distal muscle weakness in the lower limbs. In contrast, in the current study, the patient had proximal muscle weakness in the lower extremities. However, the presence of areflexia, distal muscle atrophy, spinal cord abnormalities, and foot deformity was consistent with other defined cases.

Electrophysiological studies of patients showed prolonged motor latencies and their median motor nerve conduction velocities were slowed, the amplitudes of the sensory nerve action potentials (SNAP) were absent or decreased, and their conduction velocities were sufficiently slowed in the affected nerves. Needle EMG showed muscle denervation in almost all of the patients [8, 9]. In the present case, intermediate polyneuropathy was detected moderately with predominantly axonal features evident with active and chronic denervation findings.

PLEKHG5 is predominantly expressed in the human peripheral nervous system, and the protein contains a Dbl homology (DH)–Pleckstrin homology (PH) motif, which is known as the minimal unit required for the nucleotide-exchange-promoting function of guanine nucleotide-exchange factors (GEFs) [2]. Recessive mutations in the PH domain and *PLEKHG5* gene were initially reported in a consanguineous family with early-onset diseases of the lower motor neuron (or DSMA4) and formally have been associated with intermediate CMT. In more studies, cases with either pure LMND, distal and




Fig. 2 Sanger sequencing confirming homozygous c.1844C>A mutation in the patient (A) and heterozygous mutation in his parents (B, C). D Diagram structure of *PLEKHG5* protein. Schematic of *PLEKHG5* protein shows the locations of the pathogenic variant in humans

proximal neuropathy with mild sensory involvement, or intermediate CMT have been documented [10]. Additionally, we reported the case of a 9-year-old Iraqi male who presented with CMTRIC with a novel homozygous mutation in *PLEKHG5* gene (c.1844C>A; p.Thr615Asn), and this detected mutation affects a conserved residue and supports a positive pathogenic role for this gene in causing the CMTRIC.

PLEKHG5's molecular functions and disease-associated mechanisms are limited: Previously, only one homozygous substitution mutation, c.1940 T>C; p.Phe647Ser, was reported as a pathogenic mutation in *PLEKHG5* gene associated with autosomal-recessive LMND. The site locates within the PH domain, and the mutation was suggested to affect the nuclear factor kappa B (NF-κB) activation. According to immunohistochemical results from the patient and immunocytochemical results obtained from mutants that have been cloned,

mutant proteins have lower expression levels than wild-type proteins. As a result, these data suggest that the patient with peripheral neuropathy is due to low levels of expression and NF-κB activation [2, 11]. Also, we proposed that our identified Thr615Asn variant encodes an impaired *PLEKHG5* protein that probably causes a defect in the activation of the NF-κB signaling pathway resulting in the CMTRIC.

Some studies revealed that mutations in *PLEKHG5* including c.1600-2A>G, c.1835_1860del, c.2308del and c.104del are associated with intermediate CMT disease [5, 10]. In this regard, Beijer D. et al. identified a homozygous single *PLEKHG5* mutation (p.Arg97Gln), predicting an alteration in codon translation (arginine converted to glutamine), which affects the *PLEKHG5* protein stability, resulting in CMT clinical manifestations [12]. Moreover, Kim HJ. et al., in their publication, reported a case of autosomal-recessive intermediate CMT disease

with novel compound heterozygous (p.Thr663Met and p.Gly820Arg) variants in the *PLEKHG5* gene [2]. Our report adds a novel variant to *PLEKHG5* mutation spectrum and indicates that a missense c.1844C>A; p.Thr615Asn mutation probably causes *PLEKHG5* dysfunction leading to the CMTRIC.

Finally, this detected mutation that inherited from patient's ancestors is proposed as the cause of CMTRIC in the patient and the following evidence proves that this mutation can lead to the disease: 1—WES identified only this mutation as the cause of CMTRIC. 2—As can be seen in Fig. 2A, direct Sanger sequencing proved the mutation in the proband, and based on recognized heterozygote mutations in his parents, the pattern of inheritance must be an autosomal-recessive for the *PLEKHG5* gene. 3—Bioinformatics software such as MutationTaster, PolyPhen-2, SIFT, and CADD predicted that c.1844C>A mutation is the pathogenic mutation for CMTRIC. 4—Also, a substitution mutation in exon 17 of the *PLEKHG5* gene (c.1844C>A; p.Thr615Asn) in the PH domain of the protein (Fig. 2D), which is predicted to create new codon substitutions, can create a major problem in the *PLEKHG5* protein. Based on our results, WES is an efficient approach of analyzing a patient's DNA to discover the genetic cause of CMTRIC.

Conclusion

We have successfully applied WES in an Iraqi family for mutation screening within CMT-related genes and identified a novel substitution mutation in the *PLEKHG5* gene, thereby confirming previous reports that *PLEKHG5* gene mutations are associated with CMTRIC. This study may be helpful in providing appropriate genetic counseling to the affected families and provides a new way to clarify the molecular mechanisms of the *PLEKHG5* gene in CMTRIC.

Abbreviations

ACMG	American College of Medical Genetics and Genomics
CMTRIC	Intermediate Charcot-Marie-Tooth disease type C
CPK	Creatine kinase
DH	Dbl homology
DSMA4	Distal spinal muscular atrophy IV
EMG	Electromyography
gDNA	Genomic DNA
GEFs	Guanine nucleotide-exchange factors
HMSN	Hereditary motor and sensory neuropathy
LMNDs	Lower motor neuron diseases
NF-κB	Nuclear factor kappa B
NCSS	Nerve conduction studies
PH	Pleckstrin homology
SNAP	Sensory nerve action potentials
WES	Whole-exome sequencing

Acknowledgements

The authors would like to thank the family members for their participation in this study.

Author contributions

MN, HM, AIA, JMA, and RAA analyzed and interpreted the data. MN wrote the manuscript. HM edited the manuscript. All authors read and approved the final manuscript.

Funding

None.

Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or compare ethical strand.

Consent for publication

Written informed consent was obtained from the family for this publication.

Competing interest

The authors declare that they have no conflict of interest.

Received: 1 August 2022 Accepted: 1 March 2023

Published online: 20 March 2023

References

- Reilly MM, Murphy SM, Laurá M (2011) Charcot-Marie-Tooth disease. *J Peripher Nerv Syst* 16(1):1–14. <https://doi.org/10.1111/j.1529-8027.2011.00324.x>
- Kim HJ, Hong YB, Park JM, Choi YR, Kim YJ, Yoon BR, Koo H, Yoo JH, Kim SB, Park M, Chung KW, Choi BO (2013) Mutations in the *PLEKHG5* gene is relevant with autosomal recessive intermediate Charcot-Marie-Tooth disease. *Orphanet J Rare Dis* 12(8):104. <https://doi.org/10.1186/1750-1172-8-104>
- Gonzalez-Quereda L, Pagola I, Fuentes Prior P, Bernal S, Rodriguez MJ, Torné L, Salgado Garrido J, Gallano P, Jericó I (2021) Novel *PLEKHG5* mutations in a patient with childhood-onset lower motor neuron disease. *Ann Clin Transl Neurol* 8(1):294–299. <https://doi.org/10.1002/acn3.51265>
- Miao Y, Yu M, Meng L, Zhang W, Lv H, Wang Z, Yuan Y (2021) *PLEKHG5*-related autosomal recessive lower motor neuron disease with dysmyelination in peripheral nerves. *Clin Neuropathol* 40(6):328–332. <https://doi.org/10.5414/NP301377>
- Yayici Köken Ö, Öztoprak Ü, Topçu V, Çavdarlı B, Temuçin ÇM, Aydingöz Ü et al (2021) Expanding the genotype-phenotype spectrum of autosomal recessive charcot-marie-tooth disease: a novel *plekhg5* gene mutation. *Neurol Asia* 26(3):607–612
- Choi BO, Koo SK, Park MH, Rhee H, Yang SJ, Choi KG, Jung SC, Kim HS, Hyun YS, Nakhro K, Lee HJ, Woo HM, Chung KW (2012) Exome sequencing is an efficient tool for genetic screening of Charcot-Marie-Tooth disease. *Hum Mutat* 33(11):1610–1615. <https://doi.org/10.1002/humu.22143>
- Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, Voelkerding K, Rehml HL, ACMG Laboratory Quality Assurance Committee (2015) Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 17(5):405–424. <https://doi.org/10.1038/gim.2015.30>
- Chen Z, Maroofian R, Başak AN, Shingavi L, Karakaya M, Efthymiou S, Gustavsson EK, Meier L, Polavarapu K, Vengalil S, Preethish-Kumar V, Nandeesh BN, Gökçe Güneş N, Akan O, Candan F, Schrank B, Zuchner S, Murphy D, Kapoor M, Ryten M, Wirth B, Reilly MM, Nalini A, Houlden H, Sarraf P (2021) Novel variants broaden the phenotypic spectrum of

- PLEKHG5-associated neuropathies. *Eur J Neurol* 28(4):1344–1355. <https://doi.org/10.1111/ene.14649>
9. Saporta AS, Sottile SL, Miller LJ, Feely SM, Siskind CE, Shy ME (2011) Charcot-Marie-Tooth disease subtypes and genetic testing strategies. *Ann Neurol* 69(1):22–33. <https://doi.org/10.1002/ana.22166>
 10. Villar-Quiles RN, Le VT, Leonard-Louis S, Trang NT, Huong NT, Laddada L, Francou B, Maisonobe T, Azzedine H, Stojkovic T (2021) Leukoencephalopathy and conduction blocks in PLEKHG5-associated intermediate CMT disease. *Neuromuscul Disord* 31(8):756–764. <https://doi.org/10.1016/j.nmd.2021.06.004>
 11. Maystadt I, Rezsöhazy R, Barkats M, Duque S, Vannuffel P, Remacle S, Lambert B, Najimi M, Sokal E, Munnich A, Viollet L, Verellen-Dumoulin C (2007) The nuclear factor kappaB-activator gene PLEKHG5 is mutated in a form of autosomal recessive lower motor neuron disease with childhood onset. *Am J Hum Genet* 81(1):67–76. <https://doi.org/10.1086/518900>
 12. Beijer D, Polavarapu K, Preethish-Kumar V, Bardhan M, Dohrn MF, Rebelo A, Züchner S, Nalini A (2022) Homozygous N-terminal missense variant in PLEKHG5 associated with intermediate CMT: a case report. *J Neuromuscul Dis* 9(2):347–351. <https://doi.org/10.3233/JND-210716>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ [springeropen.com](https://www.springeropen.com)
