Resecting Lymph Nodes in Colon Cancer: More than a Staging Operation?

Steven L. Chen, MD, MBA, and Anton J. Bilchik, MD, PhD, FACS

Department of Surgical Oncology, John Wayne Cancer Institute at Saint John's Health Center,
2200 Santa Monica Blvd, Santa Monica, 90404 California, USA

Erratum to: 10.1245/s10434-007-9430-8
Annals of Surgical Oncology 14(8): 2175–2176

Reference 1 appeared incorrectly in the online and print versions. The updated reference is as follows:

REFERENCE

1. Lee, H-Y, Choi, H-J, Park, K-J, et al. Prognostic significance of metastatic lymph node ratio in node-positive colon carcinoma. *Ann Surg Oncol* 14: 1712–1717; DOI: [10.1245/s10434-006-9322-3](https://doi.org/10.1245/s10434-006-9322-3) [Online: January 26, 2007].

Published online October 25, 2007.

The online version of the original article can be found under doi:
[10.1245/s10434-007-9430-8](https://doi.org/10.1245/s10434-007-9430-8)

Address correspondence and reprint requests to: Anton J. Bilchik, MD, PhD, FACS; E-mail: bilchika@jwci.org

Published by Springer Science+Business Media, LLC © 2007 The Society of Surgical Oncology, Inc.