ERRATUM

Erratum to: Clinicopathologic Features of Incident and Subsequent Tumors in Patients with Multiple Primary Cutaneous Melanomas

Rajmohan Murali, MBBS, MD, FRCPA^{1,2,3,14}, Chris Goumas, MPH⁴, Anne Krickler, PhD^{2,4}, Lynn From, MD, FRCP⁵, Klaus J. Busam, MD⁶, Colin B. Begg, PhD⁷, Terence Dwyer, MD, MPH⁸, Stephen B. Gruber, MD, PhD, MPH⁹, Peter A. Kanetsky, PhD, MPH¹⁰, Irene Orlow, PhD⁷, Stefano Rosso, PhD¹¹, Nancy E. Thomas, MD, PhD¹², Marianne Berwick, PhD, MPH¹³, Richard A. Scolyer, MBBS, MD, FRCPA, FRCP^{1,2,3}, Bruce K. Armstrong, AM, DPhil, FRACP, FAFPHM, FAA^{2,4}, For the GEM Study Group

¹Tissue Pathology and Diagnostic Oncology, Royal Prince Alfred Hospital, Sydney, Australia; ²Melanoma Institute Australia, Sydney, Australia; ³Discipline of Pathology, The University of Sydney, Sydney, Australia; ⁴School of Public Health, Sydney Medical School, The University of Sydney, Sydney, Australia; ⁵Women's College Hospital, Toronto, ON, Canada; ⁶Department of Pathology, Memorial Sloan-Kettering Cancer Center, New York; ⁷Department of Epidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, New York, NY; ⁸Murdoch Children's Research Institute, Royal Children's Hospital, Melbourne, Australia; ⁹Division of Molecular Medicine & Genetics, University of Michigan, Ann Arbor, MI; ¹⁰University of Pennsylvania School of Medicine, Philadelphia, PA; ¹¹CPO-Registro Tumori Piemonte, Turin, Italy; ¹²Department of Dermatology, Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC; ¹³Division of Epidemiology and Biostatistics, University of New Mexico, Albuquerque, NM; ¹⁴Department of Pathology and Human Oncology and Pathogenesis Program, Memorial Sloan-Kettering Cancer Center, New York, NY

ERRATUM TO: ANN SURG ONCOL (2012) 19:1024–1033 DOI 10.1245/S10434-011-2058-8

Table 3 of the article entitled “Clinicopathologic Features of Incident and Subsequent Tumors in Patients with Multiple Primary Cutaneous Melanomas” by Murali et al (*Ann Surg Oncol*: 19[3]: 1024–1033, 2012 DOI: [10.1245/s10434-011-2058-8](https://doi.org/10.1245/s10434-011-2058-8)) incorrectly re-displayed Table 1. The correct Table 3 is:

The online version of the original article can be found under doi: [10.1245/s10434-011-2058-8](https://doi.org/10.1245/s10434-011-2058-8).

© Society of Surgical Oncology 2012

Published Online: 11 September 2012

R. Murali, MBBS, MD, FRCPA
e-mail: MuraliR@mskcc.org

TABLE 3 Results of a stepwise multiple regression analysis to identify variables that independently distinguished subsequent from first melanomas in patients with multiple primary melanomas

Variables	Tumor		Prevalence ratios ^b for MPM2 (95% CI)	<i>P</i> value
	First melanoma ^a	Subsequent melanoma ^a		
Co-existing nevus				<0.001
Not identified	226	227	1.0	
Dysplastic	54	83	1.29 (1.10–1.53)	
Common acquired	54	22	0.61 (0.43–0.87)	
Congenital and others	2	4	1.78 (0.97–3.29)	
Vertical growth				0.01
Absent	157	191	1.0	
Present	179	145	0.82 (0.70–0.95)	
Site of melanoma				0.04
Head and neck	56	78	1.0	
Trunk	156	143	0.81 (0.68–0.96)	
Arms	61	48	0.78 (0.60–1.02)	
Legs	63	67	0.98 (0.81–1.19)	

^a First and subsequent melanomas are primary invasive cutaneous melanomas in patients with multiple primary melanomas

^b Prevalence ratio for subsequent melanoma compared with first melanoma in patients with multiple primary melanomas, clustered on patient and adjusted for age at diagnosis of lesion, sex, GEM center, and all other variables in table. Highest *P* value for entry 0.2; 336 cases with complete data for all variables included in the analysis