ERRATUM

Erratum to: Clinicopathology and Genetic Profile of Synchronous Multiple Small Adenocarcinomas: Implication for Surgical Treatment of an Uncommon Lung Malignancy

Mong-Wei Lin, MD^{1,5}, Chen-Tu Wu, MD^{2,5}, Shuenn-Wen Kuo, MD³, Yih-Leong Chang, MD^{2,5}, and Pan-Chyr Yang, MD, PhD^{4,5}

¹Department of Surgery, National Taiwan University Hospital Hsin-Chu Branch, Hsinchu, Taiwan; ²Department of Pathology, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan; ³Department of Surgery, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan; ⁴Department of Internal Medicine, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan; ⁵Graduate Institute of Pathology, National Taiwan University College of Medicine, Taipei, Taiwan

ERRATUM TO: ANN SURG ONCOL DOI 10.1245/S10434-014-3642-5

Errors occurred in Table 3, Table 4, and Results, which are corrected here. Specifically, the parameters of age (years) were reversed in Tables 3 and 4. The data on patient number and relapse number of smoking status was reversed in Table 4. In the section “*Correlation between Clinicopathologic Features and Disease-Free Survival*” of “Results”, one risk factor for relapse identified by univariate analysis was age > 65 years ($p = 0.020$).

The correct versions of these two tables are

TABLE 3 Univariate survival analysis of clinicopathologic features in patients with synchronous multiple small lung adenocarcinomas

Parameters	No. of patients	No. of relapses (%) / 5-year DFS (%)	p Value
Total no. of patients	64	8 (12.5)/86.1	
Age (years)			
>65	25	6 (24.0)/78.4	0.020
≤65	39	2 (5.1)/93.0	
Sex			
Male	18	3 (16.7)/82.5	0.798
Female	46	5 (10.9)/89.3	
Smoking status			
Positive	6	3 (50.0)/50.0	0.011
Negative	58	5 (8.6)/91.3	
CEA			
Normal	61	6 (9.8)/88.3	0.056
Abnormal	3	2 (66.7)/66.7	
Tumor number			
≥3	11	1 (9.1)/90.0	0.708
=2	53	7 (13.2)/87.3	
Largest tumor size (cm)			
>1 cm	40	7 (17.5)/82.7	0.152
≤1 cm	24	1 (4.2)/95.5	
Tumor site			
One lobe	23	4 (17.4)/85.3	0.828
>1 lobe, unilateral	20	2 (10.0)/86.8	
>1 lobe, bilateral	21	2 (9.5)/89.3	
CT characteristics			
Pure GGO	14	0 (0.0)/100.0	
Mixed type	42	4 (9.5)/88.7	

The online version of the original article can be found under doi:10.1245/s10434-014-3642-5.

© Society of Surgical Oncology 2014

Published Online: 1 July 2014

Y.-L. Chang, MD
e-mail: ntuhyllc@gmail.com

TABLE 3 continued

Parameters	No. of patients	No. of relapses (%) / 5-year DFS (%)	<i>p</i> Value
Solid	8	4 (50.0)/62.5	0.016
Vascular invasion			
Positive	2	1 (50.0)/0.0	0.092
Negative	62	7 (11.3)/89.3	
Pleural invasion			
Positive	5	1 (20.0)/50.0	0.358
Negative	59	7 (11.9)/89.0	
Lymph node metastasis ^a			
Positive	4	3 (75.0)/37.5	0.003
Negative	54	4 (7.4)/91.8	
Surgery			
Sublobar resection	36	2 (5.6)/96.3	

TABLE 4 Multivariate survival analysis of clinicopathologic features in patients with synchronous multiple small lung adenocarcinomas

Parameters	No. of patients	No. of relapses (%)	Hazard ratio	95% CI	<i>p</i> Value
Age (years)					
≤65	39	2 (5.1)	0.092	0.006–1.325	0.080
>65	25	6 (24.0)	1.000		
Smoking status					
Negative	58	5 (8.6)	0.055	0.005–0.601	0.017
Positive	6	3 (50.0)	1.000		
Lymph node metastasis ^a					
Negative	54	4 (7.4)	0.058	0.005–0.665	0.022
Positive	4	3 (75.0)	1.000		
Clonality assessment					
Wild type	20	2 (10.0)	1.022	0.033–31.844	0.990
SC	14	4 (28.6)	15.216	0.845–273.959	
DC	30	2 (6.7)			
CT characteristics					
Pure GGO	14	0 (0.0)	0.000	0.000	0.984
Mixed type	42	4 (9.5)	0.458	0.029–7.208	
Solid	8	4 (50.0)			

CI confidence interval, *CT* computed tomography, *DC* Different clonality, *GGO* ground glass opacity, *SC* Same clonality

^a Six patients did not undergo lymph node dissection