



CORRECTION

Correction to: Midgut Neuroendocrine Tumors with Liver-only Metastases: Benefit of Primary Tumor Resection

Alexandra Gangi, MD¹, Nicholas Manguso, MD², Jun Gong, MD³, Jessica S. Crystal, MD¹, Shirley C. Paski, MD, MSc, FRCPC⁴, Andrew E. Hendifar, MD³, and Richard Tuli, MD, PhD^{5,6}

¹Division of Surgical Oncology, Department of Surgery, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA; ²Department of Surgery, Cedars-Sinai Medical Center, Los Angeles, CA; ³Department of Gastrointestinal Malignancies, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA; ⁴Department of Gastroenterology, Cedars-Sinai Medical Center, Los Angeles, CA; ⁵Department of Radiation Oncology, Cedars-Sinai Medical Center, Los Angeles, CA; ⁶Department of Radiation Oncology, Memorial Sloan Kettering Cancer Center, New York, NY

CORRECTION TO:

ANN SURG ONCOL (2020) 27:4525–4532

[HTTPS://DOI.ORG/10.1245/S10434-020-08510-W](https://doi.org/10.1245/S10434-020-08510-W)

In the original online version of the article, the percentage for overall survival rate for resection is switched in results section of the abstract. The correct sentence is as follows:

The 5-year OS rate was 66% for the patients who underwent resection and 49% for those who did not ($p < 0.001$).

The online version of the article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1245/s10434-020-08510-w>.

© Society of Surgical Oncology 2021

Published Online: 22 March 2021

A. Gangi, MD

e-mail: Alexandra.gangi@cshs.org