ASO VISUAL ABSTRACT

ASO Visual Abstract: Surgical Merits of Open, Laparoscopic, and Robotic Gastrectomy Techniques with D2 Lymphadenectomy in Obese Patients with Gastric Cancer

Seohee Choi, MD¹, Jeong Ho Song, MD^{2,3}, Sejin Lee, MD^{2,3}, Minah Cho, MD^{2,3,4}, Yoo Min Kim, MD^{2,3,4}, Woo Jin Hyung, MD^{2,3,4}, and Hyoung-Il Kim, MD^{2,3,4}

¹Department of Surgery, Eulji General Hospital, Eulji University College of Medicine, Seoul, South Korea; ²Department of Surgery, Yonsei University College of Medicine, Seoul, Republic of Korea; ³Gastric Cancer Center, Yonsei Cancer Center, Yonsei University Health System, Seoul, Republic of Korea; ⁴Robot and MIS Center, Severance Hospital, Yonsei University Health System, Seoul, Republic of Korea

The robotic gastrectomy performed in obese patients with gastric cancer requiring D2 nodal dissection showed robust short-term surgical outcomes (<https://doi.org/10.1245/s10434-021-09952-6>). In long-term outcomes, robotic

group significantly outperformed laparoscopic group in overall survival, and open groups in overall survival and relapse-free survival.