ASO VISUAL ABSTRACT

ASO Visual Abstract: Lymphovascular Invasion—Traditional but Vital and Sensible Prognostic Factor in Early Gastric Cancer

Seohee Choi, MD^{1,2}, Jeong Ho Song, MD^{1,2}, Sejin Lee, MD^{1,2}, Minah Cho, MD^{1,2}, Yoo Min Kim, MD, PhD^{1,2}, Hyoung-Il Kim, MD, PhD^{1,2}, and Woo Jin Hyung, MD, PhD^{1,2}

¹Department of Surgery, Yonsei University College of Medicine, Seoul, Republic of Korea; ²Gastric Cancer Center, Yonsei Cancer Center, Yonsei University Health System, Seoul, Republic of Korea

Lymphovascular invasion (LVI) in gastric cancer has long been suggested as a poor prognostic indicator. Our study sought to evaluate the prognostic effect of LVI in patients with early gastric cancer (<https://doi.org/10.1245/s10434-021-10224-6>) and determined that early gastric cancer with LVI is indeed associated with poor prognosis and frequent recurrence.

DISCLOSURES Woo Jin Hyung reports receiving research grants from Medtronic and GC Pharma. He is also the chief executive officer of Hutomand and holds stocks in the company. He provided consultancy services to Ethicon and Verb Surgical outside the submitted work, and received lecture fees from Stryker. Seohee Choi, Jeong Ho Song,