ASO VISUAL ABSTRACT

ASO Visual Abstract: Temporizing Wound VAC Dressing Until Final Negative Margins are Achieved Reduces Myxofibrosarcoma Local Recurrence

Mitchell S. Fourman, MD, MPhil¹, Duncan C. Ramsey, MD, MPH¹, Justin Kleiner, MD², Anser Daud, BS³, Erik T. Newman, MD¹, Joseph H. Schwab, MD, MS¹, Yen-Lin Chen, MD⁴, Thomas F. DeLaney, MD⁴, John T. Mullen, MD⁵, Kevin A. Raskin, MD¹, and Santiago Lozano-Calderon, MD, PhD¹

¹Department of Orthopaedic Surgery, Massachusetts General Hospital, Boston, MA ; ²Department of Orthopaedic Surgery, Boston Medical Center, Boston, MA; ³Faculty of Medicine, University of Toronto, Toronto, ON, Canada; ⁴Department of Radiation Oncology, Massachusetts General Hospital Cancer Center, Boston, MA; ⁵Department of Surgery, Massachusetts General Hospital, Boston, MA

Suprafascial myxofibrosarcoma is microinvasive, and negative margins often are unclear using intraoperative frozen sectioning alone. Temporizing the wound bed with a VAC pending final margin analysis significantly reduces the local recurrence rate of suprafascial myxofibrosarcoma from 28.6% to 5.6% (<https://doi.org/10.1245/s10434-021-10242-4>).

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.