ASO VISUAL ABSTRACT

ASO Visual Abstract: Stage Migration and Survival Trends in Laryngeal Cancer

Michael M. Li, MD¹, Songzhu Zhao, MS², Antoine Eskander, MD, ScM³, Chandler Rygalski, BS⁴, Guy Brock, PhD², Anuraag S. Parikh, MD¹, Catherine T. Haring, MD¹, Brian Swendseid, MD¹, Kevin Y. Zhan, MD¹, Carol R. Bradford, MD¹, Theodoros N. Teknos, MD⁵, Ricardo L. Carrau, MD, MBA¹, Kyle K. VanKoeveering, MD¹, Nolan B. Seim, MD¹, Matthew O. Old, MD¹, James W. Rocco, MD, PhD¹, Sidharth V. Puram, MD, PhD^{6,7}, and Stephen Y. Kang, MD¹

¹Department of Otolaryngology–Head and Neck Surgery, The Ohio State University Wexner Medical Center, Columbus, OH; ²Department of Biomedical Informatics and Center for Biostatistics, The Ohio State University, Columbus, OH; ³Department of Otolaryngology–Head and Neck Surgery, Sunnybrook Health Sciences Center, Toronto, ON, Canada; ⁴College of Medicine, Ohio State University, Columbus, OH; ⁵University Hospital Seidman Cancer Center, Cleveland, OH; ⁶Department of Otolaryngology–Head and Neck Surgery, Washington University School of Medicine, St. Louis; ⁷Department of Genetics, Washington University School of Medicine, St. Louis

Although significant advances in treatment protocols were developed in the 2000s, laryngeal cancer survival did not improve from 2004 to 2016. Concurrently, stage migration

toward greater proportions of stage IV disease has occurred. Early diagnosis and treatment are critical to optimize outcomes (<https://doi.org/10.1245/s10434-021-10318-1>).