ASO VISUAL ABSTRACT

ASO Visual Abstract: Gastric Neuroendocrine Tumors: Reappraisal of Type in Predicting Outcome

Andrew Hanna, MD¹ , Caroline Kim-Kiselak, MA², Rebecca Tang, MD³, David C. Metz, MD⁴, Zhaohai Yang, MD, PhD⁵, Ronald DeMatteo, MD¹, Douglas L. Fraker, MD¹, and Robert E. Roses, MD¹

¹Department of Surgery, University of Pennsylvania, Philadelphia, PA; ²University of Pennsylvania School of Medicine, Philadelphia, PA; ³Department of Surgery, Massachusetts General Hospital, Boston, MA; ⁴Department of Medicine, University of Pennsylvania, Philadelphia, PA; ⁵Department of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA

Type 3 gastric neuroendocrine tumors (GNETs) compared with type 1 GNETs are associated with worse outcomes and often are treated surgically rather than endoscopically. This study (<https://doi.org/10.1245/s1043>

[4-021-10293-7](https://doi.org/10.1245/s10434-021-10293-7)) identified a subset of type 3 GNETs that may be treated more conservatively, similar to the treatment for type 1 GNETs.

Abstract orally presented at the Society of Surgical Oncology 18 March 2021.

© Society of Surgical Oncology 2021

First Received: 4 June 2021

Accepted: 14 June 2021;

Published Online: 15 July 2021

A. Hanna, MD

e-mail: andrewnhanna4@gmail.com

DISCLOSURE David Metz is a consultant for Crinetics and Lexicon. The other authors have no conflicts of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.