ASO VISUAL ABSTRACT

ASO Visual Abstract: Longitudinal Assessment of Quality of Life Following Molecular Testing for Indeterminate Thyroid Nodules

Max A. Schumm, MD¹, Dalena T. Nguyen, MPH¹, Jiyoung Kim, MSc², Chi-Hong Tseng, PhD³, Amy Y. Chow, MD⁴, Na Shen, MD⁴, and Masha J. Livhits, MD¹

¹Section of Endocrine Surgery, Department of Surgery, University of California Los Angeles David Geffen School of Medicine, Los Angeles, CA; ²Department of Biostatistics, UCLA Fielding School of Public Health, Los Angeles, CA; ³Division of General Internal Medicine and Health Services Research, Department of Medicine, University of California Los Angeles David Geffen School of Medicine, Los Angeles, CA; ⁴Division of Endocrinology, Diabetes, and Metabolism, Department of Medicine, University of California, Los Angeles David Geffen School of Medicine, Los Angeles, CA

Patients with indeterminate thyroid nodules with benign molecular testing reported similar quality of life (QOL) through 18 months of surveillance, suggesting reassurance that the risk of malignancy is low

(<https://doi.org/10.1245/s10434-021-10375-6>). Definitive surgery resulted in improvement of QOL in patients with suspicious test results.

DISCLOSURE The authors have no related conflicts of interest to declare.

© Society of Surgical Oncology 2021

Published Online: 22 July 2021

M. A. Schumm, MD
 e-mail: mschumm@mednet.ucla.edu

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.