ASO VISUAL ABSTRACT

ASO Visual Abstract: Margin Width and Local Recurrence in Patients Undergoing Breast Conservation after Neoadjuvant Chemotherapy

Mary Mrdutt, MD¹, Alexandra Heerdt, MD¹, Varadan Sevilimedu, MBBS, DrPH^{1,2}, Anita Mamtani, MD¹, Andrea Barrio, MD¹, and Monica Morrow, MD¹

¹Breast Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY; ²Biostatistics Service, Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY

In patients receiving breast conservation therapy after neoadjuvant chemotherapy, an adequate negative margin has not been defined. We found no difference in isolated local recurrence rates between greater and less

than 2 mm margins after neoadjuvant chemotherapy and breast conservation (<https://doi.org/10.1245/s10434-021-10533-w>).

This work was presented as an oral presentation at the SSO 2021.

© Society of Surgical Oncology 2021

Published Online: 7 August 2021

M. Morrow, MD

e-mail: morrowm@mskcc.org

DISCLOSURE Dr. Monica Morrow has received speaking honoraria from Exact Sciences and Roche. The remaining authors have no conflicts of interest to disclose.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.