




ASO VISUAL ABSTRACT

## ASO Visual Abstract: A National Survey of Breast Surgeons and Radiation Oncologists on Contemporary Axillary Management in Mastectomy Patients

Chandler S. Cortina, MD<sup>1</sup> , Carmen Bergom, MD, PhD<sup>2,3</sup>, Morgan Ashley Craft, BS<sup>1</sup>, British Fields, (MLS)ASCP<sup>1</sup>, Ruta Brazauskas, PhD<sup>4</sup>, Adam Currey, MD<sup>3</sup>, and Amanda L. Kong, MD, MS<sup>1</sup>

<sup>1</sup>Section of Breast Surgery, Division of Surgical Oncology, Department of Surgery, Medical College of Wisconsin, Milwaukee, WI; <sup>2</sup>Department of Radiation Oncology and Siteman Cancer Center, Washington University School of Medicine, St Louis, MO; <sup>3</sup>Department of Radiation Oncology, Medical College of Wisconsin, Milwaukee, WI; <sup>4</sup>Division of Biostatistics, Institute of Health and Equity, Medical College of Wisconsin, Milwaukee, WI

A national case-based survey of breast surgeons and radiation oncologists found significant heterogeneity in the treatment of breast cancer patients who were clinically node-negative, underwent upfront mastectomy, and were

found to have pathological nodal involvement on sentinel lymph node biopsy (<https://doi.org/10.1245/s10434-021-10441-z>).

