ASO VISUAL ABSTRACT

ASO Visual Abstract: Objective Assessment of Postoperative Morbidity Following Breast Cancer Treatments with Wearable Activity Monitors: The “BRACELET” Study

Nur Amalina Che Bakri, MBChB, BMedSc, MPhil^{1,2} , Richard M. Kwasnicki, PhD, MRCS^{1,2}, Kieran Dhillon, BSc¹, Naairah Khan, BSc¹, Omar Ghandour, BSc¹, Alexander Cairns, MBBS², Ara Darzi, FRCS, FRS^{1,2}, and Daniel R. Leff, PhD, MS (Hons), FRCS^{1,2}

¹Department of Surgery and Cancer, Imperial College London, London, UK; ²Academic Surgical Unit, St. Mary's Hospital, Imperial College Healthcare NHS Trust, London, UK

Upper limb functionality after breast and axillary surgery lacks quantitative morbidity measurement. Wearable activity monitors (WAMs) provide objective insights into upper limb function which may enable characterization of

recovery of breast oncoplastic procedures and aid in the development of personalized rehabilitation (<https://doi.org/10.1245/s10434-021-10458-4>).

FUNDING National Institute for Health Research (NIHR) Imperial Biomedical Research Centre (BRC). Award number: WSSS_P69945

DISCLOSURES The authors declare that they have no conflict of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.