



ASO Visual Abstract: Optimal Lymph Node Staging System in Evaluating Prognosis of Gallbladder Carcinoma: A Multi-Institutional Study

Chen Chen, MD¹, Zhang Rui, MD¹, Wu Yuhuan, MD², Zhang Yongjie, MD³, Qiu Yinghe, MD³, Yang Ning, MD³, Song Tianqiang, MD⁴, Lou Jianying, MD⁵, Li Jiangtao, MD⁵, Mao Xianhai, MD⁶, Li Shengping, MD⁷, Si Shubin, PhD², Cai Zhiqiang, PhD², Tang Zhaohui, MD⁸, and Geng Zhimin, MD¹

¹Department of Hepatobiliary Surgery, First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China; ²Department of Industrial Engineering, School of Mechanical Engineering, Northwestern Polytechnical University, Xi'an, China; ³Department of Biliary Surgery, Eastern Hepatobiliary Surgery Hospital, Affiliated to Naval Medical University, Shanghai, China; ⁴Department of Hepatobiliary Oncology, Tianjin Medical University Cancer Hospital, Tianjin, China; ⁵Department of Hepatobiliary and Pancreatic Surgery, The Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou, China; ⁶Department of Hepatobiliary Surgery, Hunan Provincial People's Hospital, Changsha, China; ⁷Department of Hepatobiliary Oncology, Sun Yat-sen University Cancer Center, Guangzhou, China; ⁸Department of General Surgery, Xinhua Hospital Affiliated to Shanghai Jiaotong University School of Medicine, Shanghai, China

Lymph node (LN) involvement is a critical prognostic factor in patients with gallbladder carcinoma (GBC). Patients from six Chinese tertiary hospitals who underwent curative-intent resection for GBC between 2008 and 2013 were analyzed retrospectively. A total of 226 patients were involved in this cohort (<https://doi.org/10.1245/s10434-021-10749-w>). Based on the Tree-augmented Naïve Bayesian model and composite importance measures, the most important factor affecting prognosis in the different LN staging systems was number of metastatic LNs (NMLN). NMLN was the optimal LN staging system in evaluating the prognosis of GBC.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1245/s10434-021-10770-z>.

FUNDING Shaanxi Key Research and Development Program (No. 2017ZDXM-SF-055, No. 2017KW-060); Shaanxi Natural Science Basic Research Program General Project (No. 2020SF-070); The Multicenter Clinical Research Project of Shanghai Jiaotong University School of Medicine (No. DLY201807).

DISCLOSURE Chen Chen, Zhang Rui, Wu Yuhuan, Zhang Yongjie, Qiu Yinghe, Yang Ning, Song Tianqiang, Lou Jianying, Li Jiangtao, Mao Xianhai, Li Shengping, Si Shubin, Cai Zhiqiang, Tang Zhaohui, and Geng Zhimin have no conflicts of interest to disclose.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Chen Chen and Zhang Rui contributed equally to this work.

© Society of Surgical Oncology 2021

Published Online: 14 September 2021

T. Zhaohui
e-mail: tangzhaohui@yahoo.com

G. Zhimin
e-mail: gengzhimin@mail.xjtu.edu.cn