ASO VISUAL ABSTRACT

ASO Visual Abstract: Optimizing Axillary Management in Clinical T1–2N0 Mastectomy Patients with Positive Sentinel Lymph Nodes

Olga Kantor, MD MS^{1,2}, Jessica Means, MD¹, Samantha Grossmith, BA¹, Tanujit Dey, PhD³, Jennifer R. Bellon, MD⁴, Elizabeth A. Mittendorf, MD PhD^{1,2}, and Tari A. King, MD^{1,2}

¹Division of Breast Surgery, Department of Surgery, Brigham and Women's Hospital, Boston, MA; ²Breast Oncology Program, Dana-Farber/Brigham and Women's Cancer Center, Boston, MA; ³Center for Surgery and Public Health, Department of Surgery, Brigham and Women's Hospital, Boston, MA; ⁴Department of Radiation Oncology, Dana-Farber Cancer Institute and Brigham and Women's Hospital, Boston, MA

Initiation of a clinical protocol to optimize axillary management of patients with cT1-2N0 breast cancer undergoing mastectomy with one to two positive sentinel lymph nodes has decreased overtreatment of the axilla without compromising outcomes in this patient population (<https://doi.org/10.1245/s10434-021-10726-3>).

FUNDING J.M. acknowledges the Pamela and Nick Gelsomini Breast Surgical Oncology Fellowship Fund for support.

DISCLOSURE T.A.K. reports speakers honoraria and advisory board participation for Exact Sciences (formerly Genomic Health); and Faculty, PrecisCA cancer information service. E.A.M. reports scientific advisory board participation for Exact Sciences (formerly Genomic Health).

Olga Kantor and Jessica Means have contributed equally.

© Society of Surgical Oncology 2021
 Published Online: 8 September 2021

T. A. King, MD
 e-mail: tking7@bwh.harvard.edu

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.