ASO VISUAL ABSTRACT

ASO Visual Abstract: The Value of Paratracheal Lymphadenectomy in Esophagectomy for Adenocarcinoma of the Esophagus or Gastroesophageal Junction: a Systematic Review of the Literature

Amaia Gantxegi, MD¹, B. Feike Kingma, MD, PhD², Jelle P. Ruurda², Grard A. P. Nieuwenhuijzen, Md, PhD³, Misha D. P. Luyer, Md, PhD³, and Richard van Hillegersberg²

¹Vall d'Hebron Hospital Universitari, Barcelona, Spain; ²Department of Surgery, University Medical Center Utrecht, P.O. Box 85500, Utrecht, The Netherlands; ³Catharina Hospital Eindhoven, Eindhoven, The Netherlands

The optimal extent of mediastinal lymphadenectomy for distal esophageal and gastroesophageal junction (GEJ) adenocarcinomas is a matter of debate, especially regarding paratracheal lymphadenectomy. This review (<https://doi.org/10.1245/s10434-021-10810-8>) aimed to provide objective evidence for the reported incidence of lymph node metastasis in the upper mediastinum.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1245/s10434-021-10895-1>.

DISCLOSURE Jelle Ruurda and Richard van Hillegersberg are proctors for Intuitive Surgical Inc. The remaining authors have no conflicts of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.