ASO VISUAL ABSTRACT

ASO Visual Abstract: Non-small Cell Lung Cancer Patients with a High Predicted Risk of Irradical Resection—can Chemoradiotherapy Offer Similar Survival?

W. Hugo van Joolingen, BSc¹, Marnix J. A. Rasing, MD¹, Max Peters, MD, PhD¹, Anne S. R. van Lindert, MD², Linda M. de Heer, MD, PhD³, Mieke J. Aarts, PhD⁴, Joost J. C. Verhoeff, MD, PhD¹, and Peter S. N. van Rossum, MD, PhD¹

¹Department of Radiation Oncology, University Medical Center Utrecht, Utrecht, The Netherlands; ²Department of Pulmonology, University Medical Center Utrecht, Utrecht, The Netherlands; ³Department of Cardiothoracic Surgery, University Medical Center Utrecht, Utrecht, The Netherlands; ⁴Netherlands Cancer Registry, Netherlands Comprehensive Cancer Organization, Utrecht, The Netherlands

Patients with NSCLC with a high predicted risk of irradical resection (per an internationally validated nomogram) treated with surgery were matched to patients who received chemoradiotherapy. Surgery yielded superior

overall survival. Additional factors besides the nomogram should dictate treatment decision-making (<https://doi.org/10.1245/s10434-021-10982-3>).

© Society of Surgical Oncology 2021

J. J. C. Verhoeff, MD, PhD
e-mail: J.J.C.Verhoeff-10@umcutrecht.nl

P. S. N. van Rossum, MD, PhD
e-mail: P.S.N.vanRossum-2@umcutrecht.nl

FUNDING No external funding source was involved in this investigation.

DISCLOSURES The authors have no conflicts of interest relevant to this work.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.