



## ASO Video Abstract: Effect of Phased Implementation of Totally Minimally Invasive Ivor Lewis Esophagectomy for Esophageal Cancer after Previous Adoption of the Hybrid Minimally Invasive Technique: Results from a French Nationwide Population-Based Cohort Study

Frederiek Nuytens, MD<sup>1,2</sup> , Xavier Lenne, MSc<sup>3</sup>, Guillaume Clément, MD<sup>3</sup>, Amelie Bruandet, MD, PhD<sup>3</sup>, Clarisse Eveno, MD, PhD<sup>1,4</sup>, and Guillaume Piessen, MD, PhD<sup>1,4</sup>

<sup>1</sup>Department of Digestive and Oncological Surgery, Claude Huriez University Hospital, University Lille, Lille, France; <sup>2</sup>Department of Digestive and Hepatobiliary/Pancreatic Surgery, AZ Groeninge Hospital, Kortrijk, Belgium; <sup>3</sup>Medical Information Department, Lille University Hospital, Lille, France; <sup>4</sup>CNRS, Inserm, CHU Lille, UMR9020-U1277 - CANTHER - Cancer Heterogeneity, Plasticity and Resistance to Therapies, University Lille, Lille, France

We evaluated nationwide the short-term outcomes of totally minimally invasive esophagectomy after previous adoption of the hybrid minimally invasive technique. Totally minimally invasive esophagectomy is associated with a reduced 90-day postoperative mortality compared with open esophagectomy and offers reduced rates of pulmonary complications compared with open esophagectomy and hybrid minimally invasive techniques (<https://doi.org/10.1245/s10434-021-11110-x>).

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1245/s10434-021-11144-1>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.