ASO VISUAL ABSTRACT

ASO Visual Abstract: Surgical Outcomes and Analysis Comparing Transduodenal Ampullectomy with Pancreaticoduodenectomy: Single-Center Study

Eun-Ki Min, MD¹, Seung Soo Hong, MD¹, Ji Su Kim, MD¹, Munseok Choi, MD², Hyeo Seong Hwang, MD³, Chang Moo Kang, MD, PhD¹, Woo Jung Lee, MD, PhD¹, Dong Sup Yoon, MD, PhD¹, and Ho Kyoung Hwang, MD, PhD¹

¹Division of Hepatobiliary and Pancreatic Surgery, Department of Surgery, Severance Hospital, Seoul, South Korea;

²Department of Surgery, Yongin Severance Hospital, Yonsei University College of Medicine, Seoul, South Korea;

³Department of Surgery, National Health Insurance Service Ilsan Hospital, Ilsan, South Korea

Transduodenal ampullectomy (TDA) with lymph node dissection can be a feasible option for patients with high-grade dysplasia (HGD)/Tis or T1 ampulla of Vater (AoV) carcinoma

in view of superior perioperative outcomes and long-term survival rates comparable with those of pancreaticoduodenectomy (PD) (<https://doi.org/10.1245/s10434-021-11190-9>).

DISCLOSURE There are no conflicts of interest.