ASO VISUAL ABSTRACT

ASO Visual Abstract: The Effect of Preoperative Treatment on the Performance of Predictive Nomograms in Primary Retroperitoneal Sarcoma (RPS)

Deanna Ng, MBBS^{1,2,3,4}, David P. Cyr, MD, MSc^{1,2,3,4}, Sally M. Burtenshaw, MSc^{1,2}, Dario Callegaro, MD⁵, Alessandro Gronchi, MD⁵, David Shultz, MD⁶, Savtaj Brar, MD, MSc^{1,2}, Peter Chung, MD⁶, Rebecca A. Gladdy, MD, PhD^{1,2,4}, Charles Catton, MD⁶, and Carol J. Swallow, MD, PhD^{1,2,3,4}

¹Department of Surgical Oncology, Princess Margaret Cancer Centre/Mount Sinai Hospital, University of Toronto, Toronto, ON, Canada; ²Department of Surgery, University of Toronto, Toronto, ON, Canada; ³Institute of Medical Science, University of Toronto, Toronto, Canada; ⁴Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, Canada; ⁵Department of Surgery, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy; ⁶Department of Radiation Oncology, Princess Margaret Cancer Centre, University of Toronto, Toronto, ON, Canada

We evaluated the predictive accuracy of existing RPS-specific nomograms in patients managed at our center, where majority of patients received preoperative radiotherapy

(<https://doi.org/10.1245/s10434-021-11156-x>). We found that preoperative radiotherapy precludes the use of the LR component of existing nomograms for primary RPS.

DISCLOSURE None of the authors has any conflict of interest to declare