ASO VISUAL ABSTRACT

ASO Visual Abstract: Clinical Relevance of the Tumor Location-Modified Laurén Classification System of Gastric Cancer in a Western Population

Jonathan L. Moore, BMedSci (Hons), BMBS, MRCS^{1,2}, Andrew R. Davies, BSc, MBChB, MD, FRCS^{1,2}, Aida Santaolalla, PhD, MSc, BSc^{2,3}, Mieke van Hemelrijck, PhD, MSc², Nick Maisey, MD, MBBS, FRCP⁴, Jesper Lagergren, MD, PhD^{1,2,5}, James A. Gossage, BSc, MBBS, MS, FRCS^{1,2}, Mark Kelly, BSc (Hons), MBBS, MRes, FRCS^{1,2}, Cara R. Baker, BA (Hons), FRCS, PhD^{1,2}, and on behalf of the Guy's and St Thomas' Esophago-gastric Research Group

¹Department of Upper Gastrointestinal and General Surgery, St Thomas' Hospital, London, UK; ²School of Cancer and Pharmaceutical Sciences, King's College, London, UK; ³Translational Oncology and Urology Research (TOUR), School of Cancer and Pharmaceutical Sciences, King's College, London, UK; ⁴Department of Medical Oncology, St Thomas' Hospital, London, UK; ⁵Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden

The Tumor Location-Modified Laurén Classification system of gastric cancer independently predicted survival in this Western cohort of patients. Patients with proximal

non-diffuse and diffuse tumors had worse survival than those with distal non-diffuse tumors (<https://doi.org/10.1245/s10434-021-11252-y>).

FUNDING No sources of funding were used for this study.