




ASO VISUAL ABSTRACT


ASO Visual Abstract: Importance of Nodal Metastases Location in Pancreatoduodenectomy for Pancreatic Ductal Adenocarcinoma: Results from a Prospective Lymphadenectomy Protocol

Giuseppe Malleo, MD¹ , Laura Maggino, MD¹, Fabio Casciani, MD¹, Gabriella Lionetto, MD¹, Sara Nobile, MD¹, Gianni Lazzarin, MD¹, Salvatore Paiella, MD¹, Alessandro Esposito, MD¹, Paola Capelli, MD², Claudio Luchini, MD², Aldo Scarpa, MD², Claudio Bassi, MD¹, and Roberto Salvia, MD¹

¹Unit of General and Pancreatic Surgery, Department of Surgery, Dentistry, Gynecology and Pediatrics, G.B. Rossi Hospital, University of Verona, Verona, Italy; ²Section of Pathology, Department of Pathology and Diagnostics, University of Verona, Verona, Italy

Prospective implementation of a lymphadenectomy protocol in 424 pancreatoduodenectomies for cancer - showed that first-echelon nodal dissection was sufficient for optimal staging. Nodal metastases occurred mostly at

stations 13/14, although second-echelon involvement was frequent. Only station 14 and jejunal mesentery nodes involvement was prognostically relevant (<https://doi.org/10.1245/s10434-022-11417-3>).



FUNDING None.

DISCLOSURE None.

© Society of Surgical Oncology 2022

Published Online: 2 March 2022

R. Salvia, MD

e-mail: roberto.salvia@univr.it

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.