ASO VISUAL ABSTRACT

ASO Visual Abstract: Development and Validation of the Optimal Circumferential Resection Margin in Pathological T3 Esophageal Cancer: A Multicenter Retrospective Study

Ryoma Haneda, MD^{1,2}, Hirotohi Kikuchi, MD, PhD¹, Yuka Nagakura, MD³, Akifumi Notsu⁴, Eisuke Booka, MD, PhD¹, Tomohiro Murakami, MD, PhD¹, Tomohiro Matsumoto, MD, PhD¹, Shuhei Mayanagi, MD, PhD², Yoshifumi Morita, MD, PhD¹, Yoshihiro Hiramatsu, MD, PhD^{1,5}, Yasuhiro Tsubosa, MD, PhD², and Hiroya Takeuchi, MD, PhD¹

¹Department of Surgery, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan; ²Division of Esophageal Surgery, Shizuoka Cancer Center, Shizuoka, Japan; ³Department of Diagnostic Pathology, Hamamatsu University School of Medicine, Shizuoka, Japan; ⁴Clinical Research Center, Shizuoka Cancer Center, Shizuoka, Japan; ⁵Department of Perioperative Functioning Care and Support, Hamamatsu University School of Medicine, Shizuoka, Japan

This study (<https://doi.org/10.1245/s10434-022-11491-7>) reported that the circumferential resection margin of 600 μ m, set between the Royal College of Pathologists criteria

and the College of American Pathologists criteria, is optimal to predict locoregional recurrence for pathological T3 esophageal squamous cell carcinoma.

DISCLOSURES The authors declare that they have no conflict of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.