





ASO VISUAL ABSTRACT

ASO Visual Abstract: Axillary Response to Neoadjuvant Therapy in Node-Positive ER+/HER2- Breast Cancer Patients—Predictors and Oncologic Outcomes

Orli Friedman-Eldar, MD^{1,2} , Tolga Ozmen, MD^{1,2}, Salah James El Haddi, MD, MS³, Neha Goel, MD², Youley Tjendra, MD⁴, Susan B. Kesmodel, MD², Mecker G. Moller, MD², Dido Franceschi, MD², Christina Layton, DO², and Eli Avisar, MD²

¹Department of Surgical Oncology, Jackson Memorial Hospital, Miami, FL; ²Division of Surgical Oncology, DeWitt Daughtry Family Department of Surgery, University of Miami Miller School of Medicine, Miami, FL; ³Department of Surgery, Charles E. Schmidt College of Medicine, Florida Atlantic University, Boca Raton, FL; ⁴Division of Surgical Pathology, University of Miami Miller School of Medicine, Miami, FL



This study (<https://doi.org/10.1245/s10434-022-11473-9>) retrospectively compared luminal node-positive patients who achieved axillary pathologic complete response after neoadjuvant treatment with those who did not, aiming to assess possible clinical and pathologic predictors of nodal response, as well as correlation between nodal response and oncologic outcomes.

FUNDING The authors received no financial support for the research, authorship, and/or publication of this article.

DISCLOSURE Orli Friedman-Eldar, Tolga Ozmen, Salah James El Haddi, Neha Goel, Youley Tjendra, Susan B. Kesmodel, Mecker G. Moller, Dido Franceschi, Christina Layton, and Eli Avisar declare that they have no conflicts of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.