



ASO VISUAL ABSTRACT


ASO Visual Abstract: Risk Factors for Anastomotic Leakage in Advanced Ovarian Cancer Surgery: A large Single-Center Experience

Barbara Costantini, MD¹, Virginia Vargiu, MD², Francesco Santullo, MD³, Andrea Rosati, MD¹, Matteo Bruno, MD¹, Valerio Gallotta, MD¹, Claudio Lodoli, MD³, Rossana Moroni⁴, Fabio Pacelli, MD^{3,5}, Giovanni Scambia, MD^{1,5}, and Anna Fagotti, PhD^{1,5}

¹Division of Gynecologic Oncology, Department Woman and Child Health and Public Health, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy; ²Department of Oncology, Gemelli Molise SpA, Campobasso, Italy; ³Surgical Unit of Peritoneum and Retroperitoneum, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy; ⁴Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy; ⁵Università Cattolica del Sacro Cuore, Rome, Italy

Anastomotic leakage is a rare but severe complication in advanced ovarian cancer surgery. Improving the patient's nutritional status and maximizing the blood

supply to the anastomotic complex could further reduce its occurrence (<https://doi.org/10.1245/s10434-022-11686-y>).



DISCLOSURES There are no conflict of interest.

© Society of Surgical Oncology 2022

Published Online: 28 April 2022

G. Scambia, MD
e-mail: giovanni.scambia@policlinicogemelli.it

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.