



ASO VISUAL ABSTRACT


## ASO Visual Abstract: The Histopathological Growth Pattern of Colorectal Liver Metastases Impacts Local Recurrence Risk and the Adequate Width of the Surgical Margin

L. Viganò, MD, PhD<sup>1,2</sup> , B. Branciforte, MD<sup>1</sup>, V. Laurenti, MD<sup>1</sup>, G. Costa, MD<sup>1</sup>, F. Procopio, MD<sup>1</sup>, M. Cimino, MD<sup>1</sup>, D. Del Fabbro, MD<sup>1</sup>, L. Di Tommaso, MD<sup>2,3</sup>, and G. Torzilli, MD, PhD, FACS, FAFC(Hon)<sup>1,2</sup>

<sup>1</sup>Division of Hepatobiliary and General Surgery, Department of Surgery, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy; <sup>2</sup>Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy; <sup>3</sup>Pathology Unit, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy

Both the margin width and histopathological growth pattern of colorectal liver metastases (CLM) influence the local recurrence risk. While narrow (1 mm) and vascular

margins are adequate for CLM with nonreplacement pattern, wider margins are needed for those with replacement pattern (<https://doi.org/10.1245/s10434-022-11717-8>).



**FUNDING** No funding to declare.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.