




ASO VISUAL ABSTRACT


ASO Visual Abstract: Management of Malignant Small Bowel Obstruction—Is Intestinal Bypass Effective Palliation?

Meagan Read, MD^{1,2}, Benjamin D. Powers, MD^{1,3,4}, Jose M. Pimiento, MD^{1,3}, Danielle Laskowitz, PA-C¹, Erin Mihelic, PA-C¹, Iman Imanirad, MD¹, Sophie Dessureault, MD^{1,2,3}, Seth Felder, MD^{1,3}, and Sean P. Dineen, MD^{1,3} 

¹Department of Gastrointestinal Oncology, Moffitt Cancer Center, Tampa, FL; ²Department of Surgery, Morsani College of Medicine, Tampa, FL; ³Department of Oncologic Sciences, Morsani College of Medicine, Tampa, FL; ⁴Department of Health Outcomes and Behavior, Moffitt Cancer Center, Tampa, FL

We investigated the outcomes of intestinal bypass for the treatment of malignant small-bowel obstruction (mSBO) (<https://doi.org/10.1245/s10434-022-12204-w>).

We demonstrated that median OS was 6.6 months. The majority of patients were discharged to home (85.5%) and were able to tolerate diet (74.6%). Nearly half of patients



Presented as a Poster at the SSO International Conference on Surgical Cancer Care, March 2022.

© Society of Surgical Oncology 2022

Published Online: 1 August 2022

S. P. Dineen, MD

e-mail: sean.dineen@moffitt.org

were able to return to therapy (49.1%) and demonstrated improved survival of 11.9 months.

DISCLOSURE Authors declare no conflicts of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.