



ASO VISUAL ABSTRACT


ASO Visual Abstract: The Influence of BMI on the Histopathology and Outcomes of Patients with a Diagnosis of Atypical Breast Lesions

Krislyn N. Miller, DO, MS^{1,2}, Samantha M. Thomas, MS³, Amanda R. Sergesketter, MD^{1,2}, Laura H. Rosenberger, MD, MS^{1,2}, Gayle DiLalla, MD^{1,2}, Astrid Botty van den Bruele, MD^{1,2}, E. Shelley Hwang, MD, MPH^{1,2}, and Jennifer K. Plichta, MD, MS^{1,2,4}

¹Department of Surgery, Duke University Medical Center, Durham, NC; ²Duke Cancer Institute, Duke University, Durham, NC; ³Biostatistics and Bioinformatics, Duke University, Durham, NC; ⁴Department of Population Health Sciences, Duke University Medical Center, Durham, NC

For patients with atypical high-risk breast lesions, select factors are associated with the risk of upstage to breast cancer, subsequent development of breast cancer, or both.

In this large cohort of women with a diagnosis of high-risk breast histology (<https://doi.org/10.1245/s10434-022-12313-6>), there was no association with body mass index (BMI).



FUNDING This article is supported by NIH (Grant No. P30CA014236).

DISCLOSURES There are no conflicts of interest.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.