




ASO VISUAL ABSTRACT

ASO Visual Abstract: Risk Factors for Lymph Node Metastasis of Rectal Neuroendocrine Tumor and its Prognostic Impact—A Single-Center Retrospective Analysis of 195 Cases with Radical Resection

Yukiharu Hiyoshi, MD, PhD, FACS¹ , Nobuya Daitoku, MD, PhD¹, Toshiki Mukai, MD, PhD¹, Toshiya Nagasaki, MD, PhD¹, Tomohiro Yamaguchi, MD, PhD¹, Takashi Akiyoshi, MD, PhD¹, Chihiro Yasue, MD², Akiko Chino, MD, PhD², Shoichi Saito, MD, PhD², Manabu Takamatsu, MD, PhD³, and Yosuke Fukunaga, MD, PhD¹

¹Gastroenterological Center, Department of Gastroenterological Surgery, The Cancer Institute Hospital of Japanese Foundation for Cancer Research, Tokyo, Japan; ²Gastroenterological Center, Department of Lower Gastrointestinal Medicine, The Cancer Institute Hospital of Japanese Foundation for Cancer Research, Tokyo, Japan; ³Department of Pathology, The Cancer Institute Hospital of Japanese Foundation for Cancer Research, Tokyo, Japan

Rectal NETs have strong potential of LN metastasis in accordance with an increase of risk factors, and LN metastasis is associated with poor prognosis. This is the first report

demonstrating multiple NETs as a risk factor for LN metastasis (<https://www.doi.org/ASO-2022-10-2888.R1>).

