




ASO VISUAL ABSTRACT


ASO Visual Abstract: Impact of Successful Implementation of an Enhanced Recovery After Surgery (ERAS) Protocol for Patients Undergoing Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy (CRS-HIPEC)

Beatrice J. Sun, MD¹ , Tiffany M. Yue, BA², Nova Xu, BS², Cedar Fowler, MD, PhD³, and Byrne Lee, MD¹

¹Department of Surgery, Section of Surgical Oncology, Stanford University School of Medicine, Stanford, CA; ²Stanford University School of Medicine, Stanford, CA; ³Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University School of Medicine, Stanford, CA

Enhanced recovery after surgery (ERAS) protocols are feasible and safe in the setting of CRS-HIPEC. This resulted in decreased hospital length of stay without increase in readmission rate; furthermore, major complications were

low. Adherence to postoperative ERAS components was 76%, which is encouraging and supports the use of ERAS programs for patients undergoing CRS-HIPEC (<https://doi.org/10.1245/s10434-023-14222-8>).



DISCLOSURES None.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.