



ASO VISUAL ABSTRACT


ASO Visual Abstract: Actual 5-Year Survival After Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy for Patients with Peritoneal Carcinomatosis of Colorectal Origin

Elad Sarfaty, MD, Nazanin Khajouejad, MD, Allen T. Yu, MD, PhD, Spiros Hiotis, MD, PhD, Benjamin J. Golas, MD, Umut Sarpel, MD, Daniel M. Labow, MD, and Noah A. Cohen, MD

Division of Surgical Oncology, Department of Surgery, Icahn School of Medicine at Mount Sinai, New York, NY

We report actual 5-year survival in patients with peritoneal carcinomatosis (PC) of colorectal (CRC) origin who undergo cytoreductive surgery–hyperthermic intraperitoneal chemotherapy (CRS/HIPEC) (<https://doi.org/10.1245/s10434-023-14608-8>), and demonstrate that well-selected patients with low PCI who undergo complete surgical resection can achieve favorable long-term survival.

and demonstrate that well-selected patients with low PCI who undergo complete surgical resection can achieve favorable long-term survival.



Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.