



ASO VISUAL ABSTRACT

ASO Visual Abstract: Local Therapy Improves Survival for Early Recurrence After Resection of Colorectal Liver Metastases

Artem Boyev, DO¹, Ching-Wei D. Tzeng, MD¹, Harufumi Maki, MD¹, Elsa M. Arvide, PA-C¹, Deborah E. Mrema, FNP-C¹, Anish J. Jain, MD¹, Antony Haddad, MD¹, Mateo Lendoire, MD¹, Neha Malik, MD, MS¹, Bruno C. Odisio, MD², Yun Shin Chun, MD¹, Hop S. Tran Cao, MD¹, Jean-Nicolas Vauthey, MD¹, and Timothy E. Newhook, MD¹

¹Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX; ²Department of Interventional Radiology, The University of Texas MD Anderson Cancer Center, Houston, TX

Early recurrence after hepatectomy for colorectal liver metastases has a worse prognosis compared to later recurrence. Local therapy is associated with improved survival regardless of timing of recurrence. Older age, extrahepatic

recurrence, and deleterious mutations are associated with worse survival after treatment with local therapy (<https://doi.org/10.1245/s10434-023-14806-4>).

