



ASO VISUAL ABSTRACT

ASO Visual Abstract: Minimally Invasive Approach Provides Oncological Benefit in Patients with High Risk of Very Early Recurrence (VER) After Surgery for Intrahepatic Cholangiocarcinoma (iCCA)

Francesca Ratti, MD^{1,2}, Cecilia Maina, MD¹, Lucrezia Clocchiatti, MD¹, Rebecca Marino, MD¹, Federica Pedica, MD³, Andrea Casadei Gardini, MD^{2,4}, Francesco De Cobelli, MD^{2,5,6}, and Luca Antonio Maria Aldrighetti, MD, PhD^{1,2}

¹Hepatobiliary Surgery Division, IRCCS San Raffaele, Milan, Italy; ²Vita-Salute San Raffaele University, Milan, Italy; ³Department of Experimental Oncology, Pathology Unit, San Raffaele Hospital, Milan, Italy; ⁴Department of Medical Oncology, IRCCS San Raffaele Hospital, Milan, Italy; ⁵Department of Radiology, IRCCS San Raffaele Hospital, Milan, Italy; ⁶Experimental Imaging Center, IRCCS San Raffaele Scientific Institute, Milan, Italy

In patients with high risk of very early recurrence (VER), laparoscopy translates into an advantage in terms of recurrence-free survival, likely related to lower biological impact of surgery, together with a shorter interval between

surgery and start of adjuvant treatments, even allowing for a higher number of patients to start adjuvant therapies within 2 months of resection (<https://doi.org/10.1245/s10434-023-14807-3>).

