GC Analysis of the Fatty Acid Composition of Yak Kidney

Qiang Wang¹, Jianping Wu^{2,3,✉}, Shenggui Zhang¹, Yubin Zhang¹, Haixia Zhang⁴, Enguo Fan^{1,5,✉}

¹ College of Food Science and Engineering, Gansu Agricultural University, 730070 Lanzhou, China

² College of Animal Science and Technology, Gansu Agricultural University, 730070 Lanzhou, China

³ Center for Engineering and Technological Research on Livestock Products of Gansu Province, 730070 Lanzhou, China;
E-Mail: jp-wu@hotmail.com

⁴ College of Agronomy, Gansu Agricultural University, 730070 Lanzhou, China

⁵ Institut für Biochemie und Molekularbiologie, Universität Freiburg, Stefan-Meier-Strasse 17, 79104 Freiburg, Germany;
E-Mail: enguo.fan@biochemie.uni-freiburg.de

Erratum to: *Chromatographia* (2009) 69:139–143 DOI 10.1365/s10337-008-0845-6

Unfortunately, in the original article in the “Experimental”/“GC Analysis” a wrong column is specified.

FAME analysis was performed with a Clarus500 gas chromatograph (Perkin–Elmer, USA) equipped with a CP Sil 88 column (120 m × 0.32 mm i.d.; film thickness 0.2 µm).

The actual column is a SGE-FFAP column (120 m × 0.25 mm i.d.; film thickness 0.25 µm).

The online version of the original article
can be found under doi:[10.1365/s10337-008-0845-6](https://doi.org/10.1365/s10337-008-0845-6).