



Identification of a novel mutation in a Chinese family with Nance-Horan syndrome by whole exome sequencing*

Nan HONG^{§1}, Yan-hua CHEN^{§1}, Chen XIE¹, Bai-sheng XU¹, Hui HUANG², Xin LI², Yue-qing YANG², Ying-ping HUANG², Jian-lian DENG², Ming QI^{†‡2,3,4}, Yang-shun GU^{†‡1}

(¹Department of Ophthalmology, the First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310003, China)

(²Beijing Genomics Institute (BGI)-Shenzhen, Shenzhen 518083, China)

(³School of Basic Medical Sciences, Zhejiang University, Hangzhou 310058, China)

(⁴Functional Genomics Center, Department of Pathology & Laboratory Medicine, University of Rochester Medical Center, West Henrietta, NY 14586, USA)

[†]E-mail: mingqi@zju.edu.cn; guyangshun_1@hotmail.com

Received Dec. 22, 2013; Revision accepted June 9, 2014; Crosschecked July 23, 2014

Abstract: Objective: Nance-Horan syndrome (NHS) is a rare X-linked disorder characterized by congenital nuclear cataracts, dental anomalies, and craniofacial dysmorphisms. Mental retardation was present in about 30% of the reported cases. The purpose of this study was to investigate the genetic and clinical features of NHS in a Chinese family. Methods: Whole exome sequencing analysis was performed on DNA from an affected male to scan for candidate mutations on the X-chromosome. Sanger sequencing was used to verify these candidate mutations in the whole family. Clinical and ophthalmological examinations were performed on all members of the family. Results: A combination of exome sequencing and Sanger sequencing revealed a nonsense mutation c.322G>T (E108X) in exon 1 of *NHS* gene, co-segregating with the disease in the family. The nonsense mutation led to the conversion of glutamic acid to a stop codon (E108X), resulting in truncation of the NHS protein. Multiple sequence alignments showed that codon 108, where the mutation (c.322G>T) occurred, was located within a phylogenetically conserved region. The clinical features in all affected males and female carriers are described in detail. Conclusions: We report a nonsense mutation c.322G>T (E108X) in a Chinese family with NHS. Our findings broaden the spectrum of *NHS* mutations and provide molecular insight into future NHS clinical genetic diagnosis.

Key words: Nance-Horan syndrome (NHS), Exome sequencing, X-linked disorder

doi:10.1631/jzus.B1300321

Document code: A

CLC number: R776.1

1 Introduction

Nance-Horan syndrome (NHS) (OMIM: 302350) is a rare X-linked syndrome, first reported in detail by Horan and Billson (1974) and Nance *et al.* (1974). Affected male patients often manifest bilateral congenital dense nuclear cataracts, characteristic dental

anomalies, craniofacial abnormalities, and, in some cases, mental retardation (MR) (Toutain *et al.*, 1997; Ding *et al.*, 2009). Congenital dense nuclear cataracts in affected males usually lead to profound vision loss; thus, cataract surgery should be performed at an early age (Burdon *et al.*, 2003). Other ophthalmological features include microcornea, microphthalmia, nystagmus, and strabismus. Dental abnormalities characteristic of NHS consist of screwdriver-shaped incisors, supernumerary maxillary incisors (mesiodens), and diastema (Walpole *et al.*, 1990). Heterozygous females display similar, but milder clinical manifestations than affected males (Ding *et al.*, 2009).

[‡] Corresponding authors

[§] The two authors contributed equally to this work

* Project supported by the Science and Technology Specific Project of Zhejiang Province (No. 2009C03010-2), China

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2014

NHS is caused by mutations in the *NHS* gene, which comprises 10 coding exons and encodes at least 5 isoforms as a result of alternative splicing (Brooks *et al.*, 2010). NHS-A and NHS-1A are two major isoforms of the NHS protein. The two isoforms are transcribed from exon 1 of the *NHS* gene, encoding a 1630-amino-acid protein and a 1651-amino-acid protein, respectively (Brooks *et al.*, 2010). The *NHS* gene is expressed in the midbrain, lens, tooth, retina, and dental primordia, in a spatially and temporally regulated pattern, and is highly conserved across vertebrate species, such as mouse, rat, and zebrafish (Huang *et al.*, 2007; Sharma *et al.*, 2008).

To date, approximately 27 mutations in the *NHS* gene have been identified (Burdon *et al.*, 2003; Brooks *et al.*, 2004; Ramprasad *et al.*, 2005; Florijn *et al.*, 2006; Huang *et al.*, 2007; Sharma *et al.*, 2008; Coccia *et al.*, 2009; Chograni *et al.*, 2011; Khan *et al.*, 2012; Tug *et al.*, 2013), including nonsense mutations, frameshift mutations, genomic rearrangements, and one missense mutation. In this study, we identified a novel nonsense mutation c.322G>T (E108X) in the *NHS* gene's exon 1 in a family with NHS by a combination of exome sequencing and Sanger sequencing. The clinical features in all affected males and female carriers are described in detail.

2 Materials and methods

2.1 Subjects and ethics approval

This study followed the tenets of the Declaration of Helsinki and was approved by the Ethics Committee of the First Affiliated Hospital, School of Medicine, Zhejiang University, China. A three-generation family with congenital cataracts, microcornea, strabismus, and nystagmus was recruited. Seven members of the family participated in the study, three affected (II:1, II:2, and III:1) and four unaffected (I:1, I:2, I:3, and II:3) members. Informed consent and blood samples were obtained from all the participants in the family. All participants underwent comprehensive ophthalmological examinations including visual acuity, slit-lamp biomicroscopy, fundus examination, intraocular pressure (IOP) measurement, and IOLMaster testing. Nonocular features related to NHS were also noted. A total of 50 healthy individuals without congenital ocular diseases were

selected serving as an ethnically-matched control. Genomic DNA was extracted using the QIAamp DNA Blood Midi kit (Qiagen, Hilden, Germany), according to the manufacturer's protocol.

2.2 Exome sequencing

One affected male (III:1) in the family was selected for exome sequencing. Exome sequencing was carried out using the Agilent SureSelect Human All Exon v4.0 (51M) kit, according to the manufacturer's protocol. The 3 μ g of genomic deoxyribonucleic acid (DNA) was randomly fragmented into pieces 200–300 bp in size using a Covaris acoustic system, followed by end-repair, A-tailing, and Covaris acoustic measurements. The adapter-ligated templates were purified using Agencourt AMPure SPRI beads and amplified by four-cycle ligation-mediated polymerase chain reaction (LM-PCR), under the following PCR condition: 2 min at 94 °C, four cycles of 10 s at 94 °C, 30 s at 62 °C, and 30 s at 72 °C, then 300 s at 72 °C. The LM-PCR products were hybridized to the Agilent Oligo pool for enrichment for 24 h at 65 °C. Hybridized fragments were bound to streptavidin beads and non-hybridized fragments were washed out. Captured LM-PCR products were amplified by PCR (2 min at 98 °C; 10–12 cycles for 10 s at 98 °C, 30 s at 60 °C, 30 s at 72 °C, followed by 300 s at 72 °C). The magnitude of enrichment of the samples was estimated with an Agilent 2100 bioanalyzer and quantitative PCR. The captured library was sequenced on Illumina HiSeq2000 analyzers for 90 cycles per read, to generate paired-end reads and 8 bp of the index tag (following the manufacturer's standard sequencing protocols).

2.3 Mapping, single-nucleotide polymorphism (SNP), and insertion and deletion (InDel) calling

Image analysis and base calling were performed according to the Illumina pipeline. Indexed primers were used for data fidelity surveillance. Short Oligonucleotide Analysis Package (SOAP) aligner (Ver. 2.21) was used to map the clean reads to the human reference genome (hg19) with a maximum fault-tolerance of 3-bp mismatches. Based on these results, the SOAPsnp software (Ver. 1.05) was used to assemble the consensus sequence and call genotypes in target regions. Candidate single-nucleotide SNPs were selected using the following criteria: SNP quality

≥ 20 , sequencing depth ≥ 4 -fold, estimated copy number ≤ 2 , and distance between two SNPs > 5 bp. For small InDels < 20 bp in exome regions, we mapped the reads to the reference genome with the Burrows Wheeler Aligner (BWA; Ver. 0.6.2), and passed the result to the Genome Analysis Tool kit (GATK) to identify breakpoints.

2.4 PCR and Sanger sequencing

Following the results of exome sequencing, Sanger sequencing was performed to verify the genetic defects. Primers flanking the mutation area of *NHS* (NM_198270) were designed, based on the reference genomic sequences of the human genome from GenBank at the National Center for Biological Information (NCBI), and synthesized by Invitrogen, Shanghai, China. The forward primer was 5'-TTCCG CCAAGCGGATCGTGGA-3', and the reverse primer was 5'-TTAGGGTCAAGCGTGCTGAGGA-3'. PCR amplification was carried out using an Applied Biosystems Inc. (ABI) 9700 Thermal Cycler. PCR products were directly sequenced on an ABI PRISM 3730 automated sequencer (Applied Biosystems). Sequence data comparisons and analyses were performed using DNASTAR SeqMan. Co-segregation analysis was subsequently performed on the DNA samples from available family members.

2.5 Bioinformatics analysis

Multiple sequence alignment of NHS protein sequences from *Homo sapiens* (GenPept accession NP_938011.1), *Mus musculus* (GenPept accession NP_001074521.1), *Bos taurus* (GenPept accession NP_001192657.1), *Macaca mulatta* (GenPept accession XP_001104229.2), *Canis lupus familiaris* (GenPept accession XP_548877.2), *Sus scrofa* (GenPept accession XP_005673517.1), and *Equus caballus* (GenPept accession XP_005614077.1) was performed using CLC Sequence Viewer 5.5 (CLC bio A/S, <http://www.clcbio.com>).

3 Results

3.1 Clinical features

The family included three affected members (II:1, II:2, and III:1), all males. A male in the third generation was adopted at birth and lost contact with

other members of the family. Information on this male member was missing (Fig. 1a). We suggest that the mode of inheritance of the pedigree was X-linked or autosomal.

A detailed description of clinical features is given in Table 1. All affected males had received bilateral lensectomy at an early age to remove congenital nuclear cataracts. All presented with bilateral microcornea, nystagmus, and strabismus. Individual III:1 (Fig. 1b) exhibited typical NHS dental anomalies (screwdriver-shaped left incisor and diastema, and the right incisor of the patient was broken in childhood, as shown in Fig. 1c), which were not noted in the other




Fig. 1 Pedigree structure and patient phenotypes of the family

(a) Pedigree of a family with Nance-Horan syndrome (NHS). The male identified by a question mark may have been affected, and was adopted out of the family. (b) A photograph of an affected male (III:1) showing a long and narrow face, broad base to the nose, and a thin nasal bridge. (c) Screwdriver-shaped incisors and diastema in individual III:1. The right incisor of the patient was broken in his childhood. (d) Slit-lamp photograph of a carrier female (II:3) showing lens opacities in the posterior Y-suture and cortical coralliform opacity in the left eye. (e, f) Bitemporal retraction in two female carriers (II:3 and I:1)

Table 1 Clinical features of the affected males and the female carriers in the NHS family

Subject	Eye	BSCVA	Ocular feature	Dental anomaly	Craniofacial dysmorphism
Male patients					
II:1	OD	20/400	Underwent bilateral lensectomy due to congenital nuclear cataract, bilateral microcornea, strabismus, and nystagmus	Screwdriver-shaped left incisor* and diastema	Long and narrow face, broad base to nose, and thin nasal bridge
	OS	20/160			
II:2	OD	20/400	Underwent bilateral lensectomy due to congenital nuclear cataract, bilateral microcornea, strabismus, and nystagmus		Long and narrow face, broad base to nose, and thin nasal bridge
	OS	20/160			
III:1	OD	20/120	Underwent bilateral lensectomy due to congenital nuclear cataract, bilateral microcornea, strabismus, and nystagmus		Long and narrow face, broad base to nose, and thin nasal bridge
	OS	20/120			
Female carriers					
I:1	OD	20/60	Opacities in the Y-suture and cortex of the lens in both eyes		Bitemporal retraction
	OS	20/40			
II:3	OD	20/20	Lens opacities in the posterior Y-suture		Bitemporal retraction
	OS	20/30	Lens opacities in the posterior Y-suture and cortical coralliform opacity		

BSCVA: best spectacle corrected visual acuity; OD: the right eye; OS: the left eye. * The right incisor of the patient was broken in his childhood

two affected males. All affected males had the same craniofacial dysmorphisms (a long, narrow face with a broad base to the nose and a thin nasal bridge). Other nonocular anomalies, such as MR, brachymetacarpalia, or cardiovascular abnormalities, were not observed in these three males.

Carrier female II:3 had bilateral fine white dots in the Y-suture of the lenses and a prominent white coralliform cortical opacity in the inferior part of the lens in the left eye (Fig. 1d). Individual I:1 exhibited cortical opacities in both eyes. The two females exhibited bitemporal retraction (Figs. 1e and 1f). No other symptoms related to NHS were observed in the two carrier females.

3.2 Exome sequencing

Using the Agilent SureSelect capture system and the bioinformatics pipeline described in Section 2.5, a mean depth of 84-fold was obtained for this sample, and 95.03% of the target exome region was covered at least 20-fold, indicating that sufficient coverage was achieved for variant calling (Table 2). In total, 103 897 single nucleotide variants (SNVs) and 11 656 small InDels were detected (Table 3). Variants leading to protein coding changes, including missense, nonsense, readthrough, and small InDel variants in the coding region, and potential splice site variants (± 10 bp of coding sequence (CDS)), were picked first. Then we used '1000 Genome' data, dbSNP, and the HapMap database to filter out polymorphism variants.

Table 2 Sequencing quality statistics of exome capture for the proband sample (individual III:1)

Parameter	Value
Total target exome region (bp)	51 391 525
Raw reads	93 225 420
Raw data yield (Mb)	8390
Reads mapped to genome	72 489 626
Reads mapped to target region ¹	55 230 353 (59.24%)
Data mapped to target region (Mb)	4316.82 (51.45%)
Capture specificity ² (%)	76.82
Mean depth of target region (fold)	84
Mean depth of chromosome X (fold)	48.24
Mean depth of chromosome Y (fold)	48.37
Rate of nucleotide mismatch (%)	0.17
Fraction of target covered (%)	
≥ 4 -fold	99.42
≥ 10 -fold	98.32
≥ 20 -fold	95.03

¹ Reads mapped to target regions are reads that are within or overlap with the target region; ² Capture specificity is defined as the percentage of uniquely mapped reads aligning to the target region

Variants with a frequency >0.01 in any of these three databases were filtered out. As a result, we identified 1268 rare variants in the affected male. We first screened these variants for inheritance in an X-linked manner and identified 12 variants on the X chromosome (Table 4). Then, we searched for mutations of known genes involved in hereditary cataracts by visual inspection. The variant c.322G>T in the *NHS* gene was identified as a candidate mutation.


Table 3 Variant statistics of exome capture sequencing for the proband sample (individual III:1) and variant filter statistics

Parameter	Number	
	SNV	InDel
Total	103897	11656
Alt_start	14	
Synonymous	12519	
Missense	8124	
Nonsense	60	
Readthrough	4	
Frameshift		92
Inframe		232
Splice	2299	362
Intron	65274	8979
5-UTR	1928	205
3-UTR	4136	588
Intergenic	9539	1198
Hom	42783	5148
Het	61114	6508
Total (SNVs+InDels)	115553	
Variants leading to protein coding change	11187	
Rare variants leading to protein coding change	1268	
Rare variants in chromosome X	12	
Rare variants in <i>NHS</i> gene	1	

SNV: single nucleotide variant; InDel: insertion and deletion; Alt_start: start codon change to an alternative start codon, which can be considered as a start codon coding-synonym; UTR: untranslated regions; Hom: homozygous; Het: heterozygous

3.3 *NHS* mutation analysis and bioinformatics analysis

A previously undescribed nonsense mutation, c.322G>T, in exon 1 of the *NHS* gene (NM_198270), was identified in the three affected males and in the two carrier females (Fig. 2), but was not detected in the unaffected males or normal controls. This mutation resulted in conversion of glutamic acid to a stop codon at position 108 (E108X). The mutation was not documented in any current database (1000 Genomes Project or HapMap), indicating that the mutation was unlikely to represent an SNP. Multiple sequence alignments were performed and we observed that codon 108, where the mutation (c.322G>T) occurred, was located within a phylogenetically conserved region (Fig. 3).

**Fig. 2 DNA sequence chromatograms**

A nonsense mutation in exon 1, c.322G>T, resulting in E108X in the three affected males and two female carriers, and the normal sequence from two unaffected males

Table 4 Rare variants leading to protein coding changes on the chromosome X

Gene	Function	Variant	Frequency			rs ID
			dbSNP	1000 Genome	HapMap	
<i>CXorf59</i>	Missense	c.1267C>G				
<i>CXorf59</i>	Frameshift	c.1267_1268insTG				
<i>EDA2R</i>	Missense	c.323C>T	0.001			rs186691097
<i>GPR112</i>	Inframe	c.7966_7968delGAT				
<i>IL13RA2</i>	Missense	c.907C>A				
<i>MAP3K15</i>	Splice	c.3567-8delT				
<i>NHS</i>	Nonsense	c.322G>T				
<i>NUDT11</i>	Splice	c.1_1_13delGCTGCCTCGAGGA				
<i>PRDX4</i>	Splice	c.242-10 A>T	0.001			rs1513572
<i>TEX11</i>	Splice	c.1202-4delT				
<i>ZBTB33</i>	Inframe	c.563_564insTGA				
<i>ZNF182</i>	Missense	c.282C>A	0.006	0.0065		rs150066211

rs ID: reference SNP ID number




Fig. 3 Multiple-sequence alignment of NHS from different species

Multiple-sequence alignment of NHS from different species revealed that codon 108, where the mutation (E108X) occurred, was located within a highly conserved region

4 Discussion

In the present study, we characterized a Chinese family with NHS. Molecular genetic analysis identified a novel nonsense mutation c.322G>T (E108X) in exon 1 of the *NHS* gene in all affected males and carrier females. The mutation was not detected in unaffected males or in normal controls. This substitution was conserved in seven tested species, confirming it as a co-segregating disease-causing mutation in the family.

NHS is a rare X-linked disorder, usually underdiagnosed by ophthalmologists with subtle nonocular manifestations that are easily overlooked (Ding *et al.*, 2009). The affected males in the Chinese family presented with congenital dense cataract, microcornea, nystagmus, and strabismus; nonocular symptoms were not investigated at the initial meeting. We performed exome sequencing of DNA from an affected male (III:1) and identified a mutation in the *NHS* gene, which was then shown to segregate with the phenotype in the pedigree. After identification of the mutation, we recalled the family to examine the pedigree members comprehensively. All three affected males manifested craniofacial dysmorphisms characteristic of NHS. Individual III:1 (Fig. 1b) also exhibited typical NHS dental anomalies (screwdriver-shaped incisors and diastema). Among the heterozygous females, carrier signs of NHS were also detected, including lens opacities in the posterior Y-sutures and subtle craniofacial dysmorphism (bitemporal retraction). All of the observations indicated the presence of an NHS phenotype in the family, consistent with the

results of the molecular genetic analysis.

NHS is a syndrome with high phenotypic heterogeneity. Tug *et al.* (2013) summarized the phenotypes of patients with different mutations and found no obvious genotype-phenotype correlations between the position of each mutation and clinical severity. In our study, nonocular features among the members of the family were also not identical; for example, dental anomalies were observed only in individual III:1 (Fig. 1b). The phenotypic heterogeneity may be due to the role of modifier genes in nonocular tissues (Sharma *et al.*, 2006).

The exact biological function and structure of the NHS protein remain unclear. On the basis of position-specific iterated BLAST (PSI-BLAST) analysis, Huang *et al.* (2006) reported that the N-terminus of the mouse *Nhs1* protein contained a poly-proline domain and a Wiskott-Aldrich syndrome protein (WASP) homology 1 (WH1) domain. Both of these domains are known to be related to targeting proteins to the cytoskeleton. Evidence from a study of Brooks *et al.* (2010) showed that the N-termini of isoforms NHS-1A and NHS-A contain a WASP family verprolin-homologous (WAVE) domain (WHD), functionally homologous to the WHD in WAVE proteins. This can interact with members of the Abelson-interactor (Abi) protein family, such as Abi1 and Abi2, playing a regulatory role in actin remodeling and maintaining cell morphology.

In the present study, we describe the first Chinese case of NHS caused by a nonsense mutation in the *NHS* gene. The mutation (c.322G>T) induces a putative premature stop codon at nucleotide 322 (E108X), producing a truncated NHS protein. The abnormal protein may initiate the nonsense-mediated mRNA decay (NMD) pathway, a well-known translation-coupled quality control system that degrades mutant mRNA containing premature termination codons (PTCs) (Khajavi *et al.*, 2006). NMD can protect cells from the toxic effects of truncated proteins caused by dominant-negative or gain-of-function effects (Khajavi *et al.*, 2006). Previously, Ramprasad *et al.* (2005) performed reverse transcription-PCR (RT-PCR) on DNA samples from an Indian NHS family with the c.115C>T (Q39X) mutation in exon 1 of the *NHS* gene and found no evidence of NMD.

It has been reported that some mRNAs harbouring PTCs located in some specific regions can

avoid the NMD pathway, leading to the translation of abnormal mRNAs (Khajavi *et al.*, 2006). If the mRNA containing the c.322G>T (E108X) mutation in the present study escaped from the NMD pathway, amounts of truncated protein with possibly destructive effects would be produced in cells. Also, the E108X mutation may damage the function of the WHD of the NHS protein, for example by disrupting the interaction between the WHD and some other functional protein, resulting in an NHS phenotype in patients. However, the pathogenic mechanism of NHS should be investigated in depth in future.

In summary, we identified a novel nonsense mutation (c.322G>T) in exon 1 of the *NHS* gene in a Chinese family, which leads to the conversion of glutamic acid to a stop codon (E108X). Our study shows that whole exome analysis in a single affected male from a family exhibiting X-linked inheritance could be used efficiently to identify causative mutations on the chromosome X, thus providing help to clinicians in making a definitive diagnosis of rare diseases.

Acknowledgements

We greatly appreciate the invaluable cooperation and participation of the family members in the present study.

Compliance with ethics guidelines

Nan HONG, Yan-hua CHEN, Chen XIE, Bai-sheng XU, Hui HUANG, Xin LI, Yue-qing YANG, Ying-ping HUANG, Jian-lian DENG, Ming QI, and Yang-shun GU declare that they have no conflict of interest.

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008 (5). Informed consent was obtained from all patients for inclusion in the study. Additional informed consent was obtained from all patients for the use of identifying information included in this article.

References

- Brooks, S.P., Ebenezer, N.D., Poopalasundaram, S., *et al.*, 2004. Identification of the gene for Nance-Horan syndrome (NHS). *J. Med. Genet.*, **41**(10):768-771. [doi:10.1136/jmg.2004.022517]
- Brooks, S.P., Coccia, M., Tang, H.R., *et al.*, 2010. The Nance-Horan syndrome protein encodes a functional wave homology domain (WHD) and is important for co-ordinating actin remodelling and maintaining cell morphology. *Hum. Mol. Genet.*, **19**(12):2421-2432. [doi:10.1093/hmg/ddq125]
- Burdon, K.P., McKay, J.D., Sale, M.M., *et al.*, 2003. Mutations in a novel gene, *NHS*, cause the pleiotropic effects of Nance-Horan syndrome, including severe congenital cataract, dental anomalies, and mental retardation. *Am. J. Hum. Genet.*, **73**(5):1120-1130. [doi:10.1086/379381]
- Chograni, M., Rejeb, I., Jemaa, L.B., *et al.*, 2011. The first missense mutation of *NHS* gene in a Tunisian family with clinical features of NHS syndrome including cardiac anomaly. *Eur. J. Hum. Genet.*, **19**(8):851-856. [doi:10.1038/ejhg.2011.52]
- Coccia, M., Brooks, S.P., Webb, T.R., *et al.*, 2009. X-linked cataract and Nance-Horan syndrome are allelic disorders. *Hum. Mol. Genet.*, **18**(14):2643-2655. [doi:10.1093/hmg/ddp206]
- Ding, X., Patel, M., Herzlich, A.A., *et al.*, 2009. Ophthalmic pathology of Nance-Horan syndrome: case report and review of the literature. *Ophthalmic Genet.*, **30**(3):127-135. [doi:10.1080/13816810902822021]
- Florijn, R.J., Loves, W., Maillette de Buy Wenniger-Prick, L.J., *et al.*, 2006. New mutations in the *NHS* gene in Nance-Horan syndrome families from the Netherlands. *Eur. J. Hum. Genet.*, **14**(9):986-990. [doi:10.1038/sj.ejhg.5201671]
- Horan, M.B., Billson, F.A., 1974. X-linked cataract and Hutchinsonian teeth. *Aust. Paediatr. J.*, **10**(2):98-102.
- Huang, K.M., Wu, J., Duncan, M.K., *et al.*, 2006. Xcat, a novel mouse model for Nance-Horan syndrome inhibits expression of the cytoplasmic-targeted *Nhs1* isoform. *Hum. Mol. Genet.*, **15**(2):319-327. [doi:10.1093/hmg/ddi449]
- Huang, K.M., Wu, J., Brooks, S.P., *et al.*, 2007. Identification of three novel *NHS* mutations in families with Nance-Horan syndrome. *Mol. Vis.*, **13**:470-474.
- Khajavi, M., Inoue, K., Lupski, J.R., 2006. Nonsense-mediated mRNA decay modulates clinical outcome of genetic disease. *Eur. J. Hum. Genet.*, **14**(10):1074-1081. [doi:10.1038/sj.ejhg.5201649]
- Khan, A.O., Aldahmesh, M.A., Mohamed, J.Y., *et al.*, 2012. Phenotype-genotype correlation in potential female carriers of X-linked developmental cataract (Nance-Horan syndrome). *Ophthalmic Genet.*, **33**(2):89-95. [doi:10.3109/13816810.2011.634881]
- Nance, W.E., Warburg, M., Bixler, D., *et al.*, 1974. Congenital X-linked cataract, dental anomalies and brachymetacarpalia. *Birth Defects Orig. Artic. Ser.*, **10**(4):285-291.
- Ramprasad, V.L., Thool, A., Murugan, S., *et al.*, 2005. Truncating mutation in the *NHS* gene: phenotypic heterogeneity of Nance-Horan syndrome in an Asian Indian family. *Invest. Ophthalmol. Vis. Sci.*, **46**(1):17-23. [doi:10.1167/iovs.04-0477]
- Sharma, S., Ang, S.L., Shaw, M., *et al.*, 2006. Nance-Horan syndrome protein, NHS, associates with epithelial cell junctions. *Hum. Mol. Genet.*, **15**(12):1972-1983. [doi:10.1093/hmg/ddl120]
- Sharma, S., Burdon, K.P., Dave, A., *et al.*, 2008. Novel causative mutations in patients with Nance-Horan syndrome

- and altered localization of the mutant NHS-A protein isoform. *Mol. Vis.*, **14**:1856-1864.
- Toutain, A., Ayrault, A.D., Moraine, C., 1997. Mental retardation in Nance-Horan syndrome: clinical and neuropsychological assessment in four families. *Am. J. Med. Genet.*, **71**(3):305-314. [doi:10.1002/(SICI)1096-8628(19970822)71:3<305::AID-AJMG11>3.0.CO;2-O]
- Tug, E., Dilek, N.F., Javadiyan, S., et al., 2013. A Turkish family with Nance-Horan syndrome due to a novel mutation. *Gene*, **525**(1):141-145. [doi:10.1016/j.gene.2013.03.094]
- Walpole, I.R., Hockey, A., Nicoll, A., 1990. The Nance-Horan syndrome. *J. Med. Genet.*, **27**(10):632-634. [doi:10.1136/jmg.27.10.632]

中文概要:

本文题目: 全外显子组测序发现一个中国 Nance-Horan 综合征家系 NHS 基因的新突变

Identification of a novel mutation in a Chinese family with Nance-Horan syndrome by whole exome sequencing

研究目的: 通过对一个中国 Nance-Horan 综合征家系的临床表型及基因突变分析, 揭示本家系的致病遗传机制。

研究方法: 对该 Nance-Horan 综合征家系的一个男性患者进行全外显子组测序, 结合此家系临床表型及遗传方式分析, 选定 X 染色体上 NHS 基因上的一个无义突变 c.322G>T (E108X) 为可疑致病突变。通过聚合酶链式反应 (PCR) 和 Sanger 测序, 对该家系内其他成员进行 NHS 基因突变分析, 同时对 50 名健康对照者的 NHS 基因的突变检测结果进行对比。另外, 将该突变的位点第 108 位氨基酸残基进行多物种 NHS 蛋白内序列比对。最后, 对该家系成员眼部及全身的临床特点进行全面检查和分析。

重要结论: 全外显子组测序结合 Sanger 测序发现 NHS 基因第一个外显子上的 c.322G>T (E108X) 突变为引起该家系临床病变的突变位点; 多物种 NHS 蛋白内序列比对发现该突变位点第 108 位氨基酸残基位于高度保守区; 临床表型分析发现该家系内存在表型异质性。此家系为国内首次报道的无义突变引起的 Nance-Horan 综合征家系。

关键词组: Nance-Horan 综合征; NHS; 外显子测序; X-连锁