



Molecular variability and evolution of a natural population of tomato yellow leaf curl virus in Shanghai, China^{*}

Xiu-ling YANG, Meng-ning ZHOU, Ya-juan QIAN, Yan XIE^{†‡}, Xue-ping ZHOU

(State Key Laboratory of Rice Biology, Institute of Biotechnology, Zhejiang University, Hangzhou 310058, China)

[†]E-mail: xiey@zju.edu.cn

Received Apr. 11, 2013; Revision accepted Aug. 21, 2013; Crosschecked Jan. 19, 2014

Abstract: Tomato yellow leaf curl virus (TYLCV), belonging to the genus *Begomovirus* of the family Geminiviridae, is emerging as the most destructive pathogen of tomato plants. Since the first report of TYLCV in Shanghai, China in 2006, TYLCV has spread rapidly to 13 provinces or autonomous regions of China. In this study, the molecular variability and evolution of TYLCV were monitored in Shanghai from its first upsurge in 2006 until 2010. Full-length genomic sequences of 26 isolates were obtained by rolling circle amplification. Sequence analysis showed that the intergenic region was the most variable, with a mean mutation rate of 4.81×10^{-3} nucleotide substitutions per site per year. Genetic differentiation was found within isolates obtained from 2006, 2009, and 2010, though a linear increase in genetic diversity over time was not evident. Whilst significant parts of TYLCV genes were under negative selection, the C4 gene embedded entirely within the C1 gene had a tendency to undergo positive selection. Our results indicate that a mechanism of independent evolution of overlapping regions could apply to the natural population of TYLCV in Shanghai, China.

Key words: Molecular variability, Tomato yellow leaf curl virus (TYLCV), Geminivirus, Evolution

doi:10.1631/jzus.B1300110

Document code: A

CLC number: S432.1

1 Introduction

Geminiviruses are a group of circular, single-stranded DNA viruses that cause devastating diseases worldwide. Four genera (*Mastrevirus*, *Begomovirus*, *Curtovirus*, and *Topocuvirus*) have been taxonomically classified within the family Geminiviridae, with the genus *Begomovirus* being the largest, containing more than 200 species. Earlier studies have demonstrated that mutation, recombination, and pseudo-

recombination are the major contributory factors driving the emergence of begomovirus variants (Czosnek and Ghanim, 2011). Although geminiviruses depend on host DNA polymerase for replication, mutation occurs frequently in geminivirus genomes and progeny populations of high variability can be generated in a short period of time. Mean rates of 1.60×10^{-3} and 1.33×10^{-4} nucleotide substitutions per site per year (subs/site/year) have been estimated for the DNA-A and DNA-B components, respectively, of the East African cassava mosaic virus (EACMV) (Duffy and Holmes, 2009). Similar rates of substitution have also been reported for artificially inoculated hosts (Ge *et al.*, 2007). It is now clear that geminiviruses evolve as rapidly as many RNA viruses, but only a few reports estimate the time required to generate such levels of genetic variability.

[‡] Corresponding author

^{*} Project supported by the Special Fund for Agro-Scientific Research in the Public Interest from the Ministry of Agriculture, China (No. 201003065), the China Postdoctoral Science Foundation (No. 2012M510200), and the National Natural Science Foundation of China (No. 31200117)

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2014

Tomato yellow leaf curl virus (TYLCV) is a monopartite begomovirus belonging to the genus *Begomovirus*, and is transmitted by the whitefly *Bemisia tabaci* in a circulative and persistent manner. The genome of TYLCV encodes two partially overlapping genes in the viral strand: *V1* encodes a coat protein, and *V2* encodes a precoat protein and acts as a suppressor of RNA silencing (Glick *et al.*, 2008; Jiang *et al.*, 2012). The complementary strand of TYLCV encodes four proteins: a replication-associated protein (C1/Rep), a transcriptional activator protein (C2/TrAP), a replication enhancer protein (C3/REn), and protein C4, which is entirely embedded in the *C1* gene, but has a different reading frame. Open reading frames (ORFs) are organized bi-directionally and are separated by an intergenic region (IR) which serves as the origin of viral replication and as a bi-directional promoter (Hanley-Bowdoin *et al.*, 1999).

Tomato yellow leaf curl disease (TYLCD) caused by TYLCV is becoming the most destructive disease in tomato plants (Moriones and Navas-Castillo, 2000). Plants infected by TYLCV during the early growth stages frequently suffer yield losses of up to 100%. Although tomato plants in Middle East countries have been severely affected by TYLCV only since the 1960s, the first report of damage caused by this disease dates back to 1931 in Israel (Cohen and Antignus, 1994). Since then, epidemics have emerged and devastated tomato crops rapidly and widely. TYLCV was first reported in Shanghai, China in 2006 (Wu *et al.*, 2006), and it has since spread rapidly to 13 provinces or autonomous regions of China, including Guangdong, Zhejiang, Jiangsu, Anhui, Shandong, Hebei, Henan, Beijing, Tianjin, Shanxi, Shaanxi, and Inner Mongolia and Xinjiang Uygur Autonomous Regions (Zhang *et al.*, 2009). Since no other begomoviruses had been detected in Shanghai before 2006, the rapid spread of TYLCV provided a great opportunity to understand the evolution of a plant DNA virus soon after its first introduction to a new area. Twenty-six TYLCV isolates were collected from the same area of Shanghai during 2006–2010. Full-length genomic sequencing allows for a more comprehensive analysis of virus variability and of the different forces driving the evolution of TYLCV.

2 Materials and methods

2.1 Collection of TYLCV-infected tomato plants

Twenty-six tomato plants (cultivar Pufen 1, formerly called 98-8) with typical TYLCV symptoms (leaf curling, yellowing, and stunted growth) were collected from Shanghai from 2006 to 2010. TYLCV isolates were named after the year in which they were collected (Table 1).

Table 1 TYLCV isolates collected in Shanghai, China during 2006–2010

Isolate	Collection date	Isolate	Collection date
SH2006-1	2006	SH2009-1	2009
SH2006-2	2006	SH2009-2	2009
SH2007-1	2007	SH2009-3	2009
SH2007-2	2007	SH2009-4	2009
SH2007-3	2007	SH2009-5	2009
SH2007-4	2007	SH2009-6	2009
SH2007-5	2007	SH2010-1	2010
SH2007-6	2007	SH2010-2	2010
SH2008-1	2008	SH2010-3	2010
SH2008-2	2008	SH2010-4	2010
SH2008-3	2008	SH2010-5	2010
SH2008-4	2008	SH2010-6	2010
SH2008-5	2008		
SH2008-6	2008		

2.2 Genomic DNA extraction and detection of TYLCV

Total genomic DNA was extracted from 0.5 g of leaf tissue using the cetyltrimethylammonium bromide (CTAB) method (Zhou *et al.*, 2001). After washing in 70% ethanol, the DNA pellets were dried at room temperature and re-suspended in about 200 μ l of distilled water. Primers TYLCV/F (5'-GAGC TCTTAGCTGCCTGAATGTTC-3') and TYLCV/R (5'-GAGCTCAACAGATGTCAAGACCTAC-3'), which were expected to specifically amplify a 2.8-kb fragment of TYLCV, were used to detect TYLCV infection.

2.3 Isolation, cloning, and sequencing of full-length genomes

Rolling circle amplification (RCA) based on the phi29 DNA polymerase available in the TempliPhi™ kit (GE Healthcare, UK) was used to amplify circular

viral genomes from 1 μg of total plant DNA extractions (Guo *et al.*, 2009). The amplified DNA concatemers (5 μl) were cleaved with *SacI* to generate full-length viral fragments, which were subsequently inserted into linearized pGEM-7Zf+ (Promega, Madison, WI). About five independent clones from each sample were randomly chosen for sequencing by an automated model 3730 DNA sequencer (Perkin-Elmer, USA) using M13 forward and reverse primers and a pair of walking primers, TYLCV-W/F (5'-TCTGCAATCCAGGACCTACC-3') and TYLCV-W/R (5'-AGTCTATCTTGCAATATGTG-3'). Sequences were assembled and edited using EditSeq (Lasergene Package of Programs from DNASTAR Inc.). The fidelity of phi29 DNA polymerase was predetermined by sequencing 20 individual clones generated directly from the TYLCV infectious clone using this enzyme during the initial amplification trials, and the mutation frequency was estimated to be 1.8×10^{-5} .

2.4 Sequence analysis

The circular genomic sequences were arranged to begin at the nick-site in the invariant nonanucleotide sequence at the origin of replication (TAA TATT-3'/5'-AC) (Padidam *et al.*, 1995). Multiple sequence alignment was generated with the aid of ClustalX 1.83 (Thompson *et al.*, 2002). The level of genetic variation was calculated by estimating Watterson's estimator of θ and the average number of nucleotide differences per site between two sequences (Pi) using DnaSP v5 (Guo *et al.*, 2008; Librado and Rozas, 2009). The ratio of pairwise nonsynonymous (d_n) to synonymous (d_s) nucleotide substitutions per site, was estimated using the Pamilo-Bianchi-Li method implemented in MEGA 5 (Tamura *et al.*, 2011). Confidence estimates for nonsynonymous and synonymous nucleotide substitutions were assessed by the bootstrap method based on 100 replicates. The extent of mutational bias and the number of unambiguous nucleotide changes of each type were calculated by MEGA 5 (Tamura *et al.*, 2011). Genetic distances within and between subpopulations were calculated by applying Kimura's two-parameter method in MEGA 5 (Tamura *et al.*, 2011). Evolutionary constraints acting on the overlapping regions were assessed using entropy values and BioEdit software (Hall, 1999). Cumulative synonymous and nonsy-

nonymous mutations were determined using Synonymous Non-synonymous Analysis Program (SNAP) (<http://www.hiv.lanl.gov/content/sequence/SNAP/SNAP.html>) as described by Korber (2002) and Melgarejo *et al.* (2013).

3 Results

3.1 Symptom observation and detection of TYLCV

From 2006 to 2010, a total of 26 samples were obtained from surveys conducted once a year in Shanghai, China. All the plants sampled showed typical TYLCV symptoms, such as leaf curling, yellowing, and stunted growth (Fig. 1a). The specific 2.8-kb DNA fragment was amplified by polymerase chain reaction (PCR) from all the samples using primers TYLCV/F and TYLCV/R (Fig. 1b), indicating that these symptomatic tomato plants were infected by TYLCV. To obtain the complete nucleotide sequence of each of the 26 isolates, RCA followed by *SacI* digestion of the DNA concatemers was performed to obtain the full-length viral fragment (Fig. 1d). About five independent clones from each sample were chosen to be sequenced and were found to be 2781 nucleotides (nts) long with a genomic organization typical of begomoviruses (Fig. 1c).

3.2 Molecular variation of the TYLCV genome

To evaluate the extent of TYLCV variation, the mean pairwise diversity index π and the Watterson parameter θ were calculated at all sites along the TYLCV genome by DnaSP Version 5.0 (Fig. 2). Using a sliding window of 100 nts with a 25-step size, we found that while the overall variability for the TYLCV genome was below 3% at the nucleotide level, distribution patterns were uneven. The peak of variability was located within the intergenic region and 5'-terminal part of V2, whereas the most conserved regions were located in the overlapping regions of V1 and V2. Estimation of the average mutation rate showed that the mutation rate of the full-length TYLCV genome reached 1.69×10^{-3} subs/site/year (Table 2), which is within the ranges for RNA viruses and EACMV (Jenkins *et al.*, 2002; Duffy and Holmes, 2009). Further analysis showed that the substitution rate was even higher for the intergenic region, with a substitution rate of 4.81×10^{-3} subs/site/year. Rates of




Fig. 1 Genome composition and detection of TYLCV-infected virus samples

(a) TYLCV-infected tomato plants showing leaf curling, yellowing, and stunted growth symptoms. (b) PCR detection of TYLCV in six of the infected tomato plants (target size, 2.8 kb). (c) Genomic organization of TYLCV. The intergenic region and six ORFs encoded by the viral strand and complementary strand are shown as indicated. (d) Full-length TYLCV fragment obtained from *SacI* digestion of rolling circle amplification products. IR: intergenic region; M: DNA marker




Fig. 2 Distribution of genetic variation estimated by nucleotide diversity (π) and Watterson's parameter (θ) for TYLCV

The sliding window size is 100 sites wide, slide by 25-site intervals. The relative positions of the six ORFs and the intergenic region of the TYLCV genome are illustrated by lines above the plot. IR: intergenic region

Table 2 Estimates of average evolutionary divergences and nucleotide substitution rates among TYLCV isolates collected from 2006 to 2010

Genomic region	Sequence length (nt)	Number of sequences	Genetic distance*	Nucleotide substitution rate (substitutions per site per year)	
				Mean value	Corrected mean value
Full genome	2781	122	0.007±0.001	1.69×10 ⁻³	1.67×10 ⁻³
IR	313	122	0.023±0.005	4.81×10 ⁻³	4.79×10 ⁻³
V1	777	122	0.005±0.002	9.52×10 ⁻⁴	9.34×10 ⁻⁴
V2	351	122	0.005±0.002	8.50×10 ⁻⁴	8.32×10 ⁻⁴
C1	1074	122	0.005±0.002	1.39×10 ⁻³	1.37×10 ⁻³
C2	408	122	0.009±0.002	2.64×10 ⁻³	2.62×10 ⁻³
C3	405	122	0.008±0.002	1.99×10 ⁻³	1.97×10 ⁻³
C4	294	122	0.005±0.002	1.38×10 ⁻³	1.36×10 ⁻³

* Genetic distance: the numbers of base substitutions per site from averaging over all sequence pairs; data are shown as mean±standard error ($n=100$), obtained by a bootstrap procedure. The analyses were conducted using the Kimura's two-parameter model

nucleotide substitution were also higher in the genes (*C1*, *C2*, *C3*, and *C4*) encoded on the complementary strand than those (*V1* and *V2*) encoded on the virion strand (Table 2). Statistical tests revealed that the substitution rates estimated for the natural population of TYLCV were significantly higher than those generated from RCA error (1.8×10^{-5}). These results indicate that the intergenic region had evolved significantly faster than the other ORFs.

To reveal further the process shaping TYLCV evolution, we tested whether specific kinds of mutational changes were overrepresented in the natural population of TYLCV. Our results showed that the shifts from C to T, T to C, G to A, and G to T were overrepresented relative to other kinds of mutational changes. These kinds of mutations constituted 62% of the mutations. However, the transversions of A to C, T to A, and C to G appeared to be underrepresented, constituting only about 10% of the mutations (Table 3). Therefore, these results indicate that transition biases exist in the evolution of TYLCV. These biases favor a shift from CG to AT genome content, especially with a bias in C to T and G to A transitions.

3.3 Genetic distances between subpopulations of TYLCV

Genetic distance refers to the average number of

Table 3 Percentages of nucleotide substitutions in TYLCV

Original base	Percentage of nucleotide substitution (%)			
	A	T	C	G
A		4.19	3.22	7.96
T	3.59		15.6	7.24
C	4.62	26.1		3.86
G	9.84	10.46	3.33	

Table 4 Estimates of average evolutionary divergence over sequence pairs within groups arranged by sample collection date

Genomic region	Genetic distance within subgroups				
	2006	2007	2008	2009	2010
IR	0.007±0.004	0.003±0.002	0.009±0.002	0.018±0.004	0.017±0.004
<i>V1</i>	0.003±0.002	0.001±0.001	0.002±0.001	0.005±0.002	0.004±0.001
<i>V2</i>	0.002±0.002	0.001±0.000	0.002±0.001	0.004±0.002	0.005±0.002
<i>C1</i>	0.004±0.001	0.001±0.000	0.002±0.000	0.004±0.001	0.004±0.001
<i>C2</i>	0.008±0.003	0.003±0.002	0.004±0.001	0.005±0.002	0.007±0.003
<i>C3</i>	0.006±0.003	0.001±0.001	0.003±0.001	0.006±0.002	0.009±0.003
<i>C4</i>	0.003±0.002	0.002±0.001	0.003±0.001	0.004±0.002	0.004±0.002

nucleotide substitutions between two randomly selected sequences in a population, and can be used to evaluate the genetic variation of a virus population. Using the Kimura's two-parameter method, the mean genetic distances for the intergenic region and individual ORFs ranged from 0.005 to 0.023. Although the individual ORFs of the TYLCV genome presented similar genetic distances, the mean values for the intergenic region showed the highest genetic distances (Table 2). To understand better the genetic variation of the 26 isolates, the full-length genomic sequences were divided into several subgroups according to the year of collection. Analysis showed that the highest genetic distance was found for the intergenic region (IR) within subgroups (Table 4). Although we did not observe a quasi-linear increase in genetic diversity across years, calculation of the F_{st} , the coefficient used to evaluate the extent of genetic differentiation or the gene flow between two populations, showed that the values of F_{st} among subgroups 2006 and 2009, 2006 and 2010, as well as 2009 and 2010 were all <0.33, suggesting frequent gene flow (Table 5).

Table 5 F_{st} values for pairs of temporal subpopulations of TYLCV

Year	F_{st}			
	2006	2007	2008	2009
2007	0.701 18			
2008	0.630 97	0.373 27		
2009	0.167 67	0.606 62	0.548 62	
2010	0.296 72	0.619 52	0.569 20	0.133 70

F_{st} is a coefficient of the extent of genetic differentiation and provides an estimate of the extent of gene flow. An absolute value of $F_{st} > 0.33$ suggests infrequent gene flow

3.4 Selection constraints acting on TYLCV genes

Pairwise genetic differences at the synonymous (d_s) and nonsynonymous (d_n) nucleotide positions were estimated according to the Pamilo-Bianchi-Li method. The ratio between nucleotide diversity values in nonsynonymous and synonymous positions (d_n/d_s) (Table 6) provides an estimation of the degree and direction of the selective constraints acting on the coding regions of TYLCV. Overall, the values of the d_n/d_s ratio for the *V1*, *V2*, *C1*, and *C2* genes were markedly low (0.154–0.44), indicating that these genes are under negative selection. However, the ratios of d_n/d_s for the *C3* and *C4* genes were greater than 1. This was probably due to the overlapping regions of *C3-C2* and *C4-C1*, respectively. In the overlapping, frame-shifted *C1/C4* region of TYLCV, the first nucleotide position in the *C1* codons corresponds to the third position in the *C4* codons (*C1-1/C4-3*), the second position in the *C1* codons corresponds to the first position in the *C4* codons (*C1-2/C4-1*), and the third position in the *C1* codons corresponds to the second position in the *C4* codons (*C1-3/C4-2*) (Fig. 3a). Independent evolution of overlapping genes has been described for several viruses, with one of the two overlapping genes being subjected to negative selection, while the other is subjected to positive selection. To evaluate the variation between *C1* and *C4*, entropy values for individual amino acids and nucleotides were determined using the BioEdit software. The amino acid changes in the Rep protein were largely determined by nucleotide mutations at the *C1-1/C4-3* sites, whereas those in the *C4* protein were primarily caused by mutations at the *C1-3/C4-2* sites (Fig. 3b). Furthermore, mutations at the *C1-2/C4-1* sites, which would affect the amino acid sequences of both the Rep and *C4* proteins, were rare. Cumulative synonymous and nonsynonymous mutations plotted using SNAP showed that the *C1* gene accumulated more synonymous mutations, while the *C4* gene accumulated more nonsynonymous mutations (Fig. 3c). Taken together, these results revealed that the overlapping regions of *C1* and *C4* are under evolutionary constraints, allowing for more rapid evolution of the *C4* protein while minimizing amino acid changes in the conserved Rep protein.

Table 6 Estimation of nucleotide diversity of different genes in TYLCV

Gene	d_n	d_s	d_n/d_s
<i>V1</i>	0.002±0.001	0.013±0.004	0.154
<i>V2</i>	0.002±0.002	0.010±0.005	0.200
<i>C1</i>	0.004±0.001	0.009±0.004	0.440
<i>C2</i>	0.006±0.003	0.023±0.013	0.260
<i>C3</i>	0.009±0.004	0.004±0.004	2.250
<i>C4</i>	0.008±0.003	0.002±0.001	4.000

d_n : nonsynonymous substitution; d_s : synonymous substitution. d_n and d_s are expressed as mean±standard error

4 Discussion

Before 2006, no begomovirus or TYLCV had been detected in Shanghai. Due to the rapid introduction and severe outbreak of the Q biotype of whitefly in China, TYLCV has now moved to several provinces of China, causing unprecedented economic losses, especially in tomato plants. Prior to this field survey, no other competitive begomovirus had been described in Shanghai. This epidemiological situation provides a great opportunity to investigate the genetic structure and evolution of TYLCV following its introduction into a new region.

This study analyzed the genetic variability of 26 TYLCV isolates collected from Shanghai over a five-year period. Analysis of the full-length genomic DNA sequences revealed that the level of genetic variation observed in the natural population of TYLCV (1.69×10^{-3} subs/site/year) was similar to that found in RNA viruses (Duffy *et al.*, 2008; Simmons *et al.*, 2008; Pagan and Holmes, 2010; Pagan *et al.*, 2010). These and other results suggest that a considerable amount of sequence variation occurs within and between populations of geminiviruses, challenging the view that RNA viruses are prone to higher mutation rates than DNA viruses (Domingo and Holland, 1997; Roossinck, 1997; Isnard *et al.*, 1998; Sanz *et al.*, 1999). It has been speculated that the high mutation rate of geminivirus genomes is in part due to improper methylation patterns occurring during their replication (Ge *et al.*, 2007; Duffy and Holmes, 2008). However, recent studies have shown that geminivirus




Fig. 3 Variation in nucleotides and amino acids of the overlapping *C1* and *C4* genes

(a) Frame-shifted positions of the overlapping *C1* and *C4* genes. Mutations in the open reading frames of the overlapping *C1* and *C4* genes result in differential rates of amino acid changes. For example, a nucleotide mutation in the third position of the *C1* codon is likely to be synonymous, allowing a nonsynonymous mutation in the *C4* codon (arrows). (b) Levels of nucleotide and amino acid variation in the three sets of nucleotides in relation to the nucleotide positions of the codons in the overlapping Rep and *C4* proteins. (c) Cumulative incidences of synonymous and nonsynonymous mutations in the overlapping region of *C1/C4*. The *x* axis represents the position of the codon in *C4*, and the *y* axis represents the cumulative values of synonymous or nonsynonymous mutations estimated at a specific codon position

genomes are subjected to DNA methylation in infected host plants (Raja *et al.*, 2008; 2010; Yang *et al.*, 2011; Zhang *et al.*, 2011). Thus, it is possible that base-excision repair may not act on geminivirus genomes because the double-stranded state lasts for a very short time during rolling circle replication. The high mutation rate is also probably in part due to base deamination occurring spontaneously or by the action of deaminating host enzymes (Duffy and Holmes, 2008; van der Walt *et al.*, 2008). This may be supported by the observation of C to T transition

biases in this study. Indeed, previous studies have also revealed the same transition biases as our genomic analysis (Ge *et al.*, 2007; Duffy and Holmes, 2008; 2009).

Although we did not observe a significant quasi-linear increase in genetic diversity across years, we found frequent gene flow between isolates obtained from the years 2006, 2009, and 2010. As was reported for TYLCV surveyed in an insular environment, the amount of mutations analyzed in this study was greater than that found in tomato yellow leaf curl

Sardinia virus (TYLCSV) over an eight-year period, using a single-strand conformation polymorphism technique (Sanchez-Campos *et al.*, 2002; Delatte *et al.*, 2007). This discrepancy might be explained by the different techniques used. The RCA technique used in this study has revolutionized the diagnosis and genomics of geminiviruses, allowing for a more efficient and reliable way to obtain full-length geminivirus DNA (Haible *et al.*, 2006). Our results also indicated that the mutation rate detected in the natural TYLCV population was greater than that caused by phi29 DNA polymerase and sequencing errors. Alternatively, this discrepancy might also come from the viruses used. It seems that TYLCV causes more severe damage to crop production, as surveys conducted between 1996 and 1998 in Málaga showed a progressive displacement of TYLCSV by TYLCV (Sanchez-Campos *et al.*, 1999). Indeed, since the first introduction of TYLCV into China, it has become the most prevalent begomovirus in China.

Estimation of nucleotide diversities using the ratio between nonsynonymous and synonymous positions (d_n/d_s) suggested that V1, V2, C1, and C2 were more tolerant to amino acid changes than C3 and C4. Among all the coding regions analyzed, the C4 gene embedded in the C1 gene appeared to be the most flexible and might be subjected to positive selection. We demonstrated that the evolution of the overlapping C1 and C4 genes occurs primarily via C1-1/C4-3 and C1-3/C4-2 sites, respectively, and mutations at the C1-2/C4-1 sites, which were most likely to alter the amino acid sequences of both the Rep and C1 proteins, were rare. Further SNAP analysis showed that positions in the C-terminal of the C4 gene, which were under positive selection, were subjected to purifying selection in the overlapping region of the C1 gene. Indeed, a similar phenomenon has been demonstrated for the evolution of tomato leaf deformation virus, an indigenous new world (NW) monopartite begomovirus emerging from the DNA-A component of an NW bipartite progenitor (Melgarejo *et al.*, 2013). We speculated that this would be correlated with the function of the C4 protein. Compared to the conserved Rep protein, the functions of the C4 protein are more divergent among monopartite and bipartite begomoviruses. For monopartite begomoviruses, C4 is a

pathogenicity determinant that is involved in movement and suppression of host defense responses, whereas the homologues of C4 encoded by most NW begomoviruses do not play a role in pathogenicity. This strategy, which allows viruses to condense a maximal amount of information into short genomes, has been widely used by a number of animal and plant-infecting viruses (Zaaijer *et al.*, 2007). Given the rapid spread of TYLCV in China, it will be interesting to monitor the evolutionary constraints *in vivo* in experimentally infected tomato plants.

Compliance with ethics guidelines

Xiu-ling YANG, Meng-ning ZHOU, Ya-juan QIAN, Yan XIE, and Xue-ping ZHOU declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Cohen, S., Antignus, Y., 1994. Tomato yellow leaf curl virus, a whitefly-borne geminivirus of tomatoes. *In: Harris, K.F. (Ed.), Advances in Disease Vector Research*. Springer-Verlag, New York, p.259-288. [doi:10.1007/978-1-4612-2590-4_10]
- Czosnek, H., Ghanim, M., 2011. *Bemisia tabaci*—tomato yellow leaf curl virus interaction causing worldwide epidemics. *In: Thompson, W.M.O. (Ed.), The Whitefly, Bemisia Tabaci (Homoptera: Aleyrodidae) Interaction with Geminivirus-Infected Host Plants*. Springer Netherlands, p.51-67. [doi:10.1007/978-94-007-1524-0_3]
- Delatte, H., Holota, H., Moury, B., *et al.*, 2007. Evidence for a founder effect after introduction of tomato yellow leaf curl virus-mild in an insular environment. *J. Mol. Evol.*, **65**(1):112-118. [doi:10.1007/s00239-007-0005-x]
- Domingo, E., Holland, J.J., 1997. RNA virus mutations and fitness for survival. *Annu. Rev. Microbiol.*, **51**(1):151-178. [doi:10.1146/annurev.micro.51.1.151]
- Duffy, S., Holmes, E.C., 2008. Phylogenetic evidence for rapid rates of molecular evolution in the single-stranded DNA begomovirus tomato yellow leaf curl virus. *J. Virol.*, **82**(2):957-965. [doi:10.1128/JVI.01929-07]
- Duffy, S., Holmes, E.C., 2009. Validation of high rates of nucleotide substitution in geminiviruses: phylogenetic evidence from East African cassava mosaic viruses. *J. Gen. Virol.*, **90**(6):1539-1547. [doi:10.1099/vir.0.009266-0]
- Duffy, S., Shackelton, L.A., Holmes, E.C., 2008. Rates of evolutionary change in viruses: patterns and determinants. *Nat. Rev. Genet.*, **9**(4):267-276. [doi:10.1038/nrg2323]

- Ge, L., Zhang, J., Zhou, X., et al., 2007. Genetic structure and population variability of tomato yellow leaf curl China virus. *J. Virol.*, **81**(11):5902-5907. [doi:10.1128/JVI.02431-06]
- Glick, E., Zrachya, A., Levy, Y., et al., 2008. Interaction with host SGS3 is required for suppression of RNA silencing by tomato yellow leaf curl virus V2 protein. *PNAS*, **105**(1):157-161. [doi:10.1073/pnas.0709036105]
- Guo, W., Jiang, T., Zhang, X., et al., 2008. Molecular variation of satellite DNA β molecules associated with *Malvastrum* yellow vein virus and their role in pathogenicity. *Appl. Environ. Microbiol.*, **74**(6):1909-1913. [doi:10.1128/AEM.02461-07]
- Guo, W., Yang, X., Xie, Y., et al., 2009. Tomato yellow leaf curl Thailand virus-[Y72] from Yunnan is a monopartite begomovirus associated with DNA β . *Virus Genes*, **38**(2):328-333. [doi:10.1007/s11262-009-0327-4]
- Haible, D., Kober, S., Jeske, H., 2006. Rolling circle amplification revolutionizes diagnosis and genomics of geminiviruses. *J. Virol. Methods*, **135**(1):9-16. [doi:10.1016/j.jviromet.2006.01.017]
- Hall, T., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids Symp. Ser.*, **41**:95-98.
- Hanley-Bowdoin, L., Settledge, S.B., Orozco, B.M., et al., 1999. Geminiviruses: models for plant DNA replication, transcription, and cell cycle regulation. *Crit. Rev. Biochem. Mol. Biol.*, **18**(1):71-106. [doi:10.1080/07352689991309162]
- Isnard, M., Granier, M., Frutos, R., et al., 1998. Quasispecies nature of three maize streak virus isolates obtained through different modes of selection from a population used to assess response to infection of maize cultivars. *J. Gen. Virol.*, **79**(12):3091-3099.
- Jenkins, G.M., Rambaut, A., Pybus, O.G., et al., 2002. Rates of molecular evolution in RNA viruses: a quantitative phylogenetic analysis. *J. Mol. Evol.*, **54**(2):156-165. [doi:10.1007/s00239-001-0064-3]
- Jiang, L., Wei, C.H., Li, Y., 2012. Viral suppression of RNA silencing. *Sci. China C Life Sci.*, **55**(2):109-118. [doi:10.1007/s11427-012-4279-x]
- Korber, B., 2002. HIV signature and sequence similarities. In: Rodrigo, A.G., Learn, J.R., Gerald, H. (Eds.), *Computational and Evolutionary Analysis of HIV Molecular Sequences*. Kluwer Academic Publishers, Boston, Massachusetts, p.55-72.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, **25**(11):1451-1452. [doi:10.1093/bioinformatics/btp187]
- Melgarejo, T.A., Kon, T., Rojas, M.R., et al., 2013. Characterization of a new world monopartite begomovirus causing leaf curl disease of tomato in Ecuador and Peru reveals a new direction in geminivirus evolution. *J. Virol.*, **87**(10):5397-5413. [doi:10.1128/JVI.00234-13]
- Moriones, E., Navas-Castillo, J., 2000. Tomato yellow leaf curl virus, an emerging virus complex causing epidemics worldwide. *Virus Res.*, **71**(1-2):123-134. [doi:10.1616/S0168-1702(00)00193-3]
- Padidam, M., Beachy, R.N., Fauquet, C.M., 1995. Classification and identification of geminiviruses using sequence comparisons. *J. Gen. Virol.*, **76**(Pt 2):249-263. [doi:10.1099/0022-1317-76-2-249]
- Pagan, I., Holmes, E.C., 2010. Long-term evolution of the *Luteoviridae*: time scale and mode of virus speciation. *J. Virol.*, **84**(12):6177-6187. [doi:10.1128/JVI.02160-09]
- Pagan, I., Firth, C., Holmes, E.C., 2010. Phylogenetic analysis reveals rapid evolutionary dynamics in the plant RNA virus genus tobamovirus. *J. Mol. Evol.*, **71**(4):298-307. [doi:10.1007/s00239-010-9385-4]
- Raja, P., Sanville, B.C., Buchmann, R.C., et al., 2008. Viral genome methylation as an epigenetic defense against geminiviruses. *J. Virol.*, **82**(18):8997-9007. [doi:10.1128/JVI.00719-08]
- Raja, P., Wolf, J.N., Bisaro, D.M., 2010. RNA silencing directed against geminiviruses: post-transcriptional and epigenetic components. *Biochim. Biophys. Acta*, **1799**(3):337-351. [doi:10.1016/j.bbagr.2010.01.004]
- Roossinck, M.J., 1997. Mechanisms of plant virus evolution. *Annu. Rev. Phytopathol.*, **35**(1):191-209. [doi:10.1146/annurev.phyto.35.1.191]
- Sanchez-Campos, S., Navas-Castillo, J., Camero, R., et al., 1999. Displacement of tomato yellow leaf curl virus (TYLCV)-Sr by TYLCV-Is in tomato epidemics in Spain. *Phytopathology*, **89**(11):1038-1043. [doi:10.1094/PHYTO.1999.89.11.1038]
- Sanchez-Campos, S., Diaz, J.A., Monci, F., et al., 2002. High genetic stability of the *Begomovirus* tomato yellow leaf curl *Sardinia* virus in southern Spain over an 8-year period. *Phytopathology*, **92**(8):842-849. [doi:10.1094/PHYTO.2002.92.8.842]
- Sanz, A.I., Fraile, A., Gallego, J.M., et al., 1999. Genetic variability of natural populations of cotton leaf curl geminivirus, a single-stranded DNA virus. *J. Mol. Evol.*, **49**(5):672-681. [doi:10.1007/PL00006588]
- Simmons, H.E., Holmes, E.C., Stephenson, A.G., 2008. Rapid evolutionary dynamics of zucchini yellow mosaic virus. *J. Gen. Virol.*, **89**(4):1081-1085. [doi:10.1099/vir.0.83543-0]
- Tamura, K., Peterson, D., Peterson, N., et al., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.*, **28**(10):2731-2739. [doi:10.1093/molbev/msr121]
- Thompson, J.D., Gibson, T.J., Higgins, D.G., 2002. Multiple sequence alignment using ClustalW and ClustalX. *Curr. Prot. Bioinf.*, **00**:2.3.1-2.3.22. [doi:10.1002/0471250953.bi0203s00]
- van der Walt, E., Martin, D.P., Varsani, A., et al., 2008. Experimental observations of rapid Maize streak virus evolution reveal a strand-specific nucleotide substitution bias. *Virol. J.*, **5**:104. [doi:10.1186/1743-422X-5-104]

- Wu, J.B., Dai, F.M., Zhou, X.P., 2006. First report of tomato yellow leaf curl virus in China. *Plant Dis.*, **90**(10): 1359-1359. [doi:10.1094/Pd-90-1359c]
- Yang, X., Xie, Y., Raja, P., et al., 2011. Suppression of methylation-mediated transcriptional gene silencing by β C1-SAHH protein interaction during geminivirus-betasatellite infection. *PLoS Pathog.*, **7**(10):e1002329. [doi:10.1371/journal.ppat.1002329]
- Zaaijer, H.L., van Hemert, F.J., Koppelman, M.H., et al., 2007. Independent evolution of overlapping polymerase and surface protein genes of hepatitis B virus. *J. Gen. Virol.*, **88**(Pt 8):2137-2143. [doi:10.1099/vir.0.82906-0]
- Zhang, H., Gong, H., Zhou, X., 2009. Molecular characterization and pathogenicity of tomato yellow leaf curl virus in China. *Virus Genes*, **39**(2):249-255. [doi:10.1007/s11262-009-0384-8]
- Zhang, Z., Chen, H., Huang, X., et al., 2011. BSCTV C2 attenuates the degradation of SAMDC1 to suppress DNA methylation-mediated gene silencing in *Arabidopsis*. *Plant Cell*, **23**(1):273-288. [doi:10.1105/tpc.110.081695]
- Zhou, X.P., Xie, Y., Zhang, Z.K., 2001. Molecular characterization of a distinct begomovirus infecting tobacco in Yunnan, China. *Arch. Virol.*, **146**(8):1599-1606. [doi:10.1007/s007050170081]

中文概要:

本文题目: 番茄黄化曲叶病毒自然种群的分子变异和进化研究

Molecular variability and evolution of a natural population of tomato yellow leaf curl virus in Shanghai, China

研究目的: 2006年我国上海首次发现番茄黄化曲叶病毒(TYLCV), 随后TYLCV迅速蔓延至全国13个省份和自治区。本研究分析了2006至2010年期间TYLCV在我国首发地上海市的分子变异规律。

创新要点: 本研究持续五年追踪田间TYLCV, 分析TYLCV的全长基因组序列、分子变异及种群遗传结构, 为防控TYLCV提供理论依据。

研究方法: 2006至2010年从上海采集了26个TYLCV的分离物, 利用高保真性的滚环扩增技术获得TYLCV分离物的全长基因组。应用MEGA5等生物信息学软件分析TYLCV的分子变异。

重要结论: TYLCV自然种群具有与RNA病毒相似的突变率, 以基因间隔区的分子变异最大, 平均突变率为 4.81×10^{-3} (见图2和表2)。TYLCV的大部分基因都处于负向选择, 但包含在C1开放阅读框内的C4, 却与C1承受着不同的选择压而处于正向选择(见图3和表6)。

关键词组: 分子变异; 番茄黄化曲叶病毒(TYLCV); 双生病毒; 进化