



Analysis of promoters of microRNAs from a *Glycine max* degradome library^{*#}

Yi-qiang HAN¹, Zheng HU², Dian-feng ZHENG³, Ya-mei GAO^{†‡1}

⁽¹⁾College of Life Science and Technology, Heilongjiang Bayi Agricultural University, Daqing 163319, China)

⁽²⁾The National Key Facility for Crop Gene Resources and Genetic Improvement, Institute of Crop Science,
 Chinese Academy of Agricultural Sciences, Beijing 100081, China)

⁽³⁾College of Agronomy, Heilongjiang Bayi Agricultural University, Daqing 163319, China)

[†]E-mail: gaoym800@126.com

Received July 1, 2013; Revision accepted Aug. 25, 2013; Crosschecked Jan. 14, 2014

Abstract: Objective: MicroRNAs (miRNAs) are genome-encoded, small non-coding RNAs that play important functions in development, biotic and abiotic stress responses, and other processes. Our aim was to explore the regulation of miRNA expression. Methods: We used bioinformatics methods to predict the core promoters of 440 miRNAs identified from a soybean (*Glycine max*) degradome library and to analyze *cis*-acting elements for 369 miRNAs. Results: The prediction results showed that 83.86% of the 440 miRNAs contained promoters in their upstream sequences, and 8.64% (38 loci) in their downstream sequences. The distributions of two core promoter elements, TATA-boxes and transcription start sites (TSSs), were similar. The *cis*-acting elements were examined to provide clues to the function and regulation of spatiotemporal expression of the miRNAs. Analyses of miRNA *cis*-elements and targets indicated a potential auxin response factor (ARF)- and gibberellin response factor (GARF)-mediated negative feedback loop for miRNA expression. Conclusions: The features of miRNAs from a *Glycine max* degradome library obtained here provide insights into the transcription regulation and functions of miRNAs in soybean.

Key words: *Glycine max*, MicroRNA (miRNA), Promoter, *Cis*-acting element, Prediction

doi:10.1631/jzus.B1300179

Document code: A

CLC number: Q522⁺.6

1 Introduction

MicroRNAs (miRNAs) are genome-encoded, about 21–23-nucleotide non-coding RNAs that negatively regulate gene expression by directing mRNA cleavage or interfering with translation (Carrington and Ambros, 2003). Studies have shown that miRNAs have the potential to regulate all physiological cell

functions (Llave *et al.*, 2002). In plants, miRNAs regulate the expression of a large number of genes in diverse biological processes (Chen, 2004; Lu *et al.*, 2005; Zeng *et al.*, 2010; Shamimuzzaman and Vodkin, 2012; Zhang and Li, 2013), as identified by computational prediction, microarray analysis, and high-throughput sequencing (Jones-Rhoades and Bartel, 2004; Liu *et al.*, 2008; Zhao *et al.*, 2010; Zhu and Luo, 2013). Most miRNAs exist as independent transcription units, which are transcribed into long primary transcripts (pri-miRNAs) by RNA polymerase II, then cleaved to miRNA precursors (pre-miRNAs) (Bartel, 2004; Lee *et al.*, 2004). miRNAs have broad regulatory effects throughout the plant. Therefore, understanding the mechanisms regulating miRNA gene expression remains an important fundamental goal.

[‡] Corresponding author

^{*} Project supported by the National Natural Science Foundation of China (No. 31271652) and the National Science Foundation for Young Scientists of China (No. 31301708)

[#] Electronic supplementary materials: The online version of this article (<http://dx.doi.org/10.1631/jzus.B1300179>) contains supplementary materials, which are available to authorized users

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2014

Class II promoters contain two parts, a core promoter and upstream elements. The core promoter consists of at least two elements: a TATA box, which begins at about the -30 bp position, and an initiator, which is centered on the transcription start site (TSS) (Liu *et al.*, 2010). Promoter elements of pre-miRNAs are similar to those of protein-coding genes (Smale, 2001). Bioinformatics analysis of the core miRNA promoters in model plants, including *Arabidopsis thaliana* and *Oryza sativa*, confirmed that the promoters of most known miRNA genes are of the same type as those of protein-coding genes and are transcribed by RNA polymerase II (Xie *et al.*, 2005; Zhou *et al.*, 2007; Cui *et al.*, 2009). TATA-less promoters were also identified in *Arabidopsis* and rice. The promoters of most miRNA genes are found within 500-bp upstream of the TSS. Although substantial work has examined the promoters of protein-coding genes, the nature of miRNA promoter elements is still one of the most interesting aspects of small RNA biology (Megraw *et al.*, 2006).

Soybean (*Glycine max*) is one of the most economically important agricultural crops in the world, and one of the most important global sources of protein and oil, both for food and for livestock feed. Soybean was the first legume species with a published high-quality draft genome sequence (Schmutz *et al.*, 2010). Because miRNAs play an important regulatory role in a wide variety of developmental and metabolic processes in plants, more and more soybean miRNAs have been identified (Zhang *et al.*, 2008; Chen *et al.*, 2009; Wang *et al.*, 2009; Guo *et al.*, 2011; Kulcheski *et al.*, 2011; Li *et al.*, 2011; Song *et al.*, 2011; Wong *et al.*, 2011; Zeng *et al.*, 2012; Hu *et al.*, 2013), but the features of the upstream sequences of these miRNAs have not been comprehensively analyzed. Core promoters and *cis*-acting elements of 82 soybean miRNAs from the miRBase database were analyzed by bioinformatics methods (Liu *et al.*, 2010), but this analysis involved a relatively small number of miRNA genes. We have identified substantial numbers of miRNAs by deep sequencing of a *Glycine max* degradome library (Hu *et al.*, 2013). In this study, the core promoters and *cis*-acting elements of those miRNAs were predicted by bioinformatics methods to reveal the features of the core promoters of soybean miRNAs. The miRNA expression regulation pathway was also analyzed. These studies provide insights into the regulation of soybean miRNA expression.

2 Materials and methods

2.1 MicroRNA data and classification

A total of 440 miRNAs were identified from a soybean cDNA degradome library that was constructed from root, stem, leaf, and inflorescence tissues. The National Center for Biotechnology Information-Gene Expression Omnibus (NCBI-GEO) accession number of the *G. max* degradome sequences is GSE33380. All soybean miRNA sequences for prediction were divided into two groups based on their genomic background: “intronic miRNAs”, which overlap protein-coding genes, and “intergenic miRNAs”, which reside between protein-coding genes. The protein-coding genes included verified and hypothetical genes. The 440 soybean miRNAs included 34 intronic miRNAs and 406 intergenic miRNAs.

2.2 Identification of potential core promoters of soybean miRNAs

For promoter prediction, the upstream and downstream sequences of the 440 miRNA genes were downloaded from the soybean genome database (<http://www.phytozome.net/index.php>) (Goodstein *et al.*, 2012). The upstream sequences of pre-microRNAs (hairpin precursors) of each miRNA gene were obtained according to Cui *et al.* (2009). The 200-bp downstream sequence of each miRNA gene was also used for promoter prediction. For 83.86% of the pre-microRNAs (hairpin precursors), the upstream sequence was 2000 bp or longer. Promoters were predicted by the plant promoter identification program TSSP (<http://www.softberry.com>), which is designed for predicting plant Pol II promoters (Shahmuradov *et al.*, 2005). The predictions were obtained at the default TSSP settings.

2.3 Analysis of *cis*-acting elements of soybean miRNAs

Potential promoter regions (from TSS to 800 bp upstream) were obtained to predict the potential *cis*-acting elements and motifs. If there were overlapping TSSs in the selected region, only the sequence between the two TSSs was analyzed to exclude redundancy (Liu *et al.*, 2010). The PlantCARE database (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html>), a database of plant *cis*-acting regulatory elements (Lescot *et al.*, 2002), was used to analyze the *cis*-acting elements of the miRNAs.

3 Results

3.1 Core promoters and enhancers of soybean miRNAs

Putative promoters were divided into TATA-box/TSS core promoters and enhancers, and counted independently. From 440 miRNAs, we identified 699 predicted core promoters, with 102 predicted enhancers in upstream sequences. There were 38 core promoters, but no enhancers, predicted in the downstream sequence.

Among the upstream sequences of the 440 miRNAs, 71 (16.14%) were found to contain no predicted promoters (Fig. 1a). Of the other 369 (83.86%) promoter-containing sequences, 127 (28.86%) were predicted to have only one promoter, 168 (38.18%) contained two, and the rest, 74 (16.90%), contained three or more predicted promoters. Among the predicted enhancers in upstream sequences of the 440 miRNAs, 345 (78.41%) were found to contain no enhancers (Fig. 1b). Among the other 95 (21.59%) enhancer-containing sequences, 88 (20.00%) were predicted to have only one enhancer, 7 (1.59%) contained two enhancers, and no sequences were predicted to contain three or more enhancers. Most pre-miRNAs examined contained no predicted enhancers.

In the downstream sequences of the 440 miRNAs, 402 (91.36%) were found to contain no predicted promoter (Fig. 1c) and 38 (8.64%) were predicted to have only one. None of the 440 miRNAs contained predicted enhancers in the downstream sequence (Fig. 1d).

Pre-miRNAs from intergenic regions generally had more core promoters than the intronic pre-miRNAs (Table S1), and two or fewer core promoters were identified for each intronic miRNA. In the pre-miRNA downstream sequences, there were few core promoters, and never more than one.

3.2 Distribution of the predicted core promoters and enhancers of soybean miRNA genes

We analyzed the specific sequences of pre-miRNAs using the TSSP program, and counted the TSS, TATA-box, and enhancers separately. For the predicted core promoters in the upstream sequences of the 440 miRNAs, a total of 699 TSSs were identified. A strong peak was observed near the start position of each transcript. For intergenic miRNAs, the distribution could be separated into three distinct regions (Fig. 2a). The first peak was found close to the pre-miRNA within the 1.0-kb upstream region, which contained 56.41% of the total predicted TSSs (663).




Fig. 1 Distribution of numbers of promoters in upstream (a) and downstream (c) sequences of miRNAs and distribution of numbers of enhancers in upstream (b) and downstream (d) sequences of miRNAs




Fig. 2 Genomic distributions of TSSs (a), TATA-boxes (b), and enhancers (c) from the 5' end of the pre-miRNA upstream sequence

A second broad region, from -1.0 to -1.6 kb, contained 27.60% of the predicted TSSs. The third region, from -1.6 to -2.0 kb, contained 15.99% of the predicted TSSs. For intronic miRNAs, there were also three peaks of TSSs (from 0 to -1.2 kb, -1.2 to -1.6 kb, and -1.6 to -2.0 kb).

The distribution of TATA-boxes was similar to the distribution of TSSs. The core promoters of four miRNAs (gm-cmiR160f, gm-cmiR319k, gm-cmiR13, and gm-cmiR4i) were found to be TATA-less. For intergenic miRNAs, we identified three distinct regions (Fig. 2b). The first peak was found close to the

pre-miRNA within the 0.6-kb upstream region, and contained 36.42% of the total (659) predicted TATA-boxes. A second broad region, from -0.6 to -1.2 kb, contained 28.68% of predicted TATA-boxes. The third region, from -1.1 to -2.0 kb, contained 34.90%. For intronic miRNAs, there were also three peaks of TATA-boxes (from 0 to -0.6 kb, -0.6 to -1.2 kb, and -1.2 to -2.0 kb).

In intergenic and intronic miRNAs, 102 enhancers were predicted. For intergenic miRNAs, most (74.51%) of the predicted enhancers were found within 1.0-kb upstream of pre-miRNAs. The remaining 15.69% occurred within -1.0 to -1.6 kb. The third region, from -1.6 to -2.0 kb, contained 9.80% of the predicted enhancers (Fig. 2c). For the predicted core promoters in downstream sequences of the 440 miRNAs, a total of 36 (8.18%) TSSs and TATA-boxes were identified.

3.3 *Cis*-acting elements in potential promoters and an miRNA expression regulation pathway related to hormone metabolism

To further elucidate the spatiotemporal expression pattern and the function of miRNAs from the *G. max* degradome library, 800 bp upstream sequences of the potential TSS of 369 miRNA genes were analyzed using the PlantCARE database, excluding 71 miRNAs that had no predicted TSS. *Cis*-acting elements and motifs of 355 miRNA promoters were predicted, excluding 14 miRNAs without sufficient sequence information. There were 126 types of *cis*-acting elements. Every miRNA contained a TATA-box core promoter element at around -30 bp from the TSS and a CAAT-box, a common *cis*-acting element, in the promoter and enhancer regions (Table S2). In addition, 99.72% of the miRNAs contained a light responsive element (LRE). *Cis*-acting elements were grouped into several classes according to their functions, such as stress-responsive elements (SREs), *cis*-acting elements regulating plant growth (GREs), hormone-regulated *cis*-acting elements (HREs), and tissue-specific *cis*-acting elements (TSEs) (Table S3) as described by Liu *et al.* (2010). Other regulatory elements, such as AT-rich DNA, *cis*-acting regulatory elements essential for anaerobic induction, and elicitor-responsive elements, were not classified. Of the miRNA sequences examined, 86.48% contained an SRE, including anaerobic induction elements (AREs), defense SREs

(TC-rich repeats), ABA (abscisic acid)-response elements (ABREs), low-temperature responsive elements (LTRs), MYB (v-myb avian myeloblastosis viral oncogene homolog) binding sites involved in drought response (MBSs), salicylic acid responsiveness elements (SAEs), heat-stress responsive elements (HSEs), and LREs. Also, 35.49% of miRNAs contained a GRE, including a leaf morphology development element (HD-Zip 2), differentiation of the palisade mesophyll cells element (HD-Zip 1), meristem-specific activation elements (OCT, NON-box, dOCT, CCGTCC-box, CAT-box), zein metabolism regulation element (O2-site), circadian control element (circadian), and cell cycle regulation elements (MSA-like). TSEs were identified in 72.39% of miRNA sequences examined, including nodule-specific factor binding site (Nodule-site2), involved in the endosperm-specific negative expression element (AACA_motif), endosperm expression elements (GCN4_motif, Skn-1_motif), and *cis*-acting regulatory elements involved in root-specific expression (as1) and seed-specific regulation (RY-element). HREs were found in 70.70% of miRNA sequences examined, including those possibly involved in the regulation of the response to plant hormones, such as gibberellin (GA), auxin-responsive element, ethylene, salicylic acid, methyl jasmonate, and flavonoids.

Among HREs, numerous copies of the auxin response factor (ARF) elements are found in the putative promoter regions of *MIR160d*, *MIR160f*, *MIR167a*, and *MIR167g*. Numerous copies of gibberellin response factor (GARF) elements are more frequently found in the putative promoter regions of *MIR160f*, *MIR160g*, *MIR160h*, *MIR167a*, *MIR167c*, *MIR167d*, and *MIR167i*. In our miRNA target gene data, multiple ARF family members, including *ARF17*, *ARF18*, *ARF6*, and *ARF8*, were predicted to be directly targeted by miR160 and miR167. Based on the above data, we propose a possible ARF- and GARF-mediated negative feedback regulatory loop involving miR160, miR167, and certain ARF family members (Fig. 3). *ARF17* and *ARF18* are post-transcriptionally regulated by miR160, and *ARF6* and *ARF8* are regulated by miR167 (Fig. 3a). In the negative feedback regulatory loop, miR160 and miR167 repress their target *ARF*, and ARFs affect miR160 and miR167 accumulation via binding to the AR regulation elements of the *MIR160* and *MIR167* promoters.

In addition to the feedback regulation of miR160 and miR167 by ARF, GARF can regulate miR160 and miR167 via binding to the GARF elements of the *MIR160* and *MIR167* promoters (Fig. 3b). According to the feedback regulation of miR160 and miR167 by ARF and GARF, we speculate that there is GA-auxin cross-talk in the pathway of the regulation of miRNA expression.




Fig. 3 An speculated miRNA expression regulation pathway related to hormone metabolism

(a) Certain auxin response factor (ARF) family members are directly regulated by miRNAs (miR160 and miR167). (b) A possible negative feedback regulatory loop involving miR160, miR167, and certain ARF family members. Numerous copies of the ARF recognition motif are found in the putative promoter regions of *MIR160d*, *MIR160f*, *MIR167a*, and *MIR167g*. Also, numerous copies of gibberellin (GA) response (GAR) elements are found in the putative promoter regions of *MIR160f*, *MIR160g*, *MIR160h*, *MIR167a*, *MIR167c*, *MIR167d*, and *MIR167i*. We speculate that GA regulates miRNAs to inhibit the formation of auxin

4 Discussion

The core promoters and *cis*-acting elements of 440 miRNAs from a *G. max* degradome library were analyzed using bioinformatics methods to reveal the general features of the promoters of a large number of soybean miRNAs, examine their *cis*-acting elements, and provide insights into the transcriptional regulation and functions of miRNAs in soybean. Recent studies of *Arabidopsis* and rice showed that most core

promoters contain a TATA-box (Megraw *et al.*, 2006; Cui *et al.*, 2009), and our results for soybean support this observation. Among 699 core promoters predicted, only 4 (0.57%) were TATA-less promoters. Also, 95 of 440 miRNAs (21.59%) contained enhancers. The promoters predicted by bioinformatics analysis will offer valuable information for the design of verification assays and for functional research on miRNAs.

The expression of genes at the transcriptional and post-transcriptional levels is often controlled by small *cis*-acting sequence elements in or near the regulated genes. *Cis*-acting elements have been analyzed extensively (Zhou *et al.*, 2007; Liu H.H. *et al.*, 2008; Liu Y.X. *et al.*, 2010). In this study, *cis*-acting elements of 369 soybean miRNAs were predicted and were classified into several classes according to their functions, such as SRE, GRE, HRE, and TSE. Among those *cis*-acting elements, HRE sites were further analyzed to reveal the potential functional correlation between the transcription factors, miRNAs, and the target genes of the miRNAs. In *Arabidopsis*, the ARF element was found to be over-represented among binding motifs in miRNA promoters. MicroRNAs with putative ARF binding sites upstream may play a role in negative feedback loops that control their own expression (Jones-Rhoades *et al.*, 2006; Megraw *et al.*, 2006; Xie *et al.*, 2010). In soybean, we also identified ARF elements in some miRNA promoters, such as miR160 and miR167. Multiple ARF family members, including *ARF17*, *ARF18*, *ARF6*, and *ARF8*, were predicted to be directly targeted by miR160 and miR167. Although the target genes of soybean miRNAs have not been experimentally verified, targeting of ARFs by soybean miRNAs was indicated by experimental results showing that *Arabidopsis* miR160 and miR167 target ARFs (Mallory *et al.*, 2005; Wang *et al.*, 2005; Wu *et al.*, 2006; Liu *et al.*, 2007). According to the negative feedback model, we propose a possible ARF-mediated negative feedback regulatory loop in soybean, which indicates that miRNAs often act as regulators with many secondary effects on gene expression. We also identified many types of GARF-binding motifs in promoters of *MIR160* and *MIR167*. Gou *et al.* (2010) found that gibberellins regulate lateral root formation in *Populus* through interacting with auxin, and auxin levels increased in GA-deficient and insensitive transgenic

roots, suggesting crosstalk between GA signaling and other hormone pathways. GA may negatively regulate auxin, but the genes and pathways responsible have remained unclear. Here, we found GARF-binding motifs in the promoters of *MIR160* and *MIR167* and propose a GARF-mediated negative feedback regulatory loop for soybean miRNA expression, suggesting that gibberellins regulate the expression of miR160 and miR167 to interplay with auxin. The functions of these computationally identified *cis*-acting elements of soybean miRNA will require experimental validation.

Although few soybean miRNAs have been studied thoroughly, the multiple types of *cis*-acting elements we have identified and the predicted target genes of these soybean miRNAs both indicate that soybean miRNAs have diverse regulatory functions. For example, the gma-miR156 family has 71 types of *cis*-acting elements and regulates 27 target genes. Moreover, the more members in each miRNA family, the more genes the miRNAs target, and the more types of *cis*-acting elements are contained in the promoter of each miRNA.

In summary, we have performed a bioinformatics analysis of the core promoter sequences (TSS, TATA-box, and enhancer) of soybean pre-miRNAs (upstream and downstream sequences) from a *G. max* degradome library. A total of 699 promoters and 102 enhancers for 440 soybean pre-miRNAs were detected. Most miRNA genes were intergenic and there were more core promoters in intergenic genes than in intronic genes. Distribution analysis showed that the locations of the TATA-box and TSS were similar. There were 126 types of *cis*-acting elements identified in 355 miRNA promoters, including SREs, GREs, HREs, and TSEs. In addition, we suggest an ARF- and GARF-mediated negative feedback regulatory loop for soybean miRNA expression. These findings improve our understanding of the specific sequences and motifs upstream of pre-miRNAs in soybean. They also provide a basis for further research on the functional role of miRNA and the regulation of miRNA expression in soybean.

Compliance with ethics guidelines

Yi-qiang HAN, Zheng HU, Dian-feng ZHENG, and Ya-mei GAO declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Bartel, D.P., 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, **116**(2):281-297. [doi:10.1016/S0092-8674(04)00045-5]
- Carrington, J.C., Ambros, V., 2003. Role of microRNAs in plant and animal development. *Science*, **301**(5631):336-338. [doi:10.1126/science.1085242]
- Chen, R., Hu, Z., Zhang, H., 2009. Identification of microRNA in wild soybean (*Glycine soja*). *J. Integr. Plant Biol.*, **51**(12):1071-1079. [doi:10.1111/j.1744-7909.2009.00887.x]
- Chen, X., 2004. A microRNA as a translational repressor of APETALA2 in *Arabidopsis* flower development. *Science*, **303**(5666):2022-2025. [doi:10.1126/science.1088060]
- Cui, X., Xu, S.M., Mu, D.S., et al., 2009. Genomic analysis of rice microRNA promoters and clusters. *Gene*, **431**(1-2):61-66. [doi:10.1016/j.gene.2008.11.016]
- Goodstein, D.M., Shu, S., Howson, R., et al., 2012. Phytozome: a comparative platform for green plant genomics. *Nucl. Acids Res.*, **40**(D1):D1178-D1186. [doi:10.1093/nar/gkr944]
- Gou, J., Strauss, S.H., Tsai, C.J., et al., 2010. Gibberellins regulate lateral root formation in *Populus* through interactions with auxin and other hormones. *Plant Cell*, **22**(3):623-639. [doi:10.1105/tpc.109.073239]
- Guo, N., Ye, W.W., Wu, X.L., et al., 2011. Microarray profiling reveals microRNAs involving soybean resistance to *Phytophthora sojae*. *Genome*, **54**(11):954-958. [doi:10.1139/g11-050]
- Hu, Z., Jiang, Q., Ni, Z., et al., 2013. Analyses of a *Glycine max* degradome library identify microRNA targets and microRNAs that trigger secondary siRNA biogenesis. *J. Integr. Plant Biol.*, **55**(2):160-176. [doi:10.1111/jipb.12002]
- Jones-Rhoades, M.W., Bartel, D.P., 2004. Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Mol. Cell*, **14**(6):787-799. [doi:10.1016/j.molcel.2004.05.027]
- Jones-Rhoades, M.W., Bartel, D.P., Bartel, B., 2006. MicroRNAs and their regulatory roles in plants. *Ann. Rev. Plant Biol.*, **57**(1):19-53. [doi:10.1146/annurev.arplant.57.032905.105218]
- Kulcheski, F.R., de Oliveira, L.F., Molina, L.G., et al., 2011. Identification of novel soybean microRNAs involved in abiotic and biotic stresses. *BMC Genomics*, **12**(1):307. [doi:10.1186/1471-2164-12-307]
- Lee, Y., Kim, M., Han, J., et al., 2004. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.*, **23**(20):4051-4060. [doi:10.1038/sj.emboj.7600385]
- Lescot, M., Dehais, P., Thijs, G., et al., 2002. PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for *in silico* analysis of promoter sequences. *Nucl. Acids Res.*, **30**(1):325-327. [doi:10.1093/nar/30.1.325]
- Li, H., Dong, Y., Yin, H., et al., 2011. Characterization of the stress associated microRNAs in *Glycine max* by deep sequencing. *BMC Plant Biol.*, **11**(1):170. [doi:10.1186/1471-2229-11-170]
- Liu, H.H., Tain, X., Li, Y.J., et al., 2008. Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*. *RNA*, **14**(5):836-843. [doi:10.1261/rna.895308]
- Liu, P.P., Montgomery, T.A., Fahlgren, N., et al., 2007. Repression of *AUXIN RESPONSE FACTOR10* by microRNA160 is critical for seed germination and post-germination stages. *Plant J.*, **52**(1):133-146. [doi:10.1111/j.1365-3113X.2007.03218.x]
- Liu, Y.X., Han, Y.P., Chang, W., et al., 2010. Genomic analysis of microRNA promoters and their cis-acting elements in soybean. *Agric. Sci. China*, **9**(11):1561-1570. [doi:10.1016/S1671-2927(09)60252-2]
- Llave, C., Xie, Z., Kasschau, K.D., et al., 2002. Cleavage of *Scarecrow-like* mRNA targets directed by a class of *Arabidopsis* miRNA. *Science*, **297**(5589):2053-2056. [doi:10.1126/science.1076311]
- Lu, S., Sun, Y.H., Shi, R., et al., 2005. Novel and mechanical stress-responsive microRNAs in *Populus trichocarpa* that are absent from *Arabidopsis*. *Plant Cell*, **17**(8):2186-2203. [doi:10.1105/tpc.105.033456]
- Mallory, A.C., Bartel, D.P., Bartel, B., 2005. MicroRNA-directed regulation of *Arabidopsis AUXIN RESPONSE FACTOR17* is essential for proper development and modulates expression of early auxin response genes. *Plant Cell*, **17**(5):1360-1375. [doi:10.1105/tpc.105.031716]
- Megraw, M., Baev, V., Rusinov, V., et al., 2006. MicroRNA promoter element discovery in *Arabidopsis*. *RNA*, **12**(9):1612-1619. [doi:10.1261/rna.130506]
- Schmutz, J., Cannon, S.B., Schlueter, J., et al., 2010. Genome sequence of the palaeopolyploid soybean. *Nature*, **463**(7278):178-183. [doi:10.1038/nature08670]
- Shahmuradov, I.A., Solovyev, V.V., Gammerman, A.J., 2005. Plant promoter prediction with confidence estimation. *Nucl. Acids Res.*, **33**(3):1069-1076. [doi:10.1093/nar/gki247]
- Shamimuzzaman, M., Vodkin, L., 2012. Identification of soybean seed developmental stage-specific and tissue-specific miRNA targets by degradome sequencing. *BMC Genomics*, **13**(1):310. [doi:10.1186/1471-2164-13-310]
- Smale, S.T., 2001. Core promoters: active contributors to combinatorial gene regulation. *Genes Dev.*, **15**(19):2503-2508. [doi:10.1101/gad.937701]
- Song, Q.X., Liu, Y.F., Hu, X.Y., et al., 2011. Identification of miRNAs and their target genes in developing soybean seeds by deep sequencing. *BMC Plant Biol.*, **11**(1):5. [doi:10.1186/1471-2229-11-5]
- Wang, J.W., Wang, L.J., Mao, Y.B., et al., 2005. Control of root cap formation by microRNA-targeted auxin response factors in *Arabidopsis*. *Plant Cell*, **17**(8):2204-2216. [doi:10.1105/tpc.105.033076]
- Wang, Y., Li, P., Cao, X., et al., 2009. Identification and expression analysis of miRNAs from nitrogen-fixing soybean nodules. *Biochem. Biophys. Res. Commun.*, **378**(4):799-803. [doi:10.1016/j.bbrc.2008.11.140]
- Wong, C.E., Zhao, Y.T., Wang, X.J., et al., 2011. MicroRNAs

- in the shoot apical meristem of soybean. *J. Exp. Bot.*, **62**(8):2495-2506. [doi:10.1093/jxb/erq437]
- Wu, M.F., Tian, Q., Reed, J.W., 2006. *Arabidopsis microRNA167* controls patterns of *ARF6* and *ARF8* expression, and regulates both female and male reproduction. *Development*, **133**(21):4211-4218. [doi:10.1242/dev.02602]
- Xie, Z., Allen, E., Fahlgren, N., et al., 2005. Expression of *Arabidopsis* MIRNA genes. *Plant Physiol.*, **138**(4): 2145-2154. [doi:10.1104/pp.105.062943]
- Xie, Z.X., Khanna, K., Ruan, S., 2010. Expression of microRNAs and its regulation in plants. *Semin. Cell Dev. Biol.*, **21**(8):790-797. [doi:10.1016/j.semcdb.2010.03.012]
- Zeng, H.Q., Zhu, Y.Y., Huang, S.Q., et al., 2010. Analysis of phosphorus-deficient responsive miRNAs and cis-elements from soybean (*Glycine max* L.). *J. Plant Physiol.*, **167**(15): 1289-1297. [doi:10.1016/j.jplph.2010.04.017]
- Zeng, Q.Y., Yang, C.Y., Ma, Q.B., et al., 2012. Identification of wild soybean miRNAs and their target genes responsive to aluminum stress. *BMC Plant Biol.*, **12**(1):182. [doi:10.1186/1471-2229-12-182]
- Zhang, B., Pan, X., Stellwag, E.J., 2008. Identification of soybean microRNAs and their targets. *Planta*, **229**(1): 161-182. [doi:10.1007/s00425-008-0818-x]
- Zhang, Y., Li, Y.K., 2013. MicroRNAs in the regulation of immune response against infections. *J. Zhejiang Univ.-Sci. B (Biomed. & Biotechnol.)*, **14**(1):1-7. [doi:10.1631/jzus. B1200292]
- Zhao, C.Z., Xia, H., Frazier, T.P., et al., 2010. Deep sequencing identifies novel and conserved microRNAs in peanuts (*Arachis hypogaea* L.). *BMC Plant Biol.*, **10**(1):3. [doi:10.1186/1471-2229-10-3]
- Zhou, X., Ruan, J., Wang, G., et al., 2007. Characterization and identification of microRNA core promoters in four model species. *PLoS Comput. Biol.*, **3**(3):e37. [doi:10.1371/journal.pcbi.0030037]
- Zhu, Q.W., Luo, Y.P., 2013. Identification of miRNAs and their targets in tea (*Camellia sinensis*). *J. Zhejiang Univ.-Sci. B (Biomed. & Biotechnol.)*, **14**(10):916-923. [doi:10.1631/jzus.B1300006]

List of electronic supplementary materials

- Table S1 Statistics of the core promoters and enhancers of soybean miRNA genes
- Table S2 Statistics of cis-acting elements in the predicted promoters
- Table S3 Function types of cis-acting elements in the predicted promoters

中文概要:

本文题目: 大豆降解组文库 microRNAs 的启动子分析

Analysis of promoters of microRNAs from a *Glycine max* degradome library

研究目的: 通过分析 miRNA 的核心启动子和顺式作用元件为进一步解析大豆 (*Glycine max*) miRNAs 表达调控及其功能研究提供重要信息。

创新要点: 利用生物信息学方法全面解析了大豆降解组文库 miRNA 的启动子特征, 并依据顺式作用元件及靶基因构建了 miRNA 的表达与生长素响应因子、赤霉素响应因子之间存在潜在的负反馈调控网络。

研究方法: 本研究利用 TSSP 程序和 PlantCARE 数据库预测了来自大豆降解组文库的 440 个 miRNA 的核心启动子以及 369 个 miRNAs 的顺式作用元件, 并依据顺式作用元件及靶基因构建 miRNA 调控网络。

重要结论: 83.86% 的 miRNA 在其上游序列中含有启动子, 8.64% 的 miRNA 在其下游序列中含有启动子, 21.59% 的 miRNA 包含增强子。核心启动子的 TATA 盒与转录起始位点 (TSSs) 的分布相似 (见图 2)。此外, 对转录起始位点 5' 端的顺式作用元件预测为 miRNAs 的可能功能和表达的时空性提供了线索。miRNAs 的顺式作用元件和靶基因的分析显示, 部分 miRNA 的表达与生长素响应因子、赤霉素响应因子之间存在潜在的负反馈调控 (见图 3)。

关键词组: 大豆; MicroRNA (miRNA); 启动子; 顺式作用元件; 预测