Metabolic Aspects and Mechanisms

7.14 Insulin Sensitivity and Cerebrovascular Disease in Elderly Non-Diabetic Hypertensive Subjects with the Metabolic Syndrome

D. Recchi,¹ P. Di Biase,² M.G. Di Niro,¹ O. Bollella,¹ P. Martone¹

(1) Geriatric Division – Regional Hospital “A. Cardarelli” – Ca Campobasso; (2)

Dietetics & Clinical Nutrition Unit – Regional Hospital “A. Cardarelli” –
Campobasso, It Italy

Introduction: To determine the association between insulin sensitivity, which is altered in subjects with Metabolic Syndrome (MS), and cerebrovascular diseases (TIA and Stroke) in a group of hypertensive treated non diabetic elderly subjects with MS.

Methods: 30 elderly non diabetic subjects with treated arterial hypertension and MS were studied. Of these, 10 subjects had a TIA or Stroke and MS while the remaining 20 subjects represented the control group (of which 12 with MS and 8 without). All subjects were free from respiratory, renal, liver, thyroid diseases or cancer and did not show signs of malnutrition, the controls were also free from cardiovascular diseases. All subjects were treated with an antihypertensive therapy (diuretics, beta-blockers, calcium channel blockers, ACE inhibitors or angiotensin II receptor blockers) at home. Only one subject was on a statin.

Blood pressure was evaluated at admission and insulin sensitivity was estimated using the HOMA index. $HOMA = (FPGSI \times FPI)/2.5$. MS was diagnosed according to the National Cholesterol Education Program Adult Treatment Panel III criteria.

Results: No differences between the two groups were found for: Age (79 ± 7.76 years vs. 79.5 ± 5.2 years, ns), Body Mass Index values (26.66 ± 2.17 vs. 25.87 ± 4.46 , ns); Systolic Blood Pressure values (141 ± 26.85 mmHg vs. 142.89 ± 28.98 mmHg, ns); Diastolic Blood Pressure values (71.5 ± 12.03 mmHg vs. 80.53 ± 13.22 mmHg, ns). Likewise, HOMA index values did not differ between the two groups (1.54 ± 0.96 vs. 1.55 ± 1.22 , ns).

Conclusions: Our data shows that there is no association between insulin sensitivity and cerebrovascular disease in hypertensive treated non diabetic elderly subjects with MS.