Discrimination functions predicted from categories in speech and music

JAMES E. CUTTING

Wesleyan University, Middletown, Connecticut 06457
and

Haskins Laboratories, New Haven, Connecticut 06511

and

BURTON S. ROSNER

University of Pennsylvania
Philadelphia, Pennsylvania 19174

Sawtooth waves differing only in rise time are identifiable as *plucked* or *bowed* notes from a stringed instrument. We previously reported (Cutting & Rosner, 1974) that these nonlinguistic sounds are perceived categorically. We also synthesized a continuum of speech sounds by varying only rise time. Listeners identified these sounds as /tʃa/ or /ʃa/, as in CHOP or SHOP, respectively, and perceived them categorically as well.

Our criteria for categorical perception were those suggested by Studdert-Kennedy, Liberman, Harris, and Cooper (1970): (a) "peaks" of high discriminability between stimuli in restricted regions along the dimension studied, (b) "troughs" of discrimination performance near chance in regions on either side of the peak, and (c) correspondence between peaks and troughs and the course of identification functions, with peaks occurring at identification boundaries and troughs occurring within each perceptual category. Categorical perception is therefore revealed by a particular combination of results from identification and discrimination tasks. This convergence between identification and discrimination is unusual: A listener generally can discriminate many more stimuli than he or she can identify absolutely (see, for example, Miller, 1956).

The correspondence between identification and discrimination can be tested quantitatively. Discrimination performance can be predicted from identification data by assuming that discrimination is no better than identification. To the extent that obtained and predicted discrimination scores do not differ significantly, categorical perception has occurred.

Our previous paper described such agreement between obtained and predicted discrimination scores for both the linguistic and musical sounds (Cutting & Rosner, 1974). Unfortunately, the pre-

dicted functions were derived through an incorrect formula. This note corrects that error.

To predict discrimination from identification of a two-category continuum, the correct formula for an ABX discrimination task is:

$$P(c) = \frac{1}{2}[1 + (p_1 - p_2)^2], \quad (1)$$

where $P(c)$ is the probability of a correct discrimination, p_1 is the probability of assigning Stimulus A to one of the categories, and p_2 is the probability of assigning Stimulus B to that same category. The original formula for the three-category case published by investigators at the Haskins Laboratories (Liberman, Harris, Hoffman, & Griffith, 1957) is incorrect; Pollack and Pisoni (1971) give proper formulae for both two- and three-category continua. We will refer to Equation 1 as the Haskins prediction.

Typically, obtained discrimination functions, even for stop consonants, systematically exceed predicted functions by as many as 10 percentage points at each comparison along the stimulus array. Thus, the strongest possible relationship between identification and discrimination is not realized (see also Barclay, 1972; Pisoni & Lazarus, 1974; and Pisoni & Tash, 1974). The discrepancy between obtained and predicted discrimination functions is even larger for more "continuously" perceived stimuli such as vowels (Pisoni, 1971, 1973, 1975). By further developing a model that Fujisaki and Kawashima (1970) formulated, Pisoni added a correction factor to prediction formulae such as (1). This factor is based on the asymptotic through discrimination value, it raises the predicted functions by several percentage points, and it can be interpreted as measuring short-term auditory storage for differences between two stimuli identified alike. For a two-category continuum in an ABX task, the proper Fujisaki-Kawashima prediction formula is

$$P(c) = \frac{1}{2}[(p_1 - p_2)^2 + p_1(1 - p_2) + p_2(1 - p_1)] \\ + [p_1p_2 + (1 - p_1)(1 - p_2)]T, \quad (2)$$

where $P(c)$, p_1 , and p_2 are the same as in Equation 1 and T is the asymptotic trough value of the obtained discrimination function. If $T = 0.50$, Equation 2 reduces to Equation 1. Like the Haskins prediction formula, the Fujisaki-Kawashima formula has suffered the misfortune of appearing incorrectly in print (Pisoni, 1971, p. 44; Pisoni, 1975, p. 13).¹

Using the correct formulae, we have recomputed both the Haskins and the Fujisaki-Kawashima predictions for our data on discrimination of sawtooth waves and of affricate-fricative speech syllables. Pre-

We thank Neil Macmillan for pointing out our error to us and for suggesting how it occurred.

Table 1
Obtained and Correctly Predicted Discrimination Values for Stimuli Differing in Rise Time

	Rise Time Comparison (msec)						
	0-20	10-30	20-40	30-50	40-60	50-70	60-80
Experiment I							
Sawtooth Wave Stimuli							
Obtained	61	64	72	78	58	60	59
Haskins Predicted	50	51	60	67	56	52	51
Fujisaki-Kawashima Predicted	58	59	65	71	61	58	58
Speech Stimuli							
Obtained	61	58	59	70	76	61	58
Haskins Predicted	51	53	55	62	64	51	50
Fujisaki-Kawashima Predicted	58	59	60	67	68	58	58
Experiment II							
Sawtooth Wave Stimuli							
Obtained	61	55	66	72	47	50	53
Haskins Predicted	50	50	66	73	54	51	50
Sine Wave Stimuli							
Obtained	54	49	56	68	56	58	53
Haskins Predicted	50	51	63	68	54	54	53

Note—The original predicted functions that appear in Cutting and Rosner (1974) are incorrect.

dictions were made for each individual listener, then averaged functions were obtained from the individual functions, as Pisoni (1971) suggests.² Table 1 shows averaged obtained and predicted discrimination scores.

The predicted functions in Table 1 are farther below the obtained functions than were those originally published (see Tables 1 and 2 in Cutting & Rosner, 1974). Nevertheless, the discrepancies between predicted and obtained scores here are not marked. Goodness-of-fit measures calculated from individual obtained and Haskins-predicted scores revealed no significant differences (see Pisoni, 1971, p. 20), although the observations per comparison may be too few to make small differences statistically reliable. The fit between the data and the correct predictions still supports our prior conclusion that musical stimuli and affricate-fricative consonants differing in rise time are each perceived categorically. Subsequent experiments have provided confirmation: Cutting, Rosner, and Foard (1976) have demonstrated that the musical sounds are perceived as categorically as stop consonants in Pisoni's (1971, 1973) variable-interval AX discrimination task.

In summary, this note presents correct predicted discrimination functions for data previously published (Cutting & Rosner, 1974). The corrections leave the principal conclusion of that study unchanged: Nonlinguistic and linguistic stimuli synthesized with different rise times are perceived categorically. In addition, this note provides correct formulae for predicting discrimination functions. Several previous sources for the formulae are in error.

REFERENCES

- BARCLAY, J. R. Noncategorical perception of a voiced stop: A replication. *Perception & Psychophysics*, 1972, 11, 269-273.
- CUTTING, J. E., & ROSNER, B. S. Categories and boundaries in speech and music. *Perception & Psychophysics*, 1974, 16, 564-570.
- CUTTING, J. E., ROSNER, B. S., & FOARD, C. F. Perceptual categories for musiclike sounds: Implications for theories of speech perception. *Quarterly Journal of Experimental Psychology*, 1976, 28, in press.
- FUJISAKI, H., & KAWASHIMA, T. Some experiments on speech perception and a model for the perceptual mechanisms. *Annual Report of the Engineering Research Institute*, Tokyo, 1970, 29, 207-214.
- LIBERMAN, A. M., HARRIS, K. S., HOFFMAN, H. S., & GRIFFITH, B. C. The discrimination of speech sounds within and across phoneme boundaries. *Journal of Experimental Psychology*, 1957, 54, 358-368.
- MILLER, G. A. The magical number seven, plus or minus two, or some limits on our capacity for processing information. *Psychological Review*, 1956, 63, 81-96.
- PISONI, D. B. On the nature of categorical perception of speech sounds. (Doctoral dissertation, University of Michigan, 1971). *Dissertation abstracts International*, 1972, 32, 6693B. (University Microfilms No. 72-14,964)
- PISONI, D. B. Auditory and phonetic memory codes in the discrimination of consonants and vowels. *Perception & Psychophysics*, 1973, 13, 253-260.
- PISONI, D. B. Auditory short-term memory and vowel perception. *Memory & Cognition*, 1975, 3, 7-18.
- PISONI, D. B., & LAZARUS, J. H. Categorical and noncategorical modes of speech perception along the voicing continuum. *Journal of the Acoustical Society of America*, 1974, 55, 328-333.
- PISONI, D. B., & TASH, J. Reaction times to comparisons within and across phonetic categories. *Perception & Psychophysics*, 1974, 15, 285-290.
- POLLACK, I., & PISONI, D. B. On the comparison between identification and discrimination tests in speech perception. *Psychonomic Science*, 1971, 24, 299-300.
- STUDDERT-KENNEDY, M., LIBERMAN, A. M., HARRIS, K. S., & COOPER, F. S. Motor theory of speech perception: A reply to Lane's critical review. *Psychological Review*, 1970, 77, 234-249.

NOTES

- Page numbers for Pisoni (1971) refer to a version published as a supplement to the Haskins Laboratories Status Report on Speech Research.
- The trough value T was not stable for individual listeners; we assumed it to be 0.60 for all listeners for both sets of stimuli represented in Table 1.

(Received for publication June 21, 1976;
accepted June 23, 1976.)