Stimulus probability and choice reaction time*

A. R. BLACKMAN

Simon Fraser University, Burnaby, British Columbia, Canada

Two experiments, in which three stimuli were paired with two responses in a 2:1, 1:1 arrangement, showed choice RT to be a decreasing function of the relative frequency of stimulus occurrence when response frequencies are equated. The size of effect varied significantly with stimulus type (Experiment 1), being larger for circular light patches of differing diameter than for sets of colors, letters, and spatially distributed light sources, but was independent of differences in the similarity of stimulus alternatives within a set (Experiment 2). The results are discussed in terms of various strategies of stimulus identification.

Several studies have demonstrated that performance in a choice reaction time (RT) task is a function of the relative frequencies of occurrence of the various stimulus-response alternatives (e.g., Hyman, 1953; Stone & Calloway, 1964). There is some disagreement, however, as to whether this relative frequency effect arises at the stimulus discrimination or at the response selection stage of the choice process. Dillon (1966) has argued for response uncertainty as the critical variable. The results of other studies point to stimulus probability as the variable determining this relative frequency effect on RT (Bertelson & Tisseyre, 1966; Biederman & Zachary, 1970; Hawkins, Thomas, & Drury, 1970; Orenstein, 1970). Yet others have implicated both stimulus and response probability (Blackman, 1972; LaBerge & Tweedy, 1964; LaBerge, Legrand, & Hobbie, 1969).

A comparison of the designs employed in these experiments reveals a number of major differences, two of which may account for the discrepant results. First, the difference in probability between the most and least frequently occurring stimulus ranges from .15 (Dillon, 1966) to .75 (Hawkins et al., 1970; LaBerge et al., 1969). Second, various types of stimuli have been used, including letters (Bertelson & Tisseyre, 1966; Dillon, 1966; Hawkins et al., 1970), colors (Biederman & Zachary, 1970; LaBerge & Tweedy, 1964; LaBerge

et al., 1969), spatially arrayed lights (Orenstein, 1970), and geometric shapes (Biederman & Zachary, 1970).

Most investigators have sought partially to unconfound stimulus and response frequency by including many:one pairings of stimulus and response, thus allowing stimulus frequencies to be manipulated while response frequencies were held constant. This technique was used in the present investigation, which was designed to provide evidence regarding the influence of degree of probability imbalance and stimulus type on the relative frequency effect.

EXPERIMENT 1

The purpose of this experiment was to determine the effect on two-choice RT of differences in stimulus probability, response requirements being held constant and equal. The second independent variable was stimulus type, the four types used being circles of differing size, colors, letters of the alphabet, and spatially distributed lights (lamps). The effects of a third variable, practice, were assessed by repeating the basic experiment with the same Ss twice.

Method

Design. A within-S design was used, each S performing in all parts of a 3 (stimulus alternatives) by 3 (conditions of stimulus frequency imbalance) by 4 (stimulus types) by 3 (weeks) factorial design.

Subjects. The Ss were four male undergraduate students, aged between 17 and 21 years, from an introductory psychology course.

Stimuli and apparatus. Four different stimulus types were used, each set consisting of three alternatives: (1) Circles—circular patches of white light with diameters of 11/32, 17/32, and 23/32 in. (2) Colors—1¼-in.-square patches of colored light; red, green, and yellow. (3) Letters—5/8-in.-high x

7/16-in.-wide uppercase letters; A, U, and T. (4) Lamps—28-V bulbs with 3/8-in.-diam translucent plastic covers, centers spaced 1½ in. apart. The circles were displayed on a multiple stimulus projector (Grason-Stadler, Model 10657), and the letters and colors were displayed on a similar unit (Model 1070R-44-L). The lamps, which were mounted vertically in a metal frame, and the projectors were situated at eye level on a large wooden screen placed about 2 ft from S. The response unit consisted of two button-operated microswitches, mounted on a wooden platform on the table in front of S. The buttons, which had a diameter of ¾ in., were centered 2 in. on either side of a "home" button on which S rested his finger between responses. A warning signal, in the form of a brief burst of clicks, was delivered to S through padded earphones.

Stimulus presentation was controlled by a 12-channel magnetic tape programmer. This device enabled different stimuli to be presented for precise durations at precise intervals. Signals were prerecorded in the appropriate channels at the required positions on the tape, and on playback, these signals activated the corresponding stimulus circuits. A chronoscope (Hunter KlockCounter, Model 120A) was triggered by stimulus onset, providing the home button was depressed, and stopped by depression of either response button, allowing response latency to be measured to the nearest millisecond. Signal lamps in each stimulus and response circuit permitted E to observe errors. S was seated in a dimly illuminated room facing the stimulus display, while E monitored the programming and timing apparatus in an adjacent room.

Procedure. In the first week, S had one practice and four experimental sessions, each on successive days, a single stimulus type being used throughout a session. The order in which stimulus types were assigned to each S's four experimental sessions was determined by a semibalanced Latin square. With the exception of the practice session, two replications using the same Ss and experimental design were carried out over the following 2 weeks. This gave each S 3 weekly sessions with each of the four stimulus types, for a total of 12 experimental sessions.

A session consisted of 300 trials, 100 in each of the three conditions of stimulus frequency imbalance. Table 1 shows the probability of occurrence of the three stimulus-response pairs in each condition. Stimulus S1, which occurred on half the trials in each condition, was paired with

*Condensed from a dissertation submitted to the Faculty of Graduate Studies and Research, McGill University, Montreal, in partial fulfillment of the requirements for the PhD degree. I would like to thank Dalbir Bindra for his critical reading of an earlier version of the manuscript and Shizuhiko Nishisato for his assistance in computer programming. This research was supported by Contract Nonr-4896(00) between the U.S. Office of Naval Research and McGill University and by Grant 9426-10 from the Defence Research Board of Canada to Dalbir Bindra.

Table 1
Probability of Occurrence of Each Stimulus-Response Pair in Each Condition of Stimulus Frequency Imbalance in Experiment 1

Condition of Stimulus Frequency Imbalance	Stimulus-Response Pair		
	S1-R _a	S2-R _b	S3-R _b
1	.50	.25	.25
2	.50	.35	.15
3	.50	.45	.05

Response R_a. In the case of the circles and lamps, S1 was always the stimulus of intermediate size or position; for the letters and colors, it was T and yellow, respectively. For two Ss, Response R_a was pressing the left button, and for the other two Ss, it was pressing the right button. Stimulus S2, whose probability of occurrence increased from Condition 1 to Condition 3, and Stimulus S3, whose probability of occurrence correspondingly decreased, were both paired with Response R_b. For each stimulus type, the particular stimulus alternatives designated as S2 and S3 were fully balanced over Ss and partially balanced over weeks.

The order of trials within each condition was randomized, with the restrictions that the average sequential response probabilities, $p(R_a \text{ following } R_a)$ and $p(R_b \text{ following } R_a)$, be set equal to .5 in all conditions, and that the largest run of repeated stimuli or responses be four. The 100 trials in each condition were split into three parts of 20, 40, and 40 trials. The first part was considered as practice, allowing S to accommodate to the particular stimulus frequency bias in the series, and the data from these trials were discarded. The order in which the second and third parts were given was varied. This feature, together with the alternating use of two different programs and the varied order of conditions within a session, was considered sufficient to make it virtually impossible for S to learn what the sequence of stimuli or responses would be on any given set of trials.

At the start of each session, S was shown examples of the relevant stimulus type. A card depicting these stimuli, and indicating the relative frequency of occurrence of each stimulus alternative, could be seen by S throughout the session. A payoff scheme was adopted in which 1c was awarded for each response that was faster than t sec (t being determined from performance on the initial practice trials in each session so as to reward S on about 75% of the trials); a penalty of 5c was levied for each mistake. Rest breaks of 1 min were given between each part, and of 5 min between each condition. The warning

signal was presented 1 sec prior to the occurrence of the stimulus. The duration of each stimulus was 100 msec and the interstimulus interval was 7 sec.

Results

Data for trials on which S made an error were not included in the analysis of RT. The harmonic mean was chosen as the best estimate of RT for each cell of the 3 by 3 by 4 by 3 matrix for each of the four Ss. The number of RTs contributing to each mean varied between 4 and 36 (except where omission of data for erroneous responses reduced these numbers), according to the frequency of occurrence of the particular stimulus-response pair.

The results are summarized in Fig. 1, which shows harmonic mean RT, averaged over weeks and Ss as a function of stimulus probability for each stimulus type. A general trend

can be observed for shorter RTs to be associated with higher stimulus probabilities. This effect is most consistent and pronounced for the circles, but is less evident, except at the extreme probability values, for the other types.

Before applying analyses of variance to the latency data, consideration was given to the comparability of results for the S1 and S2/S3 stimulus alternatives. For the lamps, letters, and colors, the shortest average RT was associated with S1, the most probable stimulus ($p = .5$). This was not the case for the circles, where the average RT to S1 was longer than that to S2 and S3, even when they occurred only half as frequently (Condition 1). This discrepancy probably stems from a difference between the confusability of S1 and that of S2 and S3 for the circles (but not for the other stimulus types). That is, for the lamps, letters, and colors, the alternatives within a set

Fig. 1. Average harmonic mean RT in milliseconds for Stimuli S1, S2, and S3 in Experiment 1, as a function of their probability of occurrence, for each stimulus type.

were considered to be of approximately equal confusability, and the particular stimulus always assigned to S1 was as readily distinguished from its alternatives as were those stimuli assigned to S2 and S3. With the circles, however, it seems likely that the inflated latency of response to S1 (always the middle-sized circle) reflects the fact that its alternatives comprised both a larger and a smaller circle, whereas the alternatives to S2 and S3 were either both smaller or both larger. This intraset confounding of confusability and probability for the circles does not obtain when comparing the overall results for S2 and S3, particularly in view of the balancing employed with each type in the assignment of the remaining two stimuli to S2 and S3. For this reason, the subsequent analyses treat the RTs for S1 separately from those for S2 and S3.

An analysis of variance (Winer, 1962, Chap. 7) was carried out on the harmonic mean RTs to S1. The only significant effects were those for stimulus types [$F(3,9) = 26.21$, $p < .001$] and weeks [$F(2,6) = 6.34$, $p < .05$], the latter indicating an overall decrease in RT to S1 from Week 1 to Week 3.

In the analysis of variance of latencies to Stimuli S2 and S3, significant main effects were found for stimulus type [$F(3,9) = 38.23$, $p < .001$] and for stimulus probability¹ [$F(4,12) = 16.95$, $p < .001$]. The only other significant term was the Type by Probability interaction [$F(12,36) = 3.71$, $p < .005$], reflecting the greater effect of probability on RT for the circles than for the other stimulus types.

The adequacy of the balancing procedures used in the assignment of particular stimuli to S2 and S3 can be assessed by comparing the average latencies to these two stimuli in Condition 1, where their probabilities of occurrence were equal ($p = .25$). For the four stimulus types, the greatest difference in average RT to S2 and S3 was 6 msec, which indicates that the design was successful in avoiding any systematic bias arising from differential latencies to particular stimulus alternatives independent of their probability.

In the S2/S3 analysis of variance, the insignificant main effect for weeks suggests that S reached a stable level of responding within the first week. Further inspection of the data, however, revealed noticeably different trends of the relative frequency effect with practice for the different stimulus types. The relative frequency effect obtained in Condition 3 for the colors, lamps, and letters was substantial in the first week of the experiment, but

showed an overall decrease with practice. By the last week, in fact, the effect for the letters had completely disappeared. On the other hand, there was no systematic change with practice in the much larger relative frequency effect observed for the circles.

Excluding practice trials, 11,520 responses were recorded. Of these, 82 were incorrect, representing an error frequency of just over 0.7%. The number of errors for each stimulus type was as follows: circles, 35; lamps, 24; colors, 13; and letters, 10.

Discussion

The results clearly indicate that the occurrence of a relative frequency effect is a function of both stimulus type and degree of stimulus probability imbalance.

The apparent discrepancy between the results of Dillon (1966) and those of LaBerge and Tweedy (1964) and Bertelson and Tisseyre (1966) may be resolved by noting that Dillon found no relative frequency effect using letters with a stimulus probability difference of .15, while the others did find an effect using letters with a probability difference of .4. Both sets of results are consonant with those of the present study, in which a significant effect for letters was found, at least in Week 1, with a probability difference of .4 (Condition 3), but not when the difference was only .2 (Condition 2). Any consideration of Dillon's data, however, should take note of the cogent criticisms of his experimental procedures offered by Biederman and Zachary (1970).

The relative frequency effect was larger, and became evident at a lower degree of probability imbalance, for the circles than for the other stimulus types. Because response latency was typically longest and error rate highest for the circles, it is plausible to interpret these intertype variations in the effect of stimulus probability in terms of differences in the level of intratype discriminability or confusability of stimulus alternatives. However, factors other than confusability also distinguished the circles from the other stimulus types, e.g., geometric vs nongeometric, familiar vs unfamiliar. The purpose of Experiment 2 was to dissociate stimulus confusability from these other variables in order to determine its influence on the size of the relative frequency effect.

EXPERIMENT 2

Two of the stimulus sets selected contained circles, the members being of such sizes that the alternatives in one set were less similar, and were intended to be less confusable than

those in the other set. These were labeled *easy circles* and *difficult circles*, respectively. It was also considered likely that stimulus discrimination would be easier if there were some qualitative basis on which S could categorize Stimuli S2 and S3 (both associated with Response R_b) separately from S1 (associated with R_a). To this end, a third set of stimuli was used in which the middle-sized member of the easy circles set was replaced by a square of approximately the same area (*circles/square* set). Finally, in an attempt to manipulate the confusability of a different stimulus type, three *letters* were selected whose forms were more similar than those used in Experiment 1. In addition, neutral density filters were used to reduce further the discriminability of the letters. It was hypothesized that if the differences in size of the relative frequency effect for the stimulus types used in Experiment 1 were the result of intertype differences in average stimulus confusability, then the effect of stimulus probability in Experiment 2 would be greater for the difficult circles and letters than for the circles/square and easy circles.

Method

Subjects. The Ss were eight male undergraduate students, aged between 17 and 24 years, from an introductory psychology course; none of them participated in Experiment 1.

Stimuli and apparatus. Four stimulus sets were used: (1) Difficult circles—circular patches of white light, 11/32, 14/32, and 17/32 in. in diam. (2) Easy circles—circular patches of white light, 3/32, 14/32, and 1 in. in diam. (3) Circles/square—same as (2) but with middle-sized circle replaced by 13/32-in.-square patch of white light. (4) Letters—5/8-in.-high $\times$ 7/16-in.-wide uppercase letters, O, U, and V, viewed through neutral density filter (Kodak, value 3.4).

For the difficult and easy circles, Stimulus S1 was always the middle-sized circle; for the circles/square set, it was the square; and for the letters, it was the letter O.

The basic experimental design was the same as that used in the first experiment, except that only one replication was included, and thus the sessions extended over 2 rather than 3 weeks. The rest of the stimulus, response, and programming apparatus, the experimental procedure, and the method of data analysis were the same as in Experiment 1.

Results

Harmonic mean RT, averaged over Ss and weeks, is shown in Fig. 2 as a function of stimulus probability for

Fig. 2. Average harmonic mean RT in milliseconds for Stimuli S1, S2, and S3 in Experiment 2, as a function of their probability of occurrence, for each stimulus set.

each stimulus set. The dependence of RT on stimulus probability is clearly apparent for all stimulus sets. As was argued for Experiment 1, it appears likely that the latencies for the highest probability stimulus (S1) were differentially influenced by variations in intraset confusability. Thus, the S1 and S2/S3 data are treated separately.

An analysis of variance of the harmonic mean RTs for S1 revealed only one significant effect, that of stimulus sets [$F(3,21) = 41.20$, $p < .001$]. The corresponding term in the analysis of the S2/S3 data was also significant [$F(3,21) = 48.00$, $p < .001$], as was the main effect for stimulus probability¹ [$F(4,28) = 16.16$, $p < .001$]. The existence of a Stimulus Sets by Probability interaction was of particular interest; however, this was not a source of significant variance [$F(12,84) = .53$, $p > .10$]. None of the remaining terms was significant at the 5% level.

Of a total of 15,360 responses, 376 (2.4%) were in error. A chi-squared test showed the observed and expected

distributions of errors over stimuli and conditions to be significantly different [$\chi^2(9) = 29.3$, $p < .01$]. This is largely due to the fewer errors than expected for Stimulus S1 and the greater number than expected for Stimulus S2 in Conditions 2 and 3. The number of errors for each stimulus set was as follows: easy circles, 48; difficult circles, 204; circles/square, 50; letters, 74.

DISCUSSION

The results of Experiment 2 provide no support for the hypothesis that the magnitude of the relative frequency effect is influenced by differences in stimulus confusability or similarity. The slopes of the function relating probability and RT are not significantly different for the three sets containing circles, although the main effects of stimulus probability and stimulus sets (similarity) are clearly apparent. While the relative frequency effect for the letters appears somewhat greater than that found in Experiment 1, direct comparison of

the two sets of data is made difficult by the difference between the two studies in the number of sessions S had with each stimulus set.

The only other experiment using a stimulus-response pairing arrangement similar to that in the present study which investigated the joint influence of stimulus probability and discriminability on RT was reported by Orenstein (1970). While he states that "... [decreased] discriminability seemed to enhance stimulus frequency effects [p. 41]" for leverpress responses to spatially arrayed lights, his data do not appear to provide unambiguous support for this conclusion. If one considers only his RT data for the two stimuli sharing the same response (Orenstein's C and R, comparable to S2 and S3 in the present study) in the two conditions in which the stimulus alternatives within a set were evenly spaced (Orenstein's 1, 4, 7 and 3, 4, 5, comparable to the easy and difficult circles, respectively), there is little evidence of an interaction between stimulus probability and discriminability. It seems that Orenstein's conclusion is based primarily on the results obtained in a third condition (3, 4, 7). But interpretation of data from this condition is confounded with intraset differences in discriminability.

Although stimulus discriminability does not appear to be a determinant of the relative frequency effect, a number of other variables have been observed to influence the effect of stimulus probability on two-choice RT. Sternberg (1966) manipulated relative stimulus frequency by varying the number of equiprobable digits in the positive set, whose membership was explicitly defined before each trial. He found stimulus probability to be a determinant of RT for the positive set but not the negative set. Varying stimulus probability directly, Hawkins and Hosking (1969) provided support for this view, but argued that the likelihood that S will treat a particular stimulus set as positive is less when the alternative set comprises a single stimulus and when the size imbalance between the two sets is minimal. Additional evidence for the latter limitation can be found in Krueger's (1970) demonstration of a relative frequency effect in both positive and negative sets with stimuli and responses paired in a 4:1, 4:1 arrangement.

Hawkins and Hosking's (1969) Ss reported rehearsing (treating as positive) only the singly paired stimulus in a 4:1, 1:1 condition. Noting that this stimulus elicited the slowest RTs and highest error rate, Hawkins and Hosking (1969) argue that although S rehearsed it before

each trial, he did not expect it to occur (it was not the most probable stimulus). The awkwardness of this position can be avoided by appealing to the influence of response probability. That is, it can be argued that their Ss did first attempt to identify the presented stimulus by comparing it with the stored representation of the singly paired stimulus, but the low probability (.25) of the associated response led to an increased latency and error rate arising at the response selection stage.

Hawkins and Hosking's (1969) concurrent demonstration of a significant relative frequency effect in both stimulus sets with a 3:1, 2:1 pairing is noteworthy. It precludes an accounting of the absence of such effects in their 4:1, 1:1 condition in terms of ineffective degree of probability imbalance or insensitive experimental design, arguments that weaken confidence in the similar findings of Dillon (1966). While the utility of the concept of positive and negative sets has been demonstrated with relatively large ($n > 5$) stimulus ensembles, its explanatory value is less apparent with two-choice tasks involving only three stimuli, such as that used in the present investigation. Since the singly paired stimulus (S1) was always the most frequently occurring, an optimal strategy would have been for S to treat it as the positive set. The relative frequency effect observed in the other set (Stimuli S2 and S3) indicates that this did not occur, at least in any consistent fashion.

An alternative strategy that S might adopt is to attempt serial comparisons of the presented stimulus with stored representations of the three stimuli in the order of their probability of occurrence. That is, S might treat the whole ensemble as positive. If it is further assumed that the consistency of this ordering of comparisons is a function of the degree of probability imbalance, then this would account for the observed relative frequency effect. Biederman and Zachary (1970) considered such an interpretation of

their data but rejected it. Their argument is based on the longer RT obtained for a stimulus with a probability of .3 in an ensemble whose probabilities were .6, .3, and .1 (variously paired 2:1 and 1:1), compared with that for stimuli of the same probability in a .4, .3, .3 condition. Ordering of the comparisons by probability would, indeed, predict the reverse relation for these RTs, but only if S consistently followed this strategy in both conditions. Their failure to demonstrate a significant relative frequency effect with probabilities of .4, .3, and .3 weakens their argument and suggests that the order in which S makes the comparisons is little influenced by such minimal differences in stimulus probability.

The evidence reviewed above indicates that the influence of stimulus probability on two-choice RT is a complex function of the size of the stimulus ensemble, the relative set size, and the presence of a singly paired stimulus. The results of the present investigation suggest that stimulus type should be added to this list. Since the size of the relative frequency effect was shown to be independent of stimulus discriminability or similarity, the nature of the critical dimension underlying the influence of stimulus type remains to be identified. One possible candidate is stimulus codability, a characteristic which Brown and Lenneberg (1954) showed to be independent of stimulus discriminability (perceptual similarity) in determining color-naming performance. Cohen (1969) has recently used this variable to account for some of the variance in recognition RT with a range of stimulus types.

REFERENCES

- BERTELSON, P., & TISSEYRE, F. Choice reaction time as a function of stimulus versus response relative frequency of occurrence. *Nature*, 1966, 212, 1069-1070.
- BIEDERMAN, I., & ZACHARY, R. A. Stimulus versus response probability in choice reaction time. *Perception & Psychophysics*, 1970, 7, 189-192.

- BLACKMAN, A. R. Influence of stimulus and response probability on decision and movement latency in a discrete choice reaction task. *Journal of Experimental Psychology*, 1972, 92, 128-133.
- BROWN, R. W., & LENNEBERG, E. H. A study in language and cognition. *Journal of Abnormal & Social Psychology*, 1954, 49, 454-462.
- COHEN, G. The effect of codability of the stimulus on recognition reaction times. *British Journal of Psychology*, 1969, 60, 25-29.
- DILLON, P. J. Stimulus versus response decisions as determinants of the relative frequency effect in disjunctive reaction-time performance. *Journal of Experimental Psychology*, 1966, 71, 321-330.
- HAWKINS, H. L., & HOSKING, K. Stimulus probability as a determinant of discrete choice reaction time. *Journal of Experimental Psychology*, 1969, 82, 435-440.
- HAWKINS, H. L., THOMAS, G. B., & DRURY, K. B. Perceptual versus response bias in discrete reaction time. *Journal of Experimental Psychology*, 1970, 84, 514-517.
- HYMAN, R. Stimulus information as a determinant of reaction time. *Journal of Experimental Psychology*, 1953, 45, 188-196.
- KRUEGER, L. E. Effect of stimulus probability on two-choice reaction time. *Journal of Experimental Psychology*, 1970, 84, 377-379.
- LaBERGE, D., LEGRAND, R., & HOBIE, R. K. Functional identification of perceptual and response biases in choice reaction time. *Journal of Experimental Psychology*, 1969, 79, 295-299.
- LaBERGE, D., & TWEEDY, J. R. Presentation probability and choice time. *Journal of Experimental Psychology*, 1964, 68, 477-481.
- ORENSTEIN, H. B. Reaction time as a function of perceptual bias, response bias, and stimulus discriminability. *Journal of Experimental Psychology*, 1970, 86, 38-42.
- STERNBERG, S. High-speed scanning in human memory. *Science*, 1966, 153, 652-654.
- STONE, G. C., & CALLOWAY, E. Effects of stimulus probability on RT in a number naming task. *Quarterly Journal of Experimental Psychology*, 1964, 16, 47-55.
- WINER, B. J. *Statistical principles of experimental design*. New York: Holt, Rinehart & Winston, 1955.

NOTE

1. For each combination of S, stimulus type (set), and week, the harmonic mean RTs for S2 and S3 were averaged in Condition 1, where they occurred with equal frequency, before analyses of variance were applied.

(Accepted for publication April 12, 1972.)