Table 1

	ECP		EEG		
	P ₂	n ₂	Alpha	Beta	Amplitude
Simple Reaction Time	.14	.45	.06	.05	.03
Visual Discrimination	—	.09	—	—	.02

The late components of the ECP, in particular n₂, were also found to be more strongly related to level of performance than any of the EEG measures. It should be noted that combined frequency and amplitude EEG measures did not substantially raise (no more than two percentage points) the strength of statistical association values reported above. Moreover, response-produced effects on the ECP are an unlikely reason for the superiority of the ECP in predicting performance. Ss whose reaction time scores exceeded the completion of the ECP showed the same relationship between the ECP and reaction time as those Ss whose responses overlapped with the occurrence of the ECP. Secondly, one would expect the latency of the various components of the ECP to be modified by response-produced effects, since such effects would tend to prolong the particular phase of the potential with which they overlapped. As noted above, however, latency scores were constant under all conditions of the experiment. Thirdly, response-produced effects should have influenced the waveform of the ECP in those cases in which S's response overlapped with the occurrence of the potential. However, the idiosyncratic fluctuations of the ECP characteristic of a given S were found under all conditions of the experiment and occurred invariably with the same latency.

Finally, the ability of the EEG and ECP to predict performance appears to be restricted to relatively simple tasks, in particular to those tasks which do not require decision processes. This finding is not altogether surprising, since nonsensory factors are known to have profound effects upon discrimination performance (Swets, Tanner, & Birdsall, 1964). It does, however, suggest that arousal factors may play a limited role in the control of complex decision processes.

REFERENCES

- DONCHIN, E., & LINDSLEY, D. B. Average evoked potentials and reaction times to visual stimuli. *Electroencephalography & Clinical Neurophysiology*, 1966, 20, 217-223.
- DUSTMAN, R. E., BOSWELL, R. S., & PORTER, P. B. Beta brain waves as an index of alertness. *Science*, 1962, 137, 533-534.
- FUSTER, J. M. Effects of stimulation of brain stem on tachistoscopic perception. *Science*, 1958, 127, 150.
- HAIDER, M., SPONG, P., & LINDSLEY, D. B. Attention, vigilance, and cortical

- evoked potentials in humans. *Science*, 1964, 145, 180-182.
- HAYS, W. L. *Statistics for psychologists*. New York: Holt, Rinehart, & Winston, 1963.
- LINDSLEY, D. B. Emotion. In S. S. Stevens (Ed.), *Handbook of experimental psychology*. New York: Wiley, 1951. Pp. 473-516.
- LINDSLEY, D. B. Psychological phenomena and the electroencephalogram. *Electroencephalography & Clinical Neurophysiology*, 1952, 4, 443-456.

- MORRELL, L. K., & MORRELL, F. Evoked potentials and reaction times: A study of intra-individual variability. *Electroencephalography & Clinical Neurophysiology*, 1966, 20, 567-575.
- SWETS, J. A., TANNER, W. P., & BIRDSALL, T. G. Decision processes in perception. In J. A. Swets (Ed.), *Signal detection and recognition by human observers*. New York: Wiley, 1964. Pp. 3-57.

NOTE

1. This report is based upon a dissertation submitted in partial fulfillment of the requirements for the PhD degree, University of Missouri, 1967. Part of the research was supported by a NASA grant to Robert S. Daniel, to whom the author is indebted for his guidance during the course of the investigation and during his entire graduate career.

Multiple stimulus discrimination learning, II. Effects of prior training*

JEAN M. MANDLER

University of California at San Diego, La Jolla, Calif. 92037

Performance on multiple-stimulus visual discriminations improved as a function of amount of prior training on another visual discrimination. Performance on the multiple-stimulus tasks, however, indicated no changes in positive and negative stimulus control as prior training was increased from criterion performance to include overtraining. It was concluded that the greater S+ control following overtraining found in previous data may not be generalized stimulus control but specific to the stimuli used in acquisition.

Following overtraining on a two-choice visual discrimination, Mandler (1968) found that changing the positive stimulus (S+) disrupted subsequent performance more than changing the negative stimulus (S-). Following mastery training only, there was a small but nonsignificantly greater disruption when S- was changed. It was concluded that one of the effects of overtraining was an increase in control by S+ and a decrease in control by S- over choice behavior. Since reacquisition following overtraining proceeds very rapidly, regardless of which stimulus is changed, it seems likely that real differences in positive and negative stimulus control might be partially masked by ceiling effects. Furthermore, when the testing procedure involves the same stimuli used in acquisition, it is not possible to determine whether stimulus control is specific to the stimuli used in acquisition or whether it is more general in nature. It was decided, therefore, to use a transfer test which would produce more errors as well as involve new visual stimuli, yet which

would still make it possible to detect differences in positive and negative stimulus control.

One way to test the relative control of S+ and S- is to vary the number of positive and negative stimuli used in a given problem. Mandler (1970a) found that a discrimination task in which there was one S- and three positive stimuli was more difficult than a standard S+S- task but easier than a task with one S+ and three negative stimuli. This type of discrimination problem weights the relative importance of S+ and S- in acquisition and should be sensitive to biases in positive and negative stimulus control established by prior training. It was decided to study acquisition of such multiple-stimulus discriminations following mastery or overtraining on a standard S+S- discrimination. If stimulus control is changing during the course of original acquisition from relatively equal control at mastery to greater control by S+ following overtraining, these differences should affect the acquisition of a task in which S+ and S- are differentially weighted. If S+ is controlling choice behavior more than S-, a task with constant S+/multiple S- should be easier than a task with a multiple S+/constant S-. The results on the multiple task should be the opposite if

*This work was supported in part by Research Grant MH 16850 from the National Institute of Mental Health.

S- is playing a major role in controlling choice. Performance on the multiple tasks should also indicate whether differences in S+ and S- control are stimulus specific, e.g., apply only to the actual stimuli used in training, or whether they are of a more general nature, involving behavior toward new visual stimuli functioning as S+ and S- in a discrimination.

Finally, the experimental design enables us to find out if overtraining of a visual discrimination facilitates acquisition of a multiple stimulus task in the same way it facilitates a standard S+S- visual task (Mandler, 1968).

METHOD

The Ss were 48 male hooded rats between 80 and 90 days old at the start of the experiment. The apparatus was a semiautomated Y-maze with a 24-in. stem. The goalbox doors, on which $3\frac{1}{2} \times 2\frac{1}{2}$ in. stimulus cards were placed, were 12 in. from the choice point of the Y. Goal- and startboxes were unpainted metal.

Three groups of 16 Ss each were used: a control group which received no prior training before learning the multiple-stimulus discrimination; a mastery group which was trained to a criterion of mastery on a black-white discrimination before learning the multiple task; and an overtrained group which received 150 trials past criterion on the black-white discrimination before training on the multiple task. Ten correction trials a day were given, with an intertrial interval of approximately 3 min. Criterion consisted of 2 consecutive days of 90% correct responding or better. Ss were run under 23-h food deprivation and were allowed to eat for 1 h after the day's run. Reinforcement consisted of one 250-mg pellet per trial.

On Task 1, half of the Ss in the mastery and overtrained groups were assigned randomly to black positive (B+) and half to white positive (W+). B and W stimuli were assigned to the left and right arms of the maze, following Gellerman orders. On Task 2, half of the animals in each group were trained on a task with a single S+ and multiple (M), i.e., three, negative stimuli (S+M-); for the other half, the task involved a single S- and three positive stimuli (S-M+). Ss were assigned randomly to Task 2, except for insuring equal numbers of B+ and W+ Ss in each condition. In Task 2, the same constant stimulus (either S+ or S-) was presented on each trial with one of the three multiple stimuli. The order of presentation of the multiple stimuli was random, with the proviso that all three should appear at least three trials per day and no one of

Table 1
Trials to Criterion on Multiple
Stimulus Discriminations

Prior Black- White Training	S+M-	S-M+
None	161.2	123.8
Mastery	101.2	96.2
Overtraining	61.2	57.5

them should appear more than two trials in a row. Control Ss were assigned randomly to one of the two multiple tasks and had no prior training on black-white.

Stimulus cards for the multiple task were constructed from equal amounts of black and white in one of the following designs: $\frac{1}{2}$ -in.-wide horizontal stripes; $\frac{1}{2}$ -in.-wide vertical stripes; $3/8$ -in.-wide diagonal stripes; and a black right triangle on a white ground, oriented in the opposite direction from the diagonal stimulus. The constant stimulus was always either horizontal or vertical stripes, randomly assigned to the Ss, but counterbalancing for previous experience with B+ or W+. When horizontal stripes was the constant stimulus, vertical stripes, diagonal stripes, and triangle were the multiple stimuli. When vertical was the constant stimulus, horizontal, diagonal, and triangle were the multiple stimuli. Thus, in the mastery and overtrained groups, Ss were transferred from a black-white task to one having either horizontal or vertical stripes as its constant stimulus. Previous experiments using this apparatus have indicated no direct transfer biases from black or white to either horizontal or vertical (Mandler, 1968).

RESULTS AND DISCUSSION

There were no significant differences in trials to criterion on Task 1 for the mastery and overtrained groups. Trials to criterion for the multiple task are shown in Table 1. It can be seen that the chief effect of increasing prior amount of black-white training was increased facilitation in acquisition of the multiple task. Analysis of variance on trials to criterion showed amount of prior training to be a significant variable, $F(2,42) = 14.75$. Facilitative effects of prior visual training on another visual task have been shown before (Mandler, 1968). However, prior training on a nonvisual discrimination does not facilitate later visual discrimination learning (Mandler, 1966). In the present experiment, facilitation took place even though the multiple-stimulus task has somewhat different requirements than the usual S+S- visual task; either the Ss must compare the constant stimulus with several variable stimuli or else they

learn the discrimination on the basis of the constant stimulus alone. The present results, then, suggest that the facilitation of later learning, due to overtraining, stems from increased ability to handle visual stimuli and is, perhaps, attentional in nature.

Turning to the question of positive and negative stimulus control, the only significant differences in acquisition of the multiple positive vs the multiple negative discrimination occur in the control groups, namely, those which had had no prior black-white training. The constant S- multiple S+ task was easier to learn than the constant S+ multiple S- task [$t(42) = 1.75$, $p < .05$, one-tailed test]. There were no significant differences in ease of learning the two kinds of multiple tasks following either mastery training or overtraining. Recent work from our laboratory has shown that multiple stimulus tasks are learned on the basis of the constant stimulus alone, whether S+ or S-, and that Ss do not discriminate the multiple stimuli in the task (Mandler, 1970b). The present results, then, in demonstrating that overtrained Ss learn a discrimination weighted toward S- as quickly as one weighted toward S+, suggest that the differences in positive and negative stimulus control following overtraining are stimulus specific. The previous data showing greater S+ control than S- control following overtraining (Mandler, 1968) involved a testing situation in which one of the acquisition stimuli was retained. In the present experiment new stimuli were used in the test situation, and no differences in S+ and S- control were found.

Although Ss with no prior training learned the multiple task with constant S- faster than one with constant S+ and this difference disappeared following discrimination training, it seems premature to conclude that S- control is greater in the early stages of discrimination learning. Further work will be needed on various stages of acquisition to determine exact changes in positive and negative stimulus control and the degree of their stimulus specificity.

REFERENCES

- MANDLER, J. M. Behavior changes during overtraining and their effects on reversal and transfer. *Psychonomic Monograph Supplements*, 1966, 1, 187-202.
- MANDLER, J. M. Overtraining and the use of positive and negative stimuli in reversal and transfer. *Journal of Comparative & Physiological Psychology*, 1968, 66, 110-115.
- MANDLER, J. M. Two-choice discrimination learning using multiple stimuli. *Learning & Motivation*, 1970a, 1, 261-266.
- MANDLER, J. M. What is learned in multiple stimulus discriminations. Paper presented at Psychonomic Society Meetings, 1970b.